

2003

# Temporally dependent C, N, and P dynamics associated with the decay of *Rhizophora mangle* L. leaf litter in oligotrophic mangrove wetlands of the Southern Everglades

Stephen E. Davis III

*Southeast Environmental Research Center, Florida International University*

Carlos Coronado-Molina

*Department of Oceanography & Coastal Sciences, Louisiana State University*

Daniel L. Childers

*Southeast Environmental Research Center, Florida International University*

John W. Day Jr.

*Department of Oceanography & Coastal Sciences, Louisiana State University*

Follow this and additional works at: [http://digitalcommons.fiu.edu/fce\\_lter\\_journal\\_articles](http://digitalcommons.fiu.edu/fce_lter_journal_articles)

---

## Recommended Citation

Davis, S.E., C. Coronado-Molina, D.L. Childers, J.W. Day. 2003. Temporally dependent C, N, and P dynamics associated with the decay of *Rhizophora mangle* L. leaf litter in oligotrophic mangrove wetlands of the Southern Everglades. *Aquatic Botany* 75(3): 199-215.

This material is based upon work supported by the National Science Foundation through the Florida Coastal Everglades Long-Term Ecological Research program under Cooperative Agreements #DBI-0620409 and #DEB-9910514. Any opinions, findings, conclusions, or recommendations expressed in the material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

This work is brought to you for free and open access by the FCE LTER at FIU Digital Commons. It has been accepted for inclusion in FCE LTER Journal Articles by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu), [jkrefft@fiu.edu](mailto:jkrefft@fiu.edu).

1 Temporally dependent C, N, and P dynamics associated with the decay of  
2 Rhizophora mangle L. leaf litter in oligotrophic mangrove wetlands of the  
3 Southern Everglades

4

5 Stephen E. Davis, III<sup>ac</sup>, Carlos Coronado-Molina<sup>bd</sup>, Daniel L. Childers<sup>a</sup>, and John W. Day, Jr.<sup>b</sup>

6

7 <sup>a</sup> Southeast Environmental Research Center, Florida International University, University Park Campus,  
8 Miami, Florida 33199, U.S.A.

9

10 <sup>b</sup> Dept. of Oceanography & Coastal Sciences, Louisiana State University, Baton Rouge, Louisiana 70803, U.S.A.

11

12 <sup>c</sup> Author for correspondence. Current address: Dept. of Wildlife and Fisheries Sciences, 210 Nagle Hall, Texas  
13 A&M University, College Station, Texas 77843, U.S.A.

14

15 <sup>d</sup> Current address: Everglades Division, South Florida Water Management District, 3301 Gun Club Road,  
16 West Palm Beach, Florida 33416, U.S.A.

1 **Abstract**

2 We performed two litter decomposition experiments using nearly-senesced red mangrove (Rhizophora  
3 mangle L.) leaves collected from an Everglades dwarf mangrove wetland to understand the short-term (3 weeks) and  
4 long-term (1 year) changes in mass, as well as carbon, nitrogen, and phosphorus contents of decomposing leaf litter.  
5 We expected that leaves decomposing in this oligotrophic environment would be short-term sources of C, N, and P,  
6 but potential long-term sinks for N and P. In May 1998, we conducted a 3-week leaching experiment, incubating  
7 fresh, individual leaves in seawater for up to 21 days. From May 1997 to May 1998, leaf litter in mesh bags  
8 decomposed on the forest floor at two dwarf mangrove sites. Leaching accounted for about 33% loss of dry mass  
9 from R. mangle leaves after 3 weeks. Leaching losses were rapid, peaking by Day 2, and large, with leachate  
10 concentrations of total organic carbon (TOC) and total phosphorus (TP) increasing by more than an order of  
11 magnitude after 3 weeks. Mean leaf C:N increased from 105 to 115 and N:P increased from a mean of 74 to 95 after  
12 21 days, reflecting the relatively large leaching losses of N and P. Loss of mass in the litterbags leveled off after 4  
13 months, with roughly 60% dry mass remaining after nearly one year of decomposition. The mass of carbon in each  
14 litterbag declined significantly after 361 days, but the mass of nitrogen and phosphorus doubled, indicating long-  
15 term accumulation of these constituents into the detritus. Subsequently, the leaf C:N ratio dropped significantly  
16 from 90 to 34 after 361 days. Following an initial 44-day increase, leaf N:P decreased from 222 to 144, reflecting  
17 high accumulation of P relative to N. A review of several estuarine macrophyte decomposition studies reveals a  
18 trend in nitrogen accumulation through time regardless of site, but suggests no clear pattern for C and P. We believe  
19 that the increase in litter P observed in this study was indicative of the P-limited status of the greater Everglades  
20 ecosystem and that decomposing mangrove litter may represent a substantial phosphorus pool in the system.

21  
22 Keywords: leaves; leaching; decomposition; carbon; nitrogen; phosphorus

## 1 **1. Introduction**

2 Litterfall from deciduous and evergreen trees is a primary mechanism by which nutrients are returned to the  
3 forest floor. Accounting for roughly 70% of the dry mass of all aboveground litter in forested ecosystems, leaves  
4 are usually the most important litter component (O'Neill and DeAngelis, 1981). In estuarine mangrove forests, leaf  
5 litter can account for 40-95% of the total pool of litterfall (e.g. Day et al., 1996; Wafar et al., 1997). This fraction of  
6 mangrove litter represents a relatively large, potentially labile source of organic matter to decomposer communities  
7 (Benner and Hodson, 1985; Benner et al., 1986).

8 Resorption of materials prior to leaf abscission can be an effective means of conserving vital elements in  
9 many mangrove species (Untawale et al., 1977; Karmarkar, 1982; Lin and Wang, 2001). However, there is still a  
10 substantial turnover of organic and inorganic materials in the system via leaf litter decomposition (Twilley et al.,  
11 1986a; Moran et al., 1991; Wafar et al., 1997). These materials can be an important source of energy and nutrition  
12 to heterotrophic communities, especially in oligotrophic wetland systems such as the Everglades.

13 The decomposition of plants typically occurs in three, often simultaneous phases: 1) leaching of soluble  
14 components, 2) microbial oxidation of refractory components such as cellulose and lignin, and 3) physical and  
15 biological fragmentation (Valiela et al. 1985). The leaching phase of mangrove leaf decomposition, as with most  
16 other litter types, is characterized by a rapid loss of soluble organic compounds (sugars, organic acids, proteins,  
17 phenolics, etc.) and inorganic minerals (K, Ca, Mg, Mn, etc.). Regardless of vegetation type, this phase lasts  
18 anywhere from a few days to a few weeks and can be responsible for substantial losses of mass, carbon, nitrogen,  
19 and phosphorus (Valiela et al., 1985; Parsons et al., 1990; Chale, 1993; Steinke et al., 1993; Taylor and Barlocher,  
20 1996; France et al., 1997). The biotic contributions in this early stage of decomposition are usually minimal and are  
21 most often limited to microbial conditioning of the litter (Nykvist, 1959; Cundell et al., 1979; France et al., 1997).

22 The second and third phases of leaf litter decomposition are characterized by microbially mediated  
23 breakdown of labile organic material and refractory structural components, all of which is enhanced by physical and  
24 biological fragmentation of the litter (Harrison and Mann, 1975). Since refractory compounds make up the bulk of  
25 leaf mass, these latter stages of decay operate over longer time scales than leaching, resulting in a gradual loss of  
26 reduced carbon over time. However, relative to carbon, there is often a considerable amount of nutrient  
27 accumulation associated with increasing microbial biomass in the detritus complex. Studies on the decay of a  
28 variety of estuarine macrophytes have shown significant accumulation of N and P relative to carbon on temporal

1 scales of a few weeks to several months (Rice and Tenore, 1981; Day et al., 1982; Twilley et al., 1986a; Twilley et  
2 al., 1986b).

3           There are a number of studies that have focused on the decomposition of estuarine macrophytes (e.g.  
4 Harrison and Mann, 1975; White et al., 1978; Rice and Tenore, 1981; Tam et al., 1990). Many of these have  
5 addressed temporal changes in the nutritional content of the litter. However, only a few have considered the time-  
6 dependence of such dynamics by focusing on both the rapid (hours—days) decay component and the long-term  
7 (weeks—months), more refractory components of decomposition (Twilley et al., 1986a; Chale, 1993). In the  
8 present study, we report the findings of simultaneous experiments that addressed both the short-term (3 week) and  
9 long-term (one year) dynamics of carbon, nitrogen, and phosphorus associated with the decay of leaves from a dwarf  
10 red mangrove (Rhizophora mangle L.) forest in the oligotrophic Southern Everglades.

11           As others have previously shown, we anticipated high leaching losses of carbon, nitrogen, and phosphorus  
12 from the leaves over the first few weeks of decomposition. We also expected that the low availability of N and P in  
13 this oligotrophic wetland would result in the long-term accumulation of these nutrients into the detritus matrix, as  
14 has been shown for nitrogen in many other systems. Finally, we compared our results with data synthesized from  
15 numerous estuarine macrophyte decomposition studies to reveal any temporal trends in litter C, N, and P dynamics.  
16 Given the trophic state of the Everglades and concerns about anthropogenic inputs of nutrients to the Everglades and  
17 Florida Bay, there is a need to understand the importance of mangrove leaf detritus as temporally dependent sources  
18 or sinks of nitrogen and phosphorus.

19

## 20 **2. Site Description**

21           The Southern Everglades mangrove zone is part of a highly oligotrophic estuarine system. Roughly 6,000  
22 ha of this area are composed primarily of the dwarf form (1-2 m canopy height) of the red mangrove (Rhizophora  
23 mangle L.; Lin and Sternberg, 1992). Leaf litter comprises about 87% ± 4% of total litterfall in this area and  
24 productivity and standing crop values are among the lowest recorded for any mangrove forest (Corronado-Molina,  
25 2000). Phosphate concentrations in the surface water and soil pore water are low and often below the limit of  
26 detection (0.01 µM), reflecting the P-limited nature of this region (Koch, 1997; Koch and Snedaker, 1997; Davis et  
27 al., 2001). Rates of herbivory on dwarf red mangrove vegetation are also low, as herbivory has been linked to

1 nutrient status (Onuf et al., 1977; Feller, 1995). As a result, much of the annual litter production (and associated  
2 nutrients) is made available to higher trophic levels only through detrital pathways.

3 The hydrology of the Southern Everglades mangrove zone is micro-tidal (< 5 cm tidal range) and seasonal  
4 in rainfall and discharge (wet and dry seasons). Therefore, the direction and velocity of flow and salinity patterns in  
5 mangrove creeks such as Taylor River are driven by the interaction of precipitation, upland runoff, and wind, the  
6 combination of which leads to a net southerly flow and oligohaline/fresh conditions throughout much of the wet  
7 season (usually from June – November) and no net flow and mesohaline/polyhaline conditions throughout the dry  
8 season (Sutula, 1999). Furthermore, dwarf mangrove wetlands in this area are characterized by persistent standing  
9 water throughout most of the year, and the soils are completely exposed to the atmosphere only during an extended  
10 drought (S. Davis, pers. observ.).

### 11 12 **3. Materials and Methods**

13 For both experiments, we collected nearly-senesced, yellow leaves from dwarf red mangrove trees along  
14 the lower Taylor River in the southern portion of Everglades National Park (Figure 1). We report the carbon,  
15 nitrogen, and phosphorus content of leaf material from the leaching and litterbag studies as relative (% or mass of  
16 constituent g dry weight<sup>-1</sup> of tissue) or absolute (mass of constituent per leaf or litterbag). Single-factor analyses of  
17 variance were used to determine a significant ( $p < 0.05$ ) change in leaf mass, C, N, or P content of leachate or leaves  
18 over time. Tukey tests were used to determine significant ( $p < 0.05$ ) differences among treatment means.

#### 19 20 3.1 Leaching Study

21 In May 1998, we incubated individual, fresh leaves in 250 ml, clear, glass bottles containing filtered (GF/F)  
22 seawater (32 ppt) for up to 21 days. The experimental leaves were not pre-dried, as drying has been shown to  
23 modify rates of leaching in many species (Taylor and Barlocher, 1997; Taylor, 1998). Bottles were incubated in the  
24 field under ambient temperature and sunlight conditions. In order to determine the abiotic contributions (leaching)  
25 to mass and nutrient loss from each individual leaf, we added 2 ml of a 1%  $\text{NaN}_3$  solution to half of the experimental  
26 units as an inhibitor of aerobic respiration. We collected 3 replicate bottles of each treatment after 1, 2, 5, 10, and 21  
27 days of decomposition. Following incubation, leaves were removed from the bottles and rinsed with de-ionized  
28 water and dried to a constant weight at 60 °C (final dry mass =  $\text{DM}_t$ ).

1            Since we chose to use fresh material, an accurate means of estimating initial dry weight was needed in  
2 order to determine change in mass over the period of decay. We used a batch of initial control leaves (n=75) to  
3 develop a simple linear regression model that could be used to estimate initial dry weight ( $DM_0$ ) of each  
4 experimental leaf from its initial fresh mass. This model—with no y-intercept—indicated that dry mass was  
5 consistently 34% of initial fresh mass ( $p < 0.0001$ ; adjusted  $r^2 = 0.99$ ).

6            Water samples collected from each bottle were stored at 4° C until analyzed for total phosphorus (TP)—  
7 according to a modification of the dry ashing, acid-hydrolysis technique of Solorzano and Sharp (1980)—total  
8 nitrogen (TN)—using an Antec 7000N total nitrogen analyzer—and total organic carbon (TOC)—using a hot  
9 platinum catalyst, direct injection analyzer; Shimadzu model TOC-5000. After final dry weight measurements were  
10 taken, all leaves were ground to a fine powder and analyzed for carbon and nitrogen content—using a Carlo Erba  
11 1500-N CHN analyzer—and TP content, as described above.

12            At the conclusion of the 21-day experiment, we calculated percent dry mass remaining (% DMR) for each  
13 leaf by dividing  $DM_t$  by  $DM_0$ . To ensure that changes in water nutrients were due to the presence of the leaves, we  
14 also incubated "control" bottles containing only filtered seawater or seawater +  $NaN_3$  for the entire 21-day length of  
15 the leaching experiment. We compared TOC, TN, and TP concentrations from these control bottles with initial  
16 values to determine changes associated with water column or photochemical processes. Paired t-tests were used to  
17 determine significant differences between initial and final concentrations ( $\alpha = 0.05$ ). Leaching rates (i.e. fluxes) of  
18 TOC, TN, and TP were calculated from concentrations in the water as moles g DW leaf<sup>-1</sup> and averaged over the  
19 number of days between each sampling interval.

### 21 3.2 Litterbag Study

22            We dried the leaves for the litterbag study at 60 °C for 48 hours and partitioned the pooled mass into  
23 replicates, each weighing approximately 5 g. We then placed samples into mesh bags fashioned from fiberglass  
24 window screen (approx. 1-2 mm mesh size). Sixteen litterbags were distributed on the forest floor at each of two  
25 sites (Sites A and B) each approximately 5 m from creek channel in the dwarf mangrove forest (Figure 1). These  
26 two sites were randomly selected to account for some of the variability in the dwarf mangrove forest along lower  
27 Taylor River. The litterbag experiment lasted from May 1997 to May 1998. At the beginning of the experiment

1 (time = 0), two bags were retained to determine relative and absolute C, N, and P content. Thereafter, duplicate  
2 samples were retrieved from each site after 13, 31, 44, 87, 128, 184, 304, and 361 days of decomposition in the field.

3 After collection, litterbags were immediately taken to the laboratory and prepared for analysis. Leaves  
4 were removed from each bag, gently washed to remove sediment and then oven-dried to constant weight at 60 °C.  
5 We calculated % DMR by dividing  $DM_t$  by  $DM_0$  and multiplying by 100. A sub-sample of dried tissue from each  
6 litterbag was ground to a fine powder and analyzed for C, N, and P content using the same procedures as described  
7 for the leaching experiment.

## 8 9 **4. Results**

### 10 4.1 Leaching study

11 There was no significant inhibitor effect on % DMR over the 21-day leaching experiment (Figure 2a), and  
12 all leaves lost a significant amount of mass (ANOVA,  $p < 0.0001$ ). After 24 hours of decomposition in the bottles,  
13 red mangrove leaves had lost  $17\% \pm 2\%$  (std dev) of the initial dry mass. The majority of these losses were  
14 attributable to leaching rather than microbially mediated processes, as aerobic respiration was inhibited with sodium  
15 azide in half of the experimental units (Figure 2a). For the next 9 days, all leaves lost approximately the same  
16 amount of dry mass per day ( $1.3\% d^{-1}$ ). However, from day 10 to day 21, leaching losses seemed to subside while  
17 microbial activity (bottles without  $NaN_3$ ) appeared to increase, as the difference between poisoned and non-poisoned  
18 bottles became more noticeable (Figure 2a). At the conclusion of the experiment, poisoned and non-poisoned leaves  
19 had lost  $33\% \pm 6\%$  and  $41\% \pm 17\%$  of their initial dry mass, respectively, but these did not differ significantly.

20 The C, N, and P content of the leaves in the leaching experiment showed little inhibitor effect (poison vs.  
21 no poison) over the 3-week experiment. In addition, we were unable to detect any statistically significant change in  
22 relative and absolute C, N, and P content of the leaves over time. Nevertheless, mean relative C of leaves increased  
23 slightly after three weeks (Figure 2b), but absolute values decreased noticeably from Day 0 to Day 21 (Figure 2c),  
24 reflecting the large losses of mass. Similarly, the mean relative N content of all leaves remained fairly steady at  
25 about 0.6% (Figure 2b), but they showed a decrease in the absolute N content after three weeks (Figure 2c). Mean  
26 % P content of leaves also declined—especially over the first few days in the presence of azide (Figure 2b). Initial  
27 values of relative P ( $0.021 \pm 0.001\%$ ) were nearly double those after 3 weeks ( $0.011 \pm 0.001\%$ ). As with C and N,  
28 absolute P content decreased throughout the 3-week study period (Figure 2c).



1 We found no statistical difference in C:N ratios between poisoned and non-poisoned leaves after 21 days of  
2 immersion. After 24 hours, leaves exhibited a decline in C:N from 105 to an overall mean of 96 (Figure 3).  
3 Thereafter, the C:N of poisoned leaves increased above initial levels, fluctuating between a mean of 110 and 115  
4 throughout the remainder of the experiment (Figure 3). The C:N of the non-poisoned leaves showed an overall  
5 decline over the first 10 days, and then increased from a mean of 82 to 105 after 21 days (Figure 3). The ratios of  
6 N:P in these same leaves showed overall increases from  $74 \pm 8$  (std dev) at Day 0 to Day 21 levels of  $95 \pm 18$  (non-  
7 poisoned) and  $104 \pm 10$  (poisoned; Figure 3). The only significant difference in N:P between treatment levels was at  
8 Day 1, when N:P of the non-poisoned leaves decreased below initial levels to  $53 \pm 10$  while poisoned leaves  
9 remained unchanged at  $76 \pm 7$  (ANOVA;  $p=0.03$ ). From Day 1 to Day 2, the N:P of non-poisoned leaves nearly  
10 doubled to a mean of 94 (Figure 3).

11 We observed substantial changes in water chemistry throughout the leaching study. Bottles with leaves  
12 showed significant increases in TOC, TN, and TP over the 3-week study (Table 1). In either treatment, mean TOC  
13 and TP concentrations increased by more than an order of magnitude after 21 days, and average concentrations of  
14 TN more than doubled in the "no azide" treatment (Table 1). Given the high N-content of our poison ( $\text{NaN}_3$ )  
15 relative to ambient water concentrations, we did not analyze poisoned samples for TN content. Fluxes for all  
16 constituents were generally highest after 1 or 2 days, then declined considerably over the next 19 to 20 days of  
17 incubation (Table 1). We found no significant change in the TOC, TN, or TP content of the water in the control  
18 bottles without leaves.

#### 19 20 4.2 Litterbag study

21 We observed a significant decrease in the % DMR over time using the litterbag technique (ANOVA;  
22  $p < 0.0001$ ; Figure 4a). Although there appeared to be a difference in the loss of dry mass between sites after 128  
23 days of decay, there was no significant difference in % DMR between sites (ANOVA;  $p > 0.05$ ; Figure 4a). The  
24 percentage of dry mass loss in individual litterbags was quite variable after 13 days of decomposition in the field,  
25 ranging from 2% to 35%. High losses of mass continued for about 3 months at Site A and 4 months at Site B, but  
26 then appeared to level off (Figure 4a). At the conclusion of the litterbag experiment, leaves decomposing at both  
27 sites had lost an average of  $40 \pm 7\%$  of their initial dry mass.

1           The C, N, and P contents of the litterbag leaves were consistent between sites. The relative carbon content  
2 of litterbag leaves increased from about 44% to as much as 55% over the first 3 months of the study, but then  
3 declined, approaching initial levels by the end of the study (Figure 4b). Normalized to dry mass of tissue, this  
4 pattern was not evident, as the absolute concentration of carbon decreased over the course of the entire experiment  
5 (Figure 4c). Patterns for nitrogen and phosphorus in these leaves were comparable with significant increases in the  
6 relative and absolute content occurring over the entire year of this study (Figures 4b and 4c). Mean relative nitrogen  
7 increased from 0.6% to 1.7% and absolute values nearly doubled (0.03 to 0.05 g). Long-term increases in  
8 phosphorus were somewhat similar, with relative P increasing from a mean of 0.009% to 0.029% and absolute P  
9 increasing from an initial average of 0.4 mg to 0.9 after 1 year (Figures 4b and 4c).

10           Molar ratios of C:N and N:P in leaf tissue were also quite similar in the two sites. Leaf C:N decreased  
11 significantly from a mean of 90 to 34 after 361 days in the field, reflecting the losses in absolute C and increases in  
12 absolute N (Figure 5). The pattern for leaf N:P was different, with large increases up through Day 44 followed by  
13 decreases to near initial levels by Day 128 (mean=144; Figure 5). At its highest, the N:P ratio of our mangrove litter  
14 averaged  $222 \pm 41$ .

## 16 **5. Discussion**

17           Leaching has been shown to be a major pathway for loss of mass and materials from senesced estuarine  
18 vegetation. In our bottle experiment, leaching accounted for more than a third of the mass lost after three weeks,  
19 and short-term leaching losses were nearly as great as the mass lost from leaves after one year in the litterbag study  
20 (Figures 2 & 4). This was remarkable given that past studies using the litterbag methodology have shown a 60%–  
21 90% loss of mangrove leaf mass in 6 months or less (Steinke et al., 1983; Twilley et al., 1986a; Tam et al., 1990;  
22 Mackey and Smail, 1996). As anticipated, leaching losses translated to substantial fluxes of TOC, TN, and TP to the  
23 water column, with highest flux rates occurring during the first few days of leaf submergence (Table 1).

24           We found no inhibitor effects on total fluxes of C, N, and P or on litter C, N, and P content throughout the  
25 study. This supported the notion that leaching was the principal process contributing to materials loss during the  
26 early phase of R. mangle leaf litter decay. After 361 days, the relative concentration of C in the litter was no  
27 different than % C in the leaching leaves. However, relative concentrations of N were significantly higher after  
28 long-term decomposition in the field ( $1.6 \pm 0.09$ ) than after 21 days of leaching ( $0.53 \pm 0.08$ ). In addition, relative P

1 was higher at the conclusion of the litterbag study ( $0.028 \pm 0.003$  vs.  $0.011 \pm 0.002$ ) even though initial % P of the  
2 leaching leaves was double the initial content of the litterbag leaves (Figures 2 and 4). This N and P enrichment  
3 over time signifies the microbial contribution to the long-term control of red mangrove leaf litter quality.

4 The early decay of estuarine macrophytes often varies as a result of inundation period, with highest losses  
5 generally occurring in low inter-tidal zones (Kruczynski et al., 1978; Valiela et al., 1985; Twilley et al, 1986a;  
6 Mackey and Smail; 1996). Our short-term losses are similar to those (25% after 14 days) reported by Steinke et al.  
7 (1993) in their study of Avicennia marina leaves. Other work on A. marina indicated a 19% loss of mass after 24  
8 hours of submergence and only an additional 11% loss after 6 weeks (Chale, 1993). Robertson (1988) determined  
9 that two species of mangrove leaves decomposed much more slowly on the exposed forest floor than when  
10 continuously submerged. In our leaching study, leaves were continually submerged for up to 3 weeks, perhaps  
11 enhancing the rate of loss. The inundation pattern at our litterbag sites was much more variable. Although we did  
12 not intensively monitor site inundation, we estimated that the litterbags at both dwarf mangrove sites were inundated  
13 at least 50% of the year.

14 The carbon content of our leaves showed little change over the first few weeks of decay. Given that the  
15 bulk of the carbon in these leaves is tied up in refractory structural tissues, this was not unexpected. Nevertheless,  
16 many estuarine macrophytes, including red mangrove leaves, are known to leach considerable amounts of labile  
17 DOC to the water column when submerged (Rice and Tenore, 1981; Benner et al., 1986; Twilley et al., 1986a; this  
18 study).

19 Early changes in N seem to be variable and may be related to soluble protein content in the leaf tissue  
20 (Table 2). Some have suggested that changes in leaf N are function of initial C:N (Harrison and Mann, 1975; Fell et  
21 al., 1984). However, initial C:N ratios alone do not seem to explain the differences we observed in early N  
22 dynamics. Cundell et al. (1979) and Fell et al. (1975) measured large, early increases in % N of submerged R.  
23 mangle leaves with initial C:N ranging from 90-110 (Table 2). We found a small, early decrease in leaf % N with  
24 initial C:N also ranging from 90-110 (Table 2). The difference, as described by Twilley et al. (1986b), may be  
25 controlled by the amount of N available to microbial decomposers. Perhaps the greater availability of N (relative to  
26 P) at our Southern Everglades sites accounted for the differences we observed in R. mangle leaf decomposition.

27 Early P dynamics in decomposing estuarine macrophytes appears more consistent, as a preponderance of  
28 the cases we reviewed showed substantial decreases in relative P in 3 weeks or less (Table 2). In this study, we

1 found a significant increase in the TP content of the leachate in the bottles that coincided with an apparent decrease  
2 in the mean % P content of the leaves. High P leaching from estuarine macrophytes is believed to be a result of the  
3 large inorganic fraction of P in leaf tissue, as opposed to N, which is mostly organic (Twilley et al., 1986b). Data in  
4 Table 2 suggest that estuarine macrophyte litter can be a rapid, short-term source of N and P to the environment,  
5 possibly enhancing the degradability and microbial conditioning of the plant litter. This may be particularly  
6 important to the degradation of litter in oligotrophic environments such as the dwarf mangrove wetlands of the  
7 Southern Everglades.

8         The long-term decay of the more refractory components of estuarine macrophyte litter is not as influenced  
9 by the inundation pattern of the site (Valiela et al., 1985). We observed a continuous loss of mass from our litterbag  
10 leaves until about Day 128, regardless of inundation pattern. From this point on, values stabilized at about 55-65%  
11 DMR for the remaining 200+ days of the study (Figure 2), suggesting that about half of the mass of these dwarf  
12 mangrove leaves is refractory. Johnson et al. (1993) demonstrated a similar trend for blades of S. patens in a  
13 mesohaline Louisiana coastal marsh with about 56% DMR after 320 days. Studies have shown that the breakdown  
14 of this recalcitrant fraction is more dependent on habitat characteristics such as the decomposer community, tidal  
15 energy, site fertility, tissue nutrient content, and air temperature (Cramer et al., 1981; Valiela et al., 1985; Robertson,  
16 1986; Twilley et al. 1986a; Robertson, 1988; Mackey and Smail, 1996; Slim et al., 1997).

17         Long-term changes in the elemental content of litter tend to reflect the importance of biotic over abiotic  
18 processes. Our experiments revealed small increases in R. mangle leaf % C with considerable increases in both % N  
19 and % P after nearly one year (Figure 4; Table 2). Increases in the relative N content of R. mangle leaves  
20 overshadowed any change in C, resulting in a large decline in leaf C:N from 90 (initial average) to 34 (final  
21 average). However, N enrichment was surpassed by increases in % P of R. mangle leaf detritus after about 50 days  
22 of decomposition in the field. At this point, N:P began declining to a mean of 132 after 361 days—slightly less than  
23 initial N:P of 142 (Figure 5).

24         A review of estuarine macrophyte decomposition studies supports our findings for tissue N dynamics, with  
25 consistent increases in the % N content of leaves over time regardless of habitat type and climate (Figure 6). This  
26 accumulation of nitrogen through time has been linked to increases in bacterial and fungal protein on the surface of  
27 estuarine macrophyte detritus (Harrison and Mann, 1975; Fell et al., 1975; Cundell et al., 1979; Rice and Tenore,  
28 1981). Nitrogen derived from the water column and soil accounts for a sizable fraction of this immobilized N, but

1 these combined sources do not always account for all accumulated nitrogen in estuarine detritus. It has been shown  
2 that this N budget deficit can be accounted for by nitrogen fixed from the atmosphere (Fell et al., 1975; Valiela et  
3 al., 1985; Twilley et al., 1986a).

4 While this concept may be useful in explaining the long-term increases in litter nitrogen in low N  
5 environments, it does not explain the long-term increases in phosphorus we observed in the P-limited dwarf  
6 mangrove wetland we studied. A few studies have noted long-term increases in litter P; however, they offered little  
7 or no explanation as to the sources or mechanisms for P-accumulation (Twilley et al., 1986b; Tam et al., 1990). In  
8 our litterbag study, the sources of immobilized phosphorus were likely limited to the water column (via direct  
9 precipitation or upland runoff) and soil.

10 Another potentially important source of phosphorus may have been from the leachate of freshly fallen  
11 leaves. When leaves abscise in this dwarf mangrove forest, they accumulate on the forest floor, as there is little  
12 particulate export in this system (S. Davis pers. observ.). It is likely that leachate from recently abscised leaves  
13 supplies a portion of the P accumulated in microbial biomass in older leaves, thus maintaining an efficient recycling  
14 of phosphorus in this system. We contend that the occurrence of P accumulation in dwarf mangrove detritus,  
15 despite extremely low phosphorus concentrations in the surrounding environment (water column TP $\approx$ 0.5  $\mu$ M; pore  
16 water SRP $\approx$ 0.2  $\mu$ M; and soil TP $<$ 25 $\mu$ g cm<sup>-3</sup>; Koch, 1997; Davis et al. 2001), reflected the P-limited status of the  
17 Southern Everglades Ecosystem.

1 **Acknowledgements**

2 We want to acknowledge numerous individuals for their assistance in the field or in the lab. These include, S. Kelly,  
3 D. Rudnick, F. Sklar, C. Madden, C. Donovan, E. Reyes, M. Sutula, B. Perez, J. Davis, L. Romero, and D. Rondeau.  
4 Funding for this research was provided by the South Florida Water Management District (contract # C-E6608) and  
5 the Tropical Biology Program at Florida International University (FIU). This material is also based upon ongoing  
6 work in the Florida Coastal Everglades Long-Term Ecological Research (FCE-LTER) Program and supported by  
7 the National Science Foundation under Grant #9910514. This manuscript was greatly enhanced by comments from  
8 two anonymous reviewers and J. Vermaat. It is SERC Contribution # XX and Contribution # XX from the Tropical  
9 Biology Program at FIU.

1 **References**

- 2 Benner, R., Peele, E.R., Hodson, R.E., 1986. Microbial utilization of dissolved organic matter from the leaves of the  
3 red mangrove, Rhizophora mangle, in the Fresh Creek Estuary, Bahamas. *Estuar. Coast. Shelf Sci.* 23, 607-619.  
4
- 5 Benner, R., Hodson, R.E., 1985. Microbial degradation of the leachable and lignocellulosic components of leaves  
6 and wood from Rhizophora mangle in a tropical mangrove swamp. *Mar. Ecol. Progr. Ser.* 23, 221-230.  
7
- 8 Chale, F.M., 1993. Degradation of mangrove leaf litter under aerobic conditions. *Hydrobiologia* 257, 177-183.
- 9 Corronado-Molina, C., 2000. Litterfall dynamics and nutrient cycling in mangrove forests of Southern Everglades,  
10 Florida and Terminos Lagoon, Mexico. Louisiana State University, Department of Oceanography and Coastal  
11 Sciences, Ph.D. Dissertation.
- 12 Cramer, G.W., Day, J.W., Conner, W.H., 1981. Productivity of four marsh sites surrounding Lake Pontchartrain,  
13 Louisiana. *Am. Mid. Nat.* 106, 65-72.
- 14 Cundell, A.M., Brown, M.S., Stanford, R., Mitchell, R., 1979. Microbial degradation of Rhizophora mangle leaves  
15 immersed in the sea. *Estuar. Coast. Shelf Sci.* 9, 281-286.
- 16 Davis, S.E., Childers, D.L., Day, J.W., Rudnick, D.T., Sklar, F.H., 2001. Carbon, nitrogen and phosphorus dynamics  
17 in a non-tidal dwarf mangrove wetland of the Southern Everglades, Florida, USA. *Estuaries* 42, 609-622.  
18
- 19 Day, J.W., Day, R.H., Barreiro, M.T., Ley-Lou, F., Madden, C.J., 1982. Primary production in the Laguna de  
20 Terminos, a tropical estuary in the Southern Gulf of Mexico. *Oceanol. Acta.* 269-276.
- 21 Day, J.W., Corronado-Molina, C., Vera-Herrera, F.R., Twilley, R., Rivera-Monroy, V.H., Alvarez-Guillen, H., Day,  
22 R., Conner, W., 1996. A 7-year record of above-ground net primary production in a southeastern Mexican  
23 mangrove forest. *Aquat. Bot.* 55, 39-60.
- 24 Fell, J.W., Cefalu, R.C., Master, I.M., Tallman, A.S., 1975. Microbial activities in the mangrove (Rhizophora  
25 mangle) leaf detrital system. In: Walsh, G.E., Snedaker, S.C., Teas, H.J. (eds.), *Proceedings of the International*  
26 *Symposium on Biology and Management of Mangroves*. Institute of Food and Agriculture, University of  
27 Florida, Gainesville. 661-679
- 28 Fell, J.W., Master, I.M., Wiegert, R.G., 1981. Litter decomposition and nutrient enrichment. In: Snedaker, S.C.,  
29 Snedaker, J.G. (eds.), *The Mangrove Ecosystem: Research methods*, 239-251. UNESCO, Bungay, U.K.
- 30 Feller, I.C., 1995. Effects of nutrient enrichment on growth and herbivory of dwarf red mangrove (Rhizophora  
31 mangle). *Ecol. Monogr.* 65, 477-505.
- 32 France, R., Culbert, H., Freeborough, C., Peters, R., 1997. Leaching and early mass loss of boreal leaves and wood  
33 in oligotrophic water. *Hydrobiologia* 345, 209-214.  
34
- 35 Harrison, P.G., Mann, K.H., 1975. Detritus formation from eelgrass (Zostera marina L.): the relative effects of  
36 fragmentation, leaching, and decay. *Limnol. Oceanogr.* 20, 924-934.  
37
- 38 Johnson, L.A., Reynolds, K.A., Foote, A.L., 1993. Production and decomposition of Spartina patens in a degrading  
39 coastal marsh. In O. Magoon, W. Wilson, H. Converse, and L. Tobin, editors, *8th Symposium on Coastal and*  
40 *Ocean Management*. American Shore and Beach Preservation Association. 349-359  
41
- 42 Karmarkar, S.M., 1982. Senescence in mangroves. In Sen, D., Rajpurohit, K. (eds.), *Tasks for Vegetation Science*,  
43 2, 173-187. Dr W. Junk Publishers, The Hague.

44

- 1 Koch, M.S., 1997. Rhizophora mangle L. seedling development into the sapling stage across resource and stress  
2 gradients in subtropical Florida. *Biotropica* 29, 427-439.
- 3 Koch, M.S., Snedaker, S.C., 1997. Factors influencing Rhizophora mangle L. seedling development in Everglades  
4 carbonate soils. *Aquat. Bot.* 59, 87-98.
- 5 Kruczynski, W.L., Subrahmanyam, C.B., Drake, S.H., 1978. Studies on the plant community of a north Florida salt  
6 marsh. *Bull.Mar. Sci.* 28, 316-334.
- 7 Lin, G., Sternberg, L., 1992. Differences in morphology, carbon isotope ratios, and photosynthesis between scrub  
8 and fringe mangroves in Florida, USA. *Aquat. Bot.* 42, 303-313.
- 9 Lin, P., Wang, W., 2001. Changes in the leaf composition, leaf mass and leaf area during leaf senescence in three  
10 species of mangroves. *Ecol. Engineer.* 16, 415-424.
- 11 Mackey, A.P, Smail, G., 1996. The decomposition of mangrove litter in a subtropical mangrove forest.  
12 *Hydrobiologia* 332, 93-98.
- 13 Moran, M.A., Wicks, R.J., Hodson, R.E., 1991. Export of dissolved organic matter from a mangrove swamp  
14 ecosystem: evidence from natural fluorescence, dissolved lignin phenols, and bacterial secondary productivity.  
15 *Mar. Ecol. Progr. Ser.* 76, 175-184.
- 16 Nykvist, N., 1959. Leaching and decomposition of litter I. Experiments on leaf litter of Fraxinus excelsior. *Oikos* 10,  
17 190-211.
- 18  
19 O'Neill, R.V., DeAngelis, D.L., 1981. Comparative productivity and biomass relations of forest ecosystems, In:  
20 Reichle, D.E. (ed.), *Dynamic properties of forest ecosystems*. 411-449. Cambridge University Press,  
21 Cambridge, England.
- 22  
23 Onuf, C.P., Teal, J.M., Valiela, I., 1977. Interactions of nutrients, plant growth and herbivory in a mangrove  
24 ecosystem. *Ecology* 58, 514-526.
- 25 Parsons, W., Taylor, B.R., Parkinson, D., 1990. Decomposition of aspen (Populus tremuloides) leaf litter modified  
26 by leaching. *Can. J. For. Res.* 20, 943-951.
- 27  
28 Rice, D.L., Tenore, K.R., 1981. Dynamics of carbon and nitrogen during the decomposition of detritus derived from  
29 estuarine macrophytes. *Estuar.Coast. Shelf Sci.* 13, 681-690.
- 30 Robertson, A.I., 1986. Leaf-burying crabs: their influence on energy flow and export from mixed mangrove forests  
31 (Rhizophora spp.) in northeastern Australia. *J. Exp. Mar. Biol. Ecol.* 102, 237-248.
- 32 Robertson, A.I., 1988. Decomposition of mangrove leaf litter in tropical Australia. *J. Exp. Mar. Biol. Ecol.* 116, 235-  
33 247.
- 34 Slim, F.J., Hemminga, M.A., Ochieng, C., Jannink, N.T., Cocheret de la Moriniere, E., van der Velde, G., 1997.  
35 Leaf litter removal by the snail Terebralia palustris L. and sesarmid crabs in an east African mangrove forest  
36 (Gazi Bay, Kenya). *J. Exp. Mar. Biol. Ecol.* 215, 35-48.
- 37 Solorzano, L., Sharp, J.H., 1980. Determination of total dissolved phosphorus and particulate phosphorus in natural  
38 waters. *Limnol. Oceanogr.* 25, 754-758.
- 39  
40 Steinke, T.D., Naidoo, G., Charles, L.M., 1983. Degradation of mangrove leaf and stem tissues in situ in Mgeni  
41 Estuary, South Africa. In: Teas, H.J. (ed.), *Biology and Ecology of Mangroves*, 8, 141-149. Dr. W. Junk  
42 Publishers, The Hague.
- 43



- 1 Steinke, T.D., Holland, A.J., Singh, Y., 1993. Leaching losses during decomposition of mangrove leaf litter. *S. Afr.*  
2 *J. Bot.* 59, 21-25.  
3
- 4 Sutula, M., 1999. Processes controlling nutrient transport in the southeastern Everglades wetlands, Florida, USA.  
5 Louisiana State University, Department of Oceanography and Coastal Sciences, Ph.D. Dissertation.  
6
- 7 Tam, N., Vrijmoed, L., Wang, Y., 1990. Nutrient dynamics associated with leaf decomposition in a small  
8 subtropical mangrove community in Hong Kong. *Bull. Mar. Sci.* 47, 68-78.  
9
- 10 Taylor, B.R., 1998. Air-drying depresses rates of leaf litter decomposition. *Soil Biol. Biochem.* 30, 403-412.  
11
- 12 Taylor, B.R., Barlocher, F., 1996. Variable effects of air-drying on leaching losses from tree leaf litter.  
13 *Hydrobiologia* 325, 173-182.  
14
- 15 Twilley, R.R., Lugo, A.E., Patterson-Zucca, C., 1986(a). Litter production and turnover in basin mangrove forests in  
16 southwest Florida. *Ecology* 67, 670-683.  
17
- 18 Twilley, R.R., Ejdung, G., Romare, P., Kemp, W.M., 1986(b). A comparative study of decomposition, oxygen  
19 consumption, and nutrient release for selected aquatic plants occurring in an estuarine environment. *Oikos* 47,  
20 190-198.  
21
- 22 Untawale, A.G., Bhosle, N.B., Dhargalkar, V.K., Matondkar, S., Bukhari, S., 1977. Biochemical changes in  
23 mangrove foliage during growth and decomposition. *Ind. J. Mar. Sci.* 6, 104-106.  
24
- 25 Valiela, I., Teal, J.M., Allen, S.D., Van Etten, R., Goehring, D., Volkman, S., 1985. Decomposition in salt marsh  
26 ecosystems: the phases and major factors affecting disappearance of above-ground organic matter. *J. Exp. Mar.*  
27 *Biol. Ecol.* 89, 29-54.  
28
- 29 Wafar, S., Untawale, A.G., Wafar, M., 1997. Litterfall and energy flux in a mangrove ecosystem. *Estuar. Coast. Shelf*  
30 *Sci.* 44, 111-124.  
31
- 32 White, D.A., Weiss, T.E., Trapani, J.M., Thien, L.B., 1978. Productivity and decomposition of the dominant salt  
33 marsh plants in Louisiana. *Ecology* 59, 751-759.

1 Table 1: Water concentrations and fluxes ( $\pm$  standard deviation) of TOC, TN, and  
 2 TP in mangrove leaf leaching bottles over time. Fluxes of C, N, and P were  
 3 calculated as change in water concentration over time normalized to grams dry  
 4 weight (gdw) of each leaf. There was a net release of all constituents from the leaf  
 5 to the water column. Different letters represent significant differences over time  
 6 ( $p < 0.05$ ) using single factor ANOVAs and Tukey post-hoc comparisons.

7

Constituent (treatment level)	Day	Concentration ( $\mu\text{mol}$ )	Letter	Flux ( $\mu\text{mol gdw}^{-1} \text{d}^{-1}$ )
TOC (no azide)	0	168 $\pm$ 1	A	
	1	781 $\pm$ 286	A	1259 $\pm$ 574
	2	919 $\pm$ 219	AB	340 $\pm$ 127
	5	1513 $\pm$ 225	AB	543 $\pm$ 174
	10	2364 $\pm$ 207	B	442 $\pm$ 109
	21	3061 $\pm$ 873	B	284 $\pm$ 131
TN (no azide)	0	11.7 $\pm$ 0.1	A	
	1	14.6 $\pm$ 0.8	A	13.9 $\pm$ 2.7
	2	15.1 $\pm$ 0.5	AB	3.2 $\pm$ 0.2
	5	19.9 $\pm$ 1.3	B	5.3 $\pm$ 1.9
	10	23 $\pm$ 1.5	B	3.5 $\pm$ 1.2
	21	29.1 $\pm$ 1.7	C	1.9 $\pm$ 0.3
TP (no azide)	0	0.03 $\pm$ 0.01	A	
	1	0.13 $\pm$ 0.03	A	0.2 $\pm$ 0.07
	2	0.19 $\pm$ 0.04	A	0.1 $\pm$ 0.02
	5	0.61 $\pm$ 0.19	B	0.2 $\pm$ 0.03
	10	0.4 $\pm$ 0.02	AB	0.08 $\pm$ 0.02
	21	0.52 $\pm$ 0.06	B	0.04 $\pm$ 0.00
TOC (azide)	0	173 $\pm$ 0	A	
	1	207 $\pm$ 7	AB	215 $\pm$ 54
	2	997 $\pm$ 465	AB	827 $\pm$ 342
	5	1186 $\pm$ 403	ABC	468 $\pm$ 196
	10	1483 $\pm$ 1032	BC	169 $\pm$ 92
	21	2964 $\pm$ 448	C	219 $\pm$ 47
TP (azide)	0	0.04 $\pm$ 0.01	A	
	1	0.11 $\pm$ 0.01	AB	0.16 $\pm$ 0.05
	2	0.43 $\pm$ 0.18	B	0.33 $\pm$ 0.19
	5	0.36 $\pm$ 0.12	B	0.15 $\pm$ 0.08
	10	0.38 $\pm$ 0.17	B	0.07 $\pm$ 0.03
	21	0.56 $\pm$ 0.02	B	0.04 $\pm$ 0.00

1 Table 2: Percent changes in relative C, N, and P content from initial values during the decomposition of a number of estuarine macrophytes. Studies  
 2 were categorized as "litterbag", if leaf material was contained within a mesh bag of some sort, or "leaching", if leaf material was incubated directly in  
 3 water without a litterbag. All changes are reported as percentages of initial relative values. Positive values indicate an increase while negative values  
 4 indicate a decrease in litter content. Some values were calculated from tabular data, but most were estimated from plotted data.  
 5

Study type	Species	Tissue treatment	Time (days)	% chg. in rel. C	% chg. in rel. N	% chg. in rel. P	Reference
leaching	<i>Avicennia marina</i>	oven-dried material, submerged	2		3	-17	Chale 1993
litterbag	<i>Rhizophora mangle</i>	fresh material, submerged	7	<1	30		Cundell et al. 1979
litterbag	<i>Aegiceras corniculatum</i>	air-dried material, tidally inundated	13	19	2	22	Tam et al. 1990 <sup>a</sup>
litterbag	<i>Avicennia marina</i>	air-dried material, tidally inundated	13	10	-18	-52	Tam et al. 1990 <sup>a</sup>
litterbag	<i>Kandelia kandel</i>	air-dried material, tidally inundated	13	9	-39	50	Tam et al. 1990 <sup>a</sup>
leaching	<i>Avicennia marina</i>	fresh material, poisoned	14		8	-72	Steinke et al. 1993
leaching	<i>Avicennia marina</i>	air-dried material, poisoned	14		1	-84	Steinke et al. 1993
litterbag	<i>Spartina alterniflora</i>	air-dried material, tidally inundated	14	6	7		Valiela et al. 1985 <sup>b</sup>
litterbag	<i>Spartina patens</i>	air-dried material, tidally inundated	14	-2	<1		Valiela et al. 1985 <sup>b</sup>
litterbag	<i>Rhizophora mangle</i>	fresh material, submerged	15	9	50		Fell et al. 1975
leaching	<i>Spartina alterniflora</i>	fresh material, submerged	15	-1	3	13	Twilley et al. 1986b
litterbag	<i>Avicennia marina</i>	fresh material, submerged	21	1	38	-42	Steinke et al. 1983
litterbag	<i>Brugueira gymnorhiza</i>	fresh material, submerged	21	-1	36	-50	Steinke et al. 1983
leaching	<i>Rhizophora mangle</i>	fresh material, submerged	21	3	-2	-34	this study <sup>c</sup>

6

1 Table 2: continued

2

Study type	Vegetation	Tissue treatment	Time (days)	% chg. in rel. C	% chg. in rel. N	% chg. in rel. P	Reference
litterbag	<i>Rhizophora mangle</i>	fresh material, submerged	70	-22	75		Cundell et al. 1979
litterbag	<i>Aegiceras corniculatum</i>	air-dried material, tidally inundated	81	-5	59	-30	Tam et al. 1990 <sup>a</sup>
litterbag	<i>Kandelia kandel</i>	air-dried material, tidally inundated	81	-9	-15	200	Tam et al. 1990 <sup>a</sup>
leaching	<i>Spartina alterniflora</i>	fresh material	93	2	90	93	Twilley et al. 1986b
leaching	<i>Avicennia marina</i>	oven-dried material	98		105	-32	Chale 1993
litterbag	<i>Avicennia marina</i>	air-dried material, tidally inundated	109	6	26	50	Tam et al. 1990 <sup>a</sup>
litterbag	<i>Rhizophora mangle</i>	fresh material, submerged	115	21	148		Fell et al. 1975
litterbag	<i>Avicennia marina</i>	fresh material, submerged	175	-10	91	-54	Steinke et al. 1983
litterbag	<i>Rhizophora mangle</i>	air-dried material, tidally-inundated	180		209	200	Day et al. 1982 <sup>d</sup>
litterbag	<i>Brugueira gymnorrhiza</i>	fresh material, submerged	189	11	164	-55	Steinke et al. 1983
litterbag	<i>Rhizophora mangle</i>	fresh material	361	6	181	207	this study
litterbag	<i>Spartina alterniflora</i>	air-dried material, tidally inundated	700	-19	157		Valiela et al. 1985 <sup>b</sup>
litterbag	<i>Spartina patens</i>	air-dried material, tidally inundated	700	-2	250		Valiela et al. 1985 <sup>b</sup>

3

4 <sup>a</sup> Freshly-picked green leaves were used in this study.

5 <sup>b</sup> Data taken from un-enriched low marsh and high marsh sites.

6 <sup>c</sup> Data from non-poisoned bottles only.

7 <sup>d</sup> Data collected from Estero Pargo, MX site.

1 **List of Figures**

2 Fig. 1. Map of the Southern region of Everglades National Park with Taylor Slough borders and indication (\*) of  
3 litter collection and decomposition sites (A & B) adjacent to Taylor River. Units for lat/long are in degrees with  
4 decimal minutes.


5  
6 Fig. 2. Cell plots of: a) % dry mass remaining, b) relative C, N, and P and c) absolute C, N, and P in red mangrove  
7 leaves over the course of the 21-day leaching experiment. Bars indicate standard deviation of three replicates.

8  
9 Fig. 3. Molar ratios of C:N (top) and N:P (bottom) in red mangrove leaves over the course of the 21-day leaching  
10 experiment. Bars indicate standard deviations of three replicates.








11  
12 Fig. 4. Cell plots of: a) % dry mass remaining, b) relative C, N, and P and c) absolute C, N, and P in red mangrove  
13 leaves over the course of the 361-day litterbag experiment. Bars indicate standard deviation of three replicates.

14  
15 Fig. 5. Molar ratios of C:N (top) and N:P (bottom) in red mangrove leaves over the course of the 361-day litterbag  
16 experiment. Bars indicate standard deviations of three replicates.


17  
18 Fig. 6. Regression of % change in litter N content versus time for a number of decomposing estuarine macrophytes  
19 (5 species of mangrove and 2 species of cordgrass). Tabular data and reference for each decomposition study are  
20 provided in Table 2. Data from this study indicated with open circles.



1












1




1







1



1