

5-31-2016

# Response to letter regarding article by Patel et al: A Novel Biomarker of Oxidative Stress is Associated with Risk of Death in Patients with Coronary Artery Disease

Riyaz S. Patel

*Emory University; University College London*

Nima Ghasemzadeh

*Emory University*

Danny J. Eapen

*Emory University*

Salman Sher

*Emory University*

Shawn Arshad

*Emory University*

*See next page for additional authors*

Follow this and additional works at: [https://digitalcommons.fiu.edu/biostatistics\\_fac](https://digitalcommons.fiu.edu/biostatistics_fac)

 Part of the [Medicine and Health Sciences Commons](#)

## Recommended Citation

Patel, Riyaz S.; Ghasemzadeh, Nima; Eapen, Danny J.; Sher, Salman; Arshad, Shawn; Ko, Yian; Veledar, Emir; Samady, Habib; Zafari, A. Maziar; Sperling, Laurence; Vaccarino, Viola; Jones, Dean P.; and Quyyumi, Arshed A., "Response to letter regarding article by Patel et al: A Novel Biomarker of Oxidative Stress is Associated with Risk of Death in Patients with Coronary Artery Disease" (2016).

*Department of Biostatistics Faculty Publications*. 30.

[https://digitalcommons.fiu.edu/biostatistics\\_fac/30](https://digitalcommons.fiu.edu/biostatistics_fac/30)

This work is brought to you for free and open access by the Robert Stempel College of Public Health & Social Work at FIU Digital Commons. It has been accepted for inclusion in Department of Biostatistics Faculty Publications by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu).

---

**Authors**

Riyaz S. Patel, Nima Ghasemzadeh, Danny J. Eapen, Salman Sher, Shawn Arshad, Yian Ko, Emir Veledar, Habib Samady, A. Maziar Zafari, Laurence Sperling, Viola Vaccarino, Dean P. Jones, and Arshed A. Quyyumi



Published in final edited form as:

*Circulation*. 2016 May 31; 133(22): e667. doi:10.1161/CIRCULATIONAHA.116.022753.

## Response to letter regarding article by Patel et al: A Novel Biomarker of Oxidative Stress is Associated with Risk of Death in Patients with Coronary Artery Disease

Riyaz S. Patel, MD<sup>1,2</sup>, Nima Ghasemzadeh, MD<sup>1</sup>, Danny J Eapen, MD<sup>1</sup>, Salman Sher, MD<sup>1</sup>, Shawn Arshad, MD<sup>1</sup>, Yi-an Ko, PhD<sup>3</sup>, Emir Veledar, PhD<sup>4,5</sup>, Habib Samady, MD<sup>1</sup>, A. Maziar Zafari, MD, PhD<sup>1,6</sup>, Laurence Sperling, MD<sup>1</sup>, Viola Vaccarino, MD, PhD<sup>1,5</sup>, Dean P Jones, PhD<sup>1</sup>, and Arshed A. Quyyumi, MD<sup>1,\*</sup>

<sup>1</sup>Dept of Medicine, Emory University School of Medicine, Atlanta, GA, USA

<sup>2</sup>Institute of Cardiovascular Science, University College London, London, UK

<sup>3</sup>Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Atlanta, GA, USA

<sup>4</sup>Dept of Medicine, Baptist Health South Florida, FI, USA

<sup>5</sup>Dept of Epidemiology, Rollins School of Public Health, Atlanta, GA, USA

<sup>6</sup>Dept of Medicine, Atlanta Veterans Affairs Medical Center, Decatur, GA, USA

---

Dear Sirs,

We thank Drs Giral and colleagues for their interest in our work.<sup>1</sup> They raise the important query of whether our findings would still persist after adjustment for Gamma Glutamyltransferase (GGT), given that GGT activity hydrolyzes GSH to produce glutamate + cysteinyl glycine. This point however is not relevant to our description of GSH/CySS as a useful biomarker of cardiovascular disease because our samples were all collected with a preservation solution containing a GGT inhibitor to prevent the very problem that they outline. Their letter is useful, however, in providing an opportunity to re-emphasize the point, that there is need for a GSH preservation solution for this assay. In our preservation solution, we use serine-borate, a well-known inhibitor of GGT that has been widely used to prevent this artifact since its discovery in 1959.<sup>2</sup> In our original development of methods, the efficacy of the preservation solution to prevent artifactual degradation was validated by showing that it completely prevented loss of GSH when known concentrations of GSH were added to plasma.<sup>3</sup> Earlier studies by CV Smith et al showed that another commonly used GGT inhibitor, acivicin, was not as effective because it required enzyme turnover to obtain complete inhibition.<sup>4</sup> In use of the serine-borate-containing preservation solution to study aminothiols we found that the distribution of GSH, Cys-Gly and CySS in freshly collected human plasma showed metabolite correlations indicating that GSH metabolism in vivo in

---

For Correspondence: Arshed A. Quyyumi, Emory University School of Medicine, 1462 Clifton Road NE, Suite 507, Atlanta, GA, 30322, USA, Fax: 404 727 8785 Tel: 404 727 3655 aquyyum@emory.edu.

Disclosures: None

plasma occurs by two mechanisms, hydrolysis by GGT to Glu + Cys-Gly and reaction of GSH with cystine to produce Cys + cysteine-GSH disulfide. We reiterated the need to inhibit GGT in our most recent description of this method.<sup>5</sup>

## References

1. Patel RS, Ghasemzadeh N, Eapen DJ, Sher S, Arshad S, Ko YA, Veledar E, Samady H, Zafari AM, Sperling L, Vaccarino V, Jones DP, Quyyumi AA. Novel biomarker of oxidative stress is associated with risk of death in patients with coronary artery disease. *Circulation*. 2016; 133:361–369. [PubMed: 26673559]
2. Revel JP, Ball EG. The reaction of glutathione with amino acids and related compounds as catalyzed by gamma-glutamyl transpeptidase. *The Journal of biological chemistry*. 1959; 234:577–582. [PubMed: 13641263]
3. Jones DP, Carlson JL, Samiec PS, Sternberg P Jr, Mody VC Jr, Reed RL, Brown LA. Glutathione measurement in human plasma. Evaluation of sample collection, storage and derivatization conditions for analysis of dansyl derivatives by hplc. *Clinica chimica acta; international journal of clinical chemistry*. 1998; 275:175–184.
4. Smith CV, Hansen TN, Martin NE, McMicken HW, Elliott SJ. Oxidant stress responses in premature infants during exposure to hyperoxia. *Pediatric research*. 1993; 34:360–365. [PubMed: 8134179]
5. Jones DP, Liang Y. Measuring the poise of thiol/disulfide couples in vivo. *Free radical biology & medicine*. 2009; 47:1329–1338. [PubMed: 19715755]