

3-16-2005

South Florida Coastal Water Quality Monitoring Network - 10-12/04 Quarterly Report (C-15397)

Joseph N. Boyer

Southeast Environmental Research Center, Florida International University, boyerj@fiu.edu

Follow this and additional works at: <https://digitalcommons.fiu.edu/sercrp>

 Part of the [Environmental Monitoring Commons](#), and the [Water Resource Management Commons](#)

Recommended Citation

Boyer, Joseph N., "South Florida Coastal Water Quality Monitoring Network - 10-12/04 Quarterly Report (C-15397)" (2005). SERC Research Reports. 23.
<https://digitalcommons.fiu.edu/sercrp/23>



Southeast Environmental Research Center
OE-148 Florida International University, Miami, FL 33199
305-348-3095, 305-348-4096 fax, <http://serc.fiu.edu>

16 March 2005

Carole Milliman Maddox
SFWMD
8894 Belvedere Road
West Palm Beach, FL 33411

Re: South Florida Coastal Water Quality Monitoring Network – 10-12/04 Quarterly Report (C-15397)

Dear Ms. Maddox:

This letter serves to transmit the South Florida Coastal Water Quality Monitoring Network Quarterly Report as per our SFWMD/SERC Cooperative Agreement #C-15397. This report consists of this letter along with corresponding tables and figures.

Project Background

This report includes water quality data collected monthly during the annual period of record (POR) Oct. – Dec. 2004 from 28 stations in Florida Bay, 22 stations in Whitewater Bay, 25 stations in Ten Thousand Islands, 25 stations in Biscayne Bay, and 28 stations in Cape Romano-Rookery Bay-Pine Island Sound. A total of 49 stations were also collected on the SW Florida Shelf on a quarterly basis. Figure 1 shows the location of the fixed sampling stations.

Water quality parameters monitored at each station include the dissolved nutrients nitrate + nitrite (NO_x), nitrite (NO_2), nitrate (NO_3), ammonium (NH_4), inorganic nitrogen (DIN), and soluble reactive phosphorus (SRP). Silicate (Si(OH)_4) was analyzed at all stations on a quarterly basis in conjunction with SW Shelf sampling. Total concentrations of nitrogen (TN), organic nitrogen (TON), phosphorus (TP), and organic carbon (TOC) were also measured. All concentrations for each of these parameters are reported as parts per million (ppm) except where noted.

Biological parameters monitored included chlorophyll a ($\mu\text{g l}^{-1}$) and alkaline phosphatase activity (APA; $\mu\text{M hr}^{-1}$). Field parameters measured at both surface and bottom of the water column include salinity, dissolved oxygen (DO; mg l^{-1}), and temperature ($^{\circ}\text{C}$). Turbidity (NTU) of the surface water was also measured.

Data Results

A previous spatial analysis of data from Florida Bay resulted in the delineation of 3 groups of stations which have robust similarities in water quality (Fig. 2). We have argued that these spatially contiguous groups of stations are the result of similar loading and processing of materials, hence we call them 'zones of similar influence'. The Eastern Bay zone (FBE) acts most like a 'conventional' estuary in that it has a quasi-longitudinal salinity gradient caused by the mixing of freshwater runoff with seawater. In contrast, the Central Bay (FBC) is a hydrographically isolated area with low and infrequent terrestrial freshwater input, a long water residence time, and high evaporative potential. The Western Bay zone (FBW) is the most influenced by the Gulf of Mexico tides and is also isolated from direct overland freshwater sources. Station #7 - Highway Creek did not cluster out with any of the Florida Bay stations and was considered separately.

Using the same statistical approach as above, the TTI-WWB complex was partitioned into 6 distinct zones of similar water quality (Fig. 3). The first cluster was composed of 13 stations in and around the Shark, Harney, Broad, and Lostmans Rivers and is called the Mangrove River (MR) group. This cluster also included a sampling station just off the Faka Union Canal. The second cluster was made up of the 8 stations enclosed within Whitewater Bay proper (WWB). Twelve stations situated mostly in and around the coastal islands of TTI-WWB formed the Gulf Island group (GI). The water quality characteristics at the Coot Bay site were sufficiently different so as to be a cluster of its own. The next cluster contained the northernmost 2 stations in the Blackwater River estuary (BLK). Finally, the Inland Wilderness Waterway zone (IWW) included 11 stations distributed throughout the inside passage as well as the Chatham River and the station off Everglades City.

Biscayne Bay was partitioned into 6 distinct ZSI using the above statistical analysis. The first cluster was composed of 2 stations closest to the shore in the south Bay (Fig. 4); they were called the Alongshore group (AS). These are stations most influenced by the Goulds, Military and Mowry Canals. The second cluster was made up of the 5 stations farther from the coast called Inshore (IS). Thirteen stations situated mostly in the bay proper were called the main Bay (MAIN) group. The next cluster contained 3 stations situated in areas of great tidal exchange (ocean channel, not shown). Two stations in Card Sound grouped together SCARD. For purposes of this report, the stations added to the area north of the Rickenbacker Causeway are defined, *a priori*, as a distinct cluster, North Bay (NBAY).

The above statistical analysis objectively classified the 49 Shelf sampling sites into 3 zones having similar water quality (Fig. 5). The first cluster was composed of only 2 stations which were closest to the shore off Cape Sable; they were called the SHARK group, after the Shark River, the main source of freshwater to the region. The second cluster was made up of the 7 more northerly stations nearest the coast and called SHOAL. The remaining stations were called the SHELF group.

Sampling in the Rookery Bay area began Jan. 1999, so we now have 5 years of data available. But because of the very heterogeneous nature of the area, we will continue to use generally accepted geomorphological characteristics to group the stations (Fig. 6). These groupings are Cocohatchee River (COCO), Estero Bay (EST), Cape Romano-Marco Island (MARC), Naples Bay (NPL), Pine Island Sound (PIS), Rookery Bay (RB), and San Carlos Bay (SCB).

Data are also reported as box-and-whiskers plots (Figs. 7-28). The center horizontal line in the box is the median of the data, the top and bottom of the box are the 25th and 75th percentiles (quartiles), and the ends of the whiskers are the 5th and 95th percentiles.

Summary statistics of all water quality parameters by ecosystem are shown in Table 1. The median was chosen because it is a more accurate measure of central tendency in non-normally distributed water quality data. The range is expressed as the minimum (Min.) and maximum (Max.) values for the POR, and *n* is the number of data points used in the analysis.

If you have any questions about the content of this report, please do not hesitate to contact me at 305-348-4076 or boyerj@fiu.edu.

Sincerely,

A handwritten signature in blue ink, appearing to read "Joseph N. Boyer".

Joseph N. Boyer, Ph.D.
Associate Director and Scientist





Figure 1: All fixed water quality stations funded by this SFWMD project.

Florida Bay Water Quality Zones



Eastern Bay (+), Central Bay, (■), Western Bay (●)

Figure 2. Florida Bay zones.




Figure 3. WWB-TTI water quality zones.





Figure 4. Biscayne Bay water quality zones.



■ Shark, ● Shoal, + Shelf

Figure 5. SW Florida Shelf water quality zones.




Figure 6. SW estuaries.

Eastern Florida Bay Zone




Figure 7. Box-and-whisker plots of water quality in Eastern Florida Bay by survey.

Central Florida Bay Zone




Figure 8. Box-and-whisker plots of water quality in Central Florida Bay by survey.

Western Florida Bay Zone




Figure 9. Box-and-whisker plots of water quality in Western Florida Bay by survey.

Whitewater Bay Zone




Figure 10. Box-and-whisker plots of water quality in WWB-TTI by survey.

Mangrove Rivers Zone




Figure 11. Box-and-whisker plots of water quality in WWB-TTI by survey.

Gulf Islands Zone




Figure 12. Box-and-whisker plots of water quality in WWB-TTI by survey.

Inner Waterway Zone




Figure 13. Box-and-whisker plots of water quality in WWB-TTI by survey.

Blackwater River Zone




Figure 14. Box-and-whisker plots of water quality in WWB-TTI by survey.

Alongshore Zone




Figure 15. Box-and-whisker plots of water quality in Biscayne Bay by survey.

Inshore Zone




Figure 16. Box-and-whisker plots of water quality in Biscayne Bay by survey.

Main Bay Zone




Figure 17. Box-and-whisker plots of water quality in Biscayne Bay by survey.

South Card Sound Zone




Figure 18. Box-and-whisker plots of water quality in Biscayne Bay by survey.

North Bay Zone




Figure 19. Box-and-whisker plots of water quality in Biscayne Bay by survey.




Figure 20. Box-and-whisker plots of water quality in SW Florida Shelf by survey.




Figure 21. Box-and-whisker plots of water quality in SW Florida Shelf by survey.




Figure 22. Box-and-whisker plots of water quality in SW Florida Shelf by survey.

Marco Zone




Figure 23. Box-and-whisker plots of water quality in RB-PIS by survey.




Figure 24. Box-and-whisker plots of water quality in RB-PIS by survey.




Figure 25. Box-and-whisker plots of water quality in RB-PIS by survey.

San Carlos Bay Zone




Figure 26. Box-and-whisker plots of water quality in RB-PIS by survey.




Figure 27. Box-and-whisker plots of water quality in RB-PIS by survey.

Pine Island Sound Zone




Figure 28. Box-and-whisker plots of water quality in RB-PIS by survey.

Table 1.

Parameter	Zone	Median	Min.	Max.	n
Alkaline Phosphatase Activity ($\mu\text{M h}^{-1}$)	Biscayne Bay	0.128	0.026	0.565	75
	Florida Bay	0.280	0.059	2.908	84
	Rookery Bay	0.067	0.028	0.230	87
	SW Shelf	0.050	0.017	0.157	49
	Ten Thousand Is.	0.060	0.022	0.956	78
	Whitewater Bay	0.437	0.037	3.741	66
Chlorophyll <i>a</i> ($\mu\text{g l}^{-1}$)	Biscayne Bay	0.451	0.124	3.140	75
	Florida Bay	0.501	0.173	9.812	84
	Rookery Bay	4.444	0.819	16.453	87
	SW Shelf	1.210	0.278	3.835	49
	Ten Thousand Is.	2.795	1.037	7.806	78
	Whitewater Bay	2.305	0.714	12.689	66
Surface Dissolved Oxygen (mg l^{-1})	Biscayne Bay	6.31	4.95	8.16	75
	Florida Bay	6.54	4.27	9.67	84
	Rookery Bay	6.06	3.64	9.16	87
	SW Shelf	6.30	5.24	7.26	49
	Ten Thousand Is.	5.94	3.43	7.87	78
	Whitewater Bay	5.59	1.34	7.17	66
Bottom Dissolved Oxygen (mg l^{-1})	Biscayne Bay	6.40	4.95	8.25	75
	Florida Bay	6.50	4.30	10.32	84
	Rookery Bay	5.83	0.95	7.53	87
	SW Shelf	6.03	5.35	6.83	49
	Ten Thousand Is.	5.88	3.40	7.96	78
	Whitewater Bay	5.25	1.08	10.19	66
NH_4^+ (ppm)	Biscayne Bay	0.013	0.001	0.158	75
	Florida Bay	0.032	0.001	0.278	84
	Rookery Bay	0.009	0.001	0.122	87
	SW Shelf	0.002	0.000	0.006	49
	Ten Thousand Is.	0.019	0.002	0.079	78
	Whitewater Bay	0.027	0.010	0.483	66

Parameter	Zone	Median	Min.	Max.	n
NO ₂ ⁻ (ppm)	Biscayne Bay	0.003	0.001	0.028	75
	Florida Bay	0.003	0.000	0.021	84
	Rookery Bay	0.001	0.000	0.023	87
	SW Shelf	0.001	0.000	0.002	49
	Ten Thousand Is.	0.004	0.001	0.010	78
	Whitewater Bay	0.004	0.000	0.011	66
NO ₃ ⁻ (ppm)	Biscayne Bay	0.012	0.000	1.173	74
	Florida Bay	0.005	0.000	0.054	84
	Rookery Bay	0.003	0.000	0.378	87
	SW Shelf	0.001	0.000	0.005	49
	Ten Thousand Is.	0.013	0.001	0.051	78
	Whitewater Bay	0.017	0.000	0.057	66
pH	Biscayne Bay	8.055	7.890	8.200	75
	Florida Bay	8.137	7.870	8.510	84
	Rookery Bay	7.910	7.345	8.150	87
	SW Shelf	7.950	7.670	8.140	49
	Ten Thousand Is.	7.720	7.240	7.980	78
	Whitewater Bay	7.775	7.130	8.345	66
Surface Salinity	Biscayne Bay	32.84	16.25	37.76	75
	Florida Bay	36.39	3.43	42.93	84
	Rookery Bay	32.87	1.87	35.94	87
	SW Shelf	35.92	31.13	38.91	49
	Ten Thousand Is.	23.91	0.98	35.84	78
	Whitewater Bay	7.22	0.31	28.48	66
Bottom Salinity	Biscayne Bay	33.77	16.32	37.75	75
	Florida Bay	36.57	4.26	43.13	84
	Rookery Bay	33.40	2.10	35.97	87
	SW Shelf	35.99	31.15	38.94	49
	Ten Thousand Is.	25.01	1.29	35.83	78
	Whitewater Bay	7.56	0.28	31.35	66
Si(OH) ₄ (ppm)	Biscayne Bay	0.197	0.049	1.203	25
	Florida Bay	0.164	0.019	1.989	28
	Rookery Bay	0.782	0.247	3.727	29
	SW Shelf	0.056	0.002	0.673	45
	Ten Thousand Is.	2.176	1.271	3.105	26
	Whitewater Bay	2.402	0.546	6.400	22

Parameter	Zone	Median	Min.	Max.	n
Soluble	Biscayne Bay	0.001	0.000	0.008	75
Reactive	Florida Bay	0.001	0.000	0.004	84
Phosphorus	Rookery Bay	0.007	0.001	0.089	87
(ppm)	SW Shelf	0.001	0.000	0.004	49
	Ten Thousand Is.	0.007	0.000	0.020	78
	Whitewater Bay	0.002	0.001	0.006	66
Surface	Biscayne Bay	23.70	16.87	28.11	75
Temperature	Florida Bay	23.29	16.39	27.43	84
(°C)	Rookery Bay	23.60	21.64	30.95	87
	SW Shelf	27.62	23.42	28.12	49
	Ten Thousand Is.	24.56	22.12	27.90	78
	Whitewater Bay	26.44	23.18	29.02	66
Bottom	Biscayne Bay	23.70	16.90	27.71	75
Temperature	Florida Bay	23.42	16.43	27.77	84
(°C)	Rookery Bay	23.60	21.37	31.01	87
	SW Shelf	27.67	23.30	28.13	49
	Ten Thousand Is.	24.46	21.98	27.74	78
	Whitewater Bay	26.39	22.66	29.30	66
Total	Biscayne Bay	0.185	0.095	1.560	74
Nitrogen	Florida Bay	0.323	0.143	0.806	84
(ppm)	Rookery Bay	0.235	0.110	1.250	87
	SW Shelf	0.150	0.075	0.552	49
	Ten Thousand Is.	0.343	0.171	1.589	78
	Whitewater Bay	0.436	0.253	1.505	66
Total	Biscayne Bay	2.921	1.231	6.186	75
Organic	Florida Bay	7.535	2.322	27.370	84
Carbon	Rookery Bay	4.159	2.202	17.640	87
(ppm)	SW Shelf	2.069	1.236	6.091	49
	Ten Thousand Is.	8.830	2.897	23.348	78
	Whitewater Bay	12.265	5.786	31.698	65
Total	Biscayne Bay	0.144	0.066	0.322	74
Organic	Florida Bay	0.251	0.095	0.764	84
Nitrogen	Rookery Bay	0.213	0.106	1.229	87
(ppm)	SW Shelf	0.148	0.072	0.548	49
	Ten Thousand Is.	0.296	0.161	1.507	78
	Whitewater Bay	0.389	0.087	1.008	66

Parameter	Zone	Median	Min.	Max.	n
Total Phosphorus (ppm)	Biscayne Bay	0.005	0.002	0.037	75
	Florida Bay	0.006	0.002	0.039	84
	Rookery Bay	0.034	0.018	0.118	87
	SW Shelf	0.011	0.004	0.023	49
	Ten Thousand Is.	0.037	0.008	0.273	78
	Whitewater Bay	0.014	0.006	0.056	66
Turbidity (NTU)	Biscayne Bay	0.72	0.20	11.53	75
	Florida Bay	2.03	0.43	18.56	84
	Rookery Bay	2.89	0.55	9.59	87
	SW Shelf	1.30	0.68	9.57	49
	Ten Thousand Is.	4.09	0.66	9.17	78
	Whitewater Bay	2.44	0.59	12.94	66