

2010

# DNA linking number change induced by sequence-specific DNA-binding proteins

Bo Chen

*Department of Chemistry and Biochemistry, Florida International University*

Yazhong Xiao

*Department of Chemistry and Biochemistry, Florida International University*

Chang Liu

*Department of Biomedical Engineering, Florida International University, chliu@fiu.edu*

Chen-zhong Li

*Department of Biomedical Engineering, Florida International University, licz@fiu.edu*

Fenfei Leng

*Department of Chemistry and Biochemistry, Florida International University, lengf@fiu.edu*

Follow this and additional works at: [https://digitalcommons.fiu.edu/chemistry\\_fac](https://digitalcommons.fiu.edu/chemistry_fac)



Part of the [Biomedical Engineering and Bioengineering Commons](#), and the [Chemistry Commons](#)

---

## Recommended Citation

DNA linking number change induced by sequence-specific DNA-binding proteins Bo Chen, Yazhong Xiao, Chang Liu, Chenzhong Li, Fenfei Leng *Nucleic Acids Res.* 2010 Jun; 38(11): 3643–3654. Published online 2010 Feb 25. doi: 10.1093/nar/gkq078

This work is brought to you for free and open access by the College of Arts, Sciences & Education at FIU Digital Commons. It has been accepted for inclusion in Department of Chemistry and Biochemistry by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu).

# DNA linking number change induced by sequence-specific DNA-binding proteins

Bo Chen<sup>1</sup>, Yazhong Xiao<sup>1</sup>, Chang Liu<sup>2</sup>, Chenzhong Li<sup>2</sup> and Fenfei Leng<sup>1,\*</sup>

<sup>1</sup>Department of Chemistry & Biochemistry, Florida International University, Miami, FL 33199 and <sup>2</sup>Department of Biomedical Engineering, Florida International University, Miami, FL 33174, USA

Received September 4, 2010; Revised January 16, 2010; Accepted January 26, 2010

## ABSTRACT

**Sequence-specific DNA-binding proteins play a key role in many fundamental biological processes, such as transcription, DNA replication and recombination. Very often, these DNA-binding proteins introduce structural changes to the target DNA-binding sites including DNA bending, twisting or untwisting and wrapping, which in many cases induce a linking number change ( $\Delta Lk$ ) to the DNA-binding site. Due to the lack of a feasible approach,  $\Delta Lk$  induced by sequence-specific DNA-binding proteins has not been fully explored. In this paper we successfully constructed a series of DNA plasmids that carry many tandem copies of a DNA-binding site for one sequence-specific DNA-binding protein, such as  $\lambda$  O, LacI, GalR, CRP and AraC. In this case, the protein-induced  $\Delta Lk$  was greatly amplified and can be measured experimentally. Indeed, not only were we able to simultaneously determine the protein-induced  $\Delta Lk$  and the DNA-binding constant for  $\lambda$  O and GalR, but also we demonstrated that the protein-induced  $\Delta Lk$  is an intrinsic property for these sequence-specific DNA-binding proteins. Our results also showed that protein-mediated DNA looping by AraC and LacI can induce a  $\Delta Lk$  to the plasmid DNA templates. Furthermore, we demonstrated that the protein-induced  $\Delta Lk$  does not correlate with the protein-induced DNA bending by the DNA-binding proteins.**

## INTRODUCTION

Sequence-specific DNA-binding proteins play a key role in many essential biological processes. For instance, DNA replication initiators, such as *Escherichia coli* DnaA protein and bacteriophage  $\lambda$  O protein, bind and restructure DNA replication origin elements (1,2) and

subsequently prepare them for chromosome duplication (3–6). Transcription is another fundamental process in biology. The control of transcription initiation by transcriptional factors is extremely important for coordinating a sophisticated response to changes in the environment (7). Transcriptional factors either stimulate or inhibit the transcription initiation, depending on different scenarios (8,9). Among the best-characterized transcriptional factors are *E. coli* lac repressor (LacI) (10), gal repressor (GalR) (11), cAMP receptor protein (CRP) (12) and AraC protein of the L-arabinose operon (8). These transcriptional factors tightly bind to their recognition sites on the chromosome, restructure the recognition sites and regulate transcription from the nearby or distant promoters (8,13).

One common property of the sequence-specific DNA-binding proteins is to induce DNA structural change to their recognition sites, which includes DNA bending, twisting or untwisting, wrapping and other distortions (14,15). Among the DNA structural changes, protein-induced DNA bending has been extensively studied (16). For example, *E. coli* LacI, GalR, CRP and AraC all sharply bend their recognition sites (17,18). In fact, the protein-induced DNA bending was considered to be necessary for their biological functions (18). Nevertheless, other protein-induced DNA structural changes by sequence-specific DNA-binding proteins are rarely addressed (14). The main reason is that a reliable strategy has not been developed to carry out a systematic study for these protein-induced DNA structural changes. For instance, there were a few attempts to determine the linking number change ( $\Delta Lk$ ) induced by LacI; however, the results were inconsistent (19,20). This controversy may be caused by the fact that each LacI causes a small  $\Delta Lk$  for each lac operator [much  $<1$  (19)], it is experimentally not feasible to use a circular plasmid DNA containing one or two lac operators for these studies. In this case, it is required to construct a plasmid DNA template containing multiple tandem copies of the lac operator to determine LacI-induced  $\Delta Lk$ . As demonstrated by Sadler *et al.* (21), it was difficult to construct a stable plasmid containing

\*To whom correspondence should be addressed. Tel: +1 305 348 3277; Fax: +1 305 348 3772; Email: lengf@fiu.edu

many tandem copies of the *lac O*<sub>1</sub> operator, which made the determination of  $\Delta Lk$  challenging. In this article, we overcame this difficulty and constructed a series of circular plasmid DNA templates that carry many tandem copies of one DNA-binding site of a sequence-specific DNA-binding protein, such as LacI, GalR, CRP, AraC and  $\lambda$  O protein. Utilizing these unique plasmid DNA templates, we were able to simultaneously determine the protein-induced  $\Delta Lk$  and the DNA-binding constant for  $\lambda$  O protein and GalR. We also found that the protein-induced  $\Delta Lk$  is a unique property for each DNA-binding protein and does not correlate with the protein-induced DNA bending. In addition, protein-mediated DNA looping by AraC and LacI can introduce  $\Delta Lk$  to the DNA-binding sites.

## MATERIALS AND METHODS

### Purified proteins

Bacteriophage  $\lambda$  O protein was a generous gift of R. McMacken (Johns Hopkins University). *Escherichia coli* GalR was kindly provided by S. Adhya (National Institutes of Health). *Escherichia coli* CRP was a gift of J. C. Lee (The University of Texas Medical Branch, Galveston, TX, USA). *Escherichia coli* AraC protein is a gift of R. Schleif (Johns Hopkins University). *Escherichia coli* LacI was purified by the method of Chen and Matthews [(22); *E. coli* strains containing plasmids overexpressing these LacI mutants were kindly provided by K. S. Matthews at Rice University]. All restriction enzymes, T4 DNA ligase and calf intestinal alkaline phosphatase were purchased from New England Biolabs (Beverly, MA, USA). Recombinant Human topoisomerase I was obtained from TopoGen, Inc. (Port Orange, FL, USA).

### Plasmid DNA templates

All plasmids are derived from low copy number plasmids pACYC184 or pACYCDuet-1. Plasmid pYZX43 was constructed by the insertion of a 24-bp synthetic DNA fragment, containing a BglII site, into the unique BspHI site of pACYC184. A 323-bp BamHI–BglII DNA fragment carrying phage  $\lambda$  sequences from 38974 to 39207, which includes the  $\lambda$  replication origin with four iterons (O protein-recognition sites), was cloned into the BamHI–BglII sites of pYZX43 to create pCB2. Next, the same 323-bp BamHI–BglII DNA fragment was inserted into the unique BglII site of pCB2 to produce pCB3 that contains two tandem copies of the  $\lambda$  replication origin. Plasmid pCB5 was made by the insertion of a 646-bp BamHI–BglII fragment, which contains two tandem copies of the  $\lambda$  replication origin, into the unique BglII site of pCB3. In this case, pCB5 contains four tandem copies of the  $\lambda$  replication origin, which has 16 O-recognition sites. Plasmid pCB21 was constructed by inserting a 145-bp DNA fragment, containing five tandem copies of iteron III of the  $\lambda$  replication origin, into the BamHI–BglII sites of pYZX43. The 145-bp

fragment is the head-to-tail ligation product of a 29-bp synthetic BamHI–BglII DNA fragment (top strand: 5'-GATCCCTCAAATTGGGGGATTGCGCTGAA-3'; the underlined sequence is iteron III of  $\lambda$  replication origin). Since the O-recognition sequence iteron III is a 19-bp DNA sequence, the space between each iteron III is 10 bp. Plasmid pCB22 was produced by the insertion of the same 145-bp DNA fragment into the unique BglII site of pCB21 and contains 10 tandem copies of iteron III. Plasmid pCB28 was constructed by inserting a 290-bp BamHI–BglII fragment into the unique BglII site of pCB22. In this case, pCB28 contains 20 copies of head-to-tail iteron III multimers (20 O-recognition sites). Using similar approaches, the following plasmids have been made: pCB18, pCB32, pCB34 and pCB37, which contain 15, 20, 24 and 16 copies of head-to-tail iteron III multimers, respectively. The space between each iteron III is 5, 15, 20 and 25 bp for these plasmids, respectively (Table 1). These plasmids were used to examine the phase and distance effect on the O-protein-induced  $\Delta Lk$  or  $\beta$ .

Plasmids pCB42 and pCB46 were constructed by the insertion, respectively, of a 648 or 738 bp BamHI–BglII DNA fragment into the BamHI–BglII sites of pYZX43. Both plasmids contain 18 tandem copies of the *gal O*<sub>E</sub> operator. The difference between these two plasmids is the space between each *gal O*<sub>E</sub> operator, which is 20 and 25 bp, respectively (the head-to-tail lengths for each repeating sequence are 36 and 41 bp, respectively). Plasmids pCB51 and pCB55 were made by insertion of an 1152 or 1272 bp BamHI–BglII DNA fragment into the BamHI–BglII site of pYZX43, respectively. Both plasmids contain 24 tandem copies of the CRP binding site of the *lac P*<sub>1</sub> promoter. The difference between these two plasmids is also the space between each neighboring CRP sites, which is 24 and 29 bp, respectively. Plasmid pYZX12 was constructed by the insertion of a 22-bp DNA fragment, containing a nicking restriction endonuclease Nt.BbvC1 site, into the unique BstEII site of pACYCDuet-1. A 476-bp XhoI fragment carrying seven copies of the head-to-tail araI site (a head-to-tail ligation product of a 68 bp synthetic XhoI DNA fragment containing one araI site of the L-arabinose operon) was then cloned into the unique XhoI site of pYZX12 to yield plasmid pYZX36. Plasmid pYZX42 was constructed by the insertion of a 588 bp EcoRV–AvrII fragment, containing the seven tandem copies of the araI site of pYZX36, between the BsaAI and NheI sites of pYZX36. In this case, pYZX42 contains 14 araI sites.

Plasmid pYZX46 was constructed in a few steps. In the first step, a synthetic oligonucleotide (92 bp; annealing product of oligonucleotides FL313 and FL314) containing two *lac O*<sub>1</sub> operators was inserted into the unique BamHI site of pYZX12 to produce pYZX29, which contains four *lac O*<sub>1</sub> operators (pACYCDuet-1 has two *lac O*<sub>1</sub> operators). Using similar approaches, we have made the following plasmids: pYZX34 (six *lac O*<sub>1</sub> operators), pYZX37 (eight *lac O*<sub>1</sub> operators) and pYZX40 (10 *lac O*<sub>1</sub> operators). Next, pYZX40 was digested by BamHI and BglII to produce 567, 790 and 3041 bp fragments. The 567-bp

**Table 1.** Plasmids constructed in this study

Plasmid	DNA-binding site	Space between each DNA-binding site (bp)	Copy number of the DNA-binding site	The DNA-binding sequences (top strand)
pCB5	O binding site	4	16	$\lambda$ DNA replication origin <sup>a</sup>
pCB18	Iteron III <sup>b</sup>	5	15	5'-ATCCCTCAAATTGGGGGAT-3'
pCB28	Iteron III <sup>b</sup>	10	20	5'-ATCCCTCAAATTGGGGGAT-3'
pCB32	Iteron III <sup>b</sup>	15	20	5'-ATCCCTCAAATTGGGGGAT-3'
pCB34	Iteron III <sup>b</sup>	20	24	5'-ATCCCTCAAATTGGGGGAT-3'
pCB37	Iteron III <sup>b</sup>	25	16	5'-ATCCCTCAAATTGGGGGAT-3'
pYZX36	<i>araI</i>	30	7	5'-TAGCATTTTTATCCATAAGATTAGCGGATCCTACCTGA-3'
pYZX42	<i>araI</i>	30	14	5'-TAGCATTTTTATCCATAAGATTAGCGGATCCTACCTGA-3'
pCB42	<i>gal O<sub>E</sub></i>	20	18	5'-GTGTAACGATTCCAC-3'
pCB46	<i>gal O<sub>E</sub></i>	25	18	5'-GTGTAACGATTCCAC-3'
pCB51	The CRP binding site <sup>c</sup>	20	24	5'-CAATTAATGTGAGTTAGCTCACTCATTA-3'
pCB55	The CRP binding site <sup>c</sup>	25	24	5'-CAATTAATGTGAGTTAGCTCACTCATTA-3'
pYZX46	<i>lac O<sub>1</sub></i>	25	19	5'-AATTGTGAGCGGATAACAATT-3'

<sup>a</sup> $\lambda$  DNA origin contains four DNA-binding sites for  $\lambda$  O protein.

<sup>b</sup>Iteron III is a DNA-binding site of  $\lambda$  O protein from  $\lambda$  DNA replication origin (the third repeating sequence of  $\lambda$  DNA replication origin).

<sup>c</sup>The CRP binding site is derived from the CRP binding site of *E. coli lac P<sub>1</sub>* promoter.

fragment, which contains nine *lac O<sub>1</sub>* operators, was then inserted into the BamHI sites of pYZX40 to generate pYZX46 that contains 19 head-to-tail tandem copies of *lac O<sub>1</sub>* operators.

Plasmids pBend2- $\lambda$ O, pBend2-*lacO<sub>1</sub>*, pBend2-*galO<sub>E</sub>* and pBend2-CRP were created by insertion, respectively, of a synthetic XbaI DNA fragment containing either iteron III of  $\lambda$  DNA replication origin or *lac O<sub>1</sub>* operator or *gal O<sub>E</sub>* operator or the CRP binding site of the *lac P<sub>1</sub>* promoter into the unique XbaI site of pBend2 [plasmid pBend2 is a generous gift of S. Adhya (National Institutes of Health)].

All DNA elements were confirmed by DNA sequencing.

### The determination of protein-induced DNA $\Delta Lk$ or unwinding angle ( $\beta$ )

In order to determine the protein-induced  $\Delta Lk$ , we used plasmids containing multiple tandem copies of a DNA-binding site of  $\lambda$  O protein, GalR, CRP, AraC or LacI (Table 1). Two methods were used: the ligation method and the Topo I (topoisomerase I) method. Since these plasmids each contain one nicking enzyme recognition site (Nb.BsrDI or Nt.BbvCI), in the ligation method, we nicked these plasmids using one of the nicking restriction enzymes, Nb.BsrDI or Nt.BbvCI, and then titrated the nicked plasmid with one of the DNA-binding proteins, i.e.  $\lambda$  O protein, GalR, CRP, AraC or LacI. After 30 min incubation, the DNA templates were ligated by T4 DNA ligase. The linking number (Lk) of the ligated DNA products was determined with 1% agarose gel electrophoresis in the presence of chloroquine (23) and calculated from the gel images stained with ethidium bromide or SYBR Gold using KODAK 1D Image Analysis Software. In the Topo I method, the supercoiled plasmid DNA templates were titrated with one of the DNA-binding proteins and then relaxed by human topoisomerase I. The relaxation reaction was stopped by phenol extraction. The linking number of the DNA templates was also determined with 1% agarose gel

containing chloroquine as described above. The observed linking number change was calculated by  $\Delta Lk_{\text{obs}} = Lk - Lk^{\circ}$ , where Lk and  $Lk^{\circ}$  are the Boltzmann centers of the topoisomers band in the presence of the DNA-binding protein and in its absence, respectively. If we define the apparent protein-induced DNA-unwinding angle ( $\beta_{\text{obs}}$ ) equals to  $360 \times \Delta Lk_{\text{obs}}$ , i.e.  $\beta_{\text{obs}} = 360 \times \Delta Lk_{\text{obs}}$ , then the protein-induced DNA-unwinding angle ( $\beta$ ) per binding site was determined by the following equation if we assume that no interaction exists among the bound DNA-binding protein and context-dependent differences in binding affinity are ignored (for proteins cooperatively binding to its DNA-binding sites, more complex equations have to be derived) and the DNA-binding density ( $r$ ) is equal to  $\beta_{\text{obs}}/\beta$  (see below for its derivation):

$$\frac{\beta_{\text{obs}}}{C_f} = K(n\beta - \beta_{\text{obs}}) \quad (1)$$

where  $K$  is the DNA-binding constant,  $n$  is the binding site size ( $n$  is the copy number of the DNA-binding sites on the plasmids that can be determined by DNA sequencing) and  $C_f$  is the free concentration of the DNA-binding protein (for the low protein concentrations, the free protein concentration was estimated using the DNA-binding affinity of the protein. For the high-protein concentrations,  $C_f$  is approximated to the total ligand concentration  $C_T$ ). This analysis should give us the protein-induced  $\beta$  and  $K$  simultaneously. The intrinsic  $\Delta Lk$  per binding site can be calculated by

$$\Delta Lk = \frac{\beta}{360} \quad (2)$$

To derive Equation 1, we consider a binding reaction between a sequence-specific DNA-binding protein and  $n$  independent binding sites on a plasmid DNA template with an intrinsic binding constant of  $K$ . Each of the  $n$  sites has the same affinity for the DNA-binding protein as any other and the binding is non-cooperative.

The binding properties can be described by Scatchard Equation:

$$\frac{r}{C_f} = K(n - r) \quad (3)$$

where  $r$  is the binding density (the number of moles of the DNA-binding protein bound per mole of the DNA template) and  $C_f$  is the concentration of the DNA-binding protein in the free form. If the DNA-binding protein binds to each of its  $n$  binding sites to cause the unwinding of  $\beta$  degrees of the DNA template, the DNA-binding density  $r$  can be calculated by

$$r = \frac{\beta_{\text{obs}}}{\beta} \quad (4)$$

Substitution of Equation 4 into Equation 3 yields Equation 1.

### The determination of the protein-induced DNA bending angle

The protein-induced DNA bending angles were determined using the circular permutation assay (24). Plasmids pBend2 derivatives containing a recognition sequence of a sequence-specific DNA-binding protein, such as  $\lambda$  O, GalR, CRP and LacI were digested by restriction enzymes MluI, BglII, NheI, SpeI, XhoI, DraI, EcoRV, NruI and BamHI to produce a set of fragments with identical length and base composition in which the position of the protein binding site is variable. The DNA fragments were labeled with  $^{32}\text{P}$  at 5'-termini by T4 polynucleotide kinase in the presence of  $\gamma\text{-}^{32}\text{P}\text{-ATP}$ . The DNA-protein complexes were formed by addition of  $1.0$  to  $3.0 \times 10^{-9}$  M of the protein to a solution containing  $1 \times 10^{-10}$  M of  $^{32}\text{P}$ -labeled DNA. After equilibration for 60 min at  $22^\circ\text{C}$ , the samples were loaded on a polyacrylamide gel to determine the mobility of DNA-protein complexes, which is dependent upon the position of the bound protein, with the lowest mobility present when the protein is bound to the center of the fragment. The bending angle  $\alpha$  by which the DNA is bent from linearity was estimated by

$$\text{Cos}\left(\frac{\alpha}{2}\right) = \frac{\mu_M}{\mu_E} \quad (5)$$

where  $\mu_M$  and  $\mu_E$  are the mobility of the complex with protein bound at the center and the end of DNA, respectively (25).

### Electrophoretic mobility shift assay

Electrophoretic mobility shift assays (EMSA) were used to determine the apparent DNA-binding constant of  $\lambda$  O protein, CRP, GalR and LacI. DNA oligomers containing a DNA-binding site of one of the DNA-binding proteins (the EcoRV fragment of the pBend2 derivatives) were labeled with  $^{32}\text{P}$  at 5'-termini by T4 polynucleotide kinase in the presence of  $\gamma\text{-}^{32}\text{P}\text{-ATP}$ . The DNA-protein complexes were formed by addition of appropriate amounts of the protein to a solution containing  $0.1$  nM of  $^{32}\text{P}$ -labeled DNA in the  $1 \times$  DNA-binding buffer

containing 20 mM Tris-HCl (pH 8.0), 200 mM NaCl, 0.5 mM EDTA, 1 mM DTT, 5 mM  $\text{MgCl}_2$  and 5% glycerol. After equilibration for 60 min at  $22^\circ\text{C}$ , the samples were loaded on an 8% native polyacrylamide gel in  $0.5 \times$  TBE buffer [0.045 M Tris-borate (pH 8.3) and 1 mM EDTA] to separate free and bound DNA. The gels were subsequently dried and visualized by autoradiography or quantified using a Fuji FLA 3000 image analyzer. The radioactivity of the free and bound DNA was determined and used to calculate the binding ratio ( $R$ ), which is equal to the ratio of the radioactivity of the bound DNA divided by the sum of the radioactivity of the bound and free DNA. The apparent DNA-binding constant ( $K_{\text{app}}$ ) was obtained by nonlinear-least-squares fitting the following equation using the program Scientist.

$$R = \frac{(a + x + 1/K_{\text{app}}) - \sqrt{(a + x + 1/K_{\text{app}})^2 - 4ax}}{2a} \quad (6)$$


where  $a$  and  $x$  represent the total DNA and the total protein concentration, respectively.

## RESULTS

### A strategy to study protein-induced DNA $\Delta Lk$ of circular plasmid DNA by sequence-specific DNA-binding proteins

Here, we successfully made a series of DNA plasmids containing many tandem copies of one DNA-binding site, such as *lac*  $O_1$  operator, *gal*  $O_E$  operator, the CRP binding site of *lac*  $P1$  promoter, *araI* site of the L-arabinose operon and iteron III of phage  $\lambda$  DNA replication origin (a DNA-binding site of  $\lambda$  O protein). Our cloning strategy was to use a low-copy number plasmid pACYC184 or pACYCDuet-1 to minimize the recombination among the tandem copies of the DNA-binding sequences. In this case, plasmids carrying multiple copies of one DNA-binding site are stable in *E. coli* strains, such as DH5 $\alpha$  and Top10. We also cloned one nicking endonuclease recognition site (either Nt.BbvCI or Nb.BsrDI) into these plasmids. The distance between the nicking endonuclease recognition site and the DNA-binding sites is  $\sim 0.8$  or  $1.8$  kb (Figure 1) to minimize the interference between the specific binding of the DNA-binding proteins to the multiple DNA-binding sites and the ligation reaction for determining the protein-induced  $\Delta Lk$  (see below for details). Figure 1 and Table 1 summarize the main properties of the plasmids.

As described in 'Materials and Methods' section, we employed two methods, a ligation method and a Topo I method to determine the protein-induced  $\Delta Lk$  by sequence-specific DNA-binding proteins. Since  $\Delta Lk$  of  $-1$  is equivalent to an unwinding angle of  $360^\circ$  for a circular plasmid DNA template, we chose to use the unwinding angle ( $\beta$ ) for our calculation, i.e.  $\beta = 360 \times \Delta Lk$ . If the concentration of the DNA template is sufficiently low ( $\ll K_d$ ) and the interaction between individual DNA-binding proteins is negligible, we should be able to simultaneously determine  $\beta$  or  $\Delta Lk$  and the DNA-binding constant ( $K$ ). Since the initial




**Figure 1.** Plasmids used to determine protein-induced  $\Delta Lk$  by  $\lambda$  O protein, GalR, CRP, AraC and LacI. Plasmids pCB34, pCB46 and pCB55 were derived from pACYC184; plasmids pYZX36, pYZX42 and pYZX46 were derived from pACYCDuet-1. These plasmids were constructed as detailed under 'Materials and Methods' section. The restriction enzyme sites for BamHI, BglIII, XhoI, AvrII, Nb.BsrDI and Nt.BbvCI are shown. The small closed rectangles represent the site specific recognition sequence for  $\lambda$  O protein (Iteron III), GalR (*gal O<sub>E</sub>*), CRP, AraC (*araI*) and LacI (*lacO<sub>1</sub>*).

topological status of the DNA templates is quite different for these two methods,  $\beta$  or  $\Delta Lk$  determined by these two methods may also be different (see below for details). Our strategy is summarized in Supplementary Figure S1.


### Simultaneous determination of protein-induced $\Delta Lk$ or $\beta$ and the DNA-binding constant of $\lambda$ O protein and GalR

Due to availability, we first decided to determine  $\lambda$  O protein-induced  $\Delta Lk$  or  $\beta$ .  $\lambda$  O protein is the DNA replication initiator of bacteriophage  $\lambda$  and specifically binds to the four repeating sequences (iterons) of  $\lambda$  DNA replication origin (26). Upon  $\lambda$  O binding, the  $\lambda$  DNA replication origin is wrapped around O protein to form a nucleoprotein complex, the 'O-some' (6) and subsequently unwind the DNA sequence of  $\lambda$  DNA replication origin (27). In this case,  $\lambda$  O protein should induce a  $\Delta Lk$  to the DNA templates containing multiple copies of its binding sites. Figure 2A shows results of  $\lambda$  O protein titrating into a solution containing 0.0625 nM of the nicked plasmid pCB5 carrying four  $\lambda$  replication origins (16  $\lambda$  O protein binding sites; the concentration of the  $\lambda$  O protein binding site is 1 nM). After the ligation reaction, the plasmid DNA samples were subjected to electrophoresis in a chloroquine-agarose gel to determine the linking number (Lk). In the absence of  $\lambda$  O protein, the DNA template is relaxed with a superhelical density,  $\sigma$ , of approximately zero. In the presence of the DNA intercalator chloroquine (0.5  $\mu$ g/ml), this DNA migrated during agarose gel electrophoresis as if it contained a few (+) supercoils



**Figure 2.** Simultaneous determination of  $\lambda$  O protein-induced DNA unwinding angle ( $\beta$ ) or  $\Delta Lk$  and the DNA-binding constant ( $K$ ). (A) and (C)  $\lambda$  O protein titration assays were performed as described under 'Materials and Methods' section using the ligation method (A) or the Topo I method (C). In addition to 0.0625 nM of plasmid pCB5, the reaction mixtures for the DNA samples applied to lanes 1–10 also contain 0, 1.5, 2.0, 3.0, 4.0, 5.0, 6.3, 7.5, 10.0, 12.5 nM of  $\lambda$  O protein (as dimer), respectively. DNA topoisomers were resolved with electrophoresis in 1% agarose gel containing 0.5  $\mu$ g/ml chloroquine. (B) and (D) Quantification analysis of the binding data from the  $\lambda$  O titration experiments.  $\beta$  and  $K$  were calculated according to Equation 1.  $\Delta Lk$  was calculated by Equation 2.

(Figure 2A, lane 1). Binding of  $\lambda$  O protein to the  $\lambda$  O binding sites on plasmid pCB5 introduced about 4 (–) supercoils into the DNA template (Figure 2A, lanes 1–10). The  $\lambda$  O-induced  $\Delta Lk$  or  $\beta$  required the presence of



**Figure 3.** The Determination of the DNA-binding constant ( $K$ ) and the protein-induced DNA bending angle for  $\lambda$  O protein. The EMSA assays and the DNA bending assays were performed as described under ‘Materials and Methods’ section. (A) Binding of  $\lambda$  O protein to the DNA oligomer containing the iteron III of coliphage  $\lambda$  DNA replication origin. A 145 bp  $^{32}\text{P}$ -labeled EcoRV fragment of pBend2-Iteron III was incubated with increasing concentrations of  $\lambda$  O protein in 50  $\mu\text{l}$  of  $1\times$  EMSA binding buffer containing 10 mM Tris-HCl (pH 7.4), 20 mM KCl, 0.5 mM EDTA, 1 mM DTT, 5 mM  $\text{MgCl}_2$  and 5% glycerol. The autoradiogram was shown. The radioactivities were quantified with a PhosphorImager. Lane 1 is free DNA fragment. In addition to the  $^{32}\text{P}$ -labeled DNA fragment, the reaction mixtures for the DNA samples applied to lanes 2–11 also contains 0, 1, 2.5, 5, 7.5, 10, 15, 20, 25, 50 and 100 nM of  $\lambda$  O protein (as dimer), respectively. (B) Quantification analysis of the binding data from the EMSA experiments. The bound ratio of DNA was plotted against the protein concentration. The curve was generated by fitting the data to Equation 6 as described under ‘Materials and Methods’ section to yield the DNA-binding constant of  $2.5\times 10^8 \text{M}^{-1}$ . (C) The DNA bending assay. After the binding of  $\lambda$  O protein to the permuted DNA fragments, an 8% polyacrylamide gel was used to separate the bound and free DNA fragments. The autoradiogram of  $^{32}\text{P}$ -labeled DNA fragments was shown. Lanes 1–5 are the  $\lambda$  iteron III fragments produced by digestion of plasmid pBend2-Iteron III with restriction enzymes of MluI, NheI, EcoRV, SspI and BamHI, respectively. The DNA fragments in the bottom of the gels are free DNA, and those in the upper part are protein–DNA complexes.

O binding sites in the DNA templates (Figure S2). When plasmid pYZX43, which is identical to pCB5 except that it does not contain  $\lambda$  O binding sites, was used in the ligation method,  $\lambda$  O protein was not able to induce a  $\Delta Lk$  or  $\beta$  to the plasmid (Figure S2A). The results in Figure 2 were used to fit Equation 1 to yield the  $\lambda$  O protein-induced  $\beta$  ( $80.0^\circ \pm 5.6^\circ$ ) or  $\Delta Lk$  ( $-0.222 \pm 0.016$ ) and the DNA-binding constant [ $3.0(\pm 1.5)\times 10^8 \text{M}^{-1}$ ] (Figure 2B). The DNA-binding constant of  $\lambda$  O protein is almost identical to the one determined with EMSA assay (Figure 3). Similar results were also obtained using the Topo I assay (Figure 2C and D; Table 2).

We next varied the distance between the individual O binding sites (iteron III of  $\lambda$  replication origin) to examine whether the phase match or the distance between these DNA-binding sites affects the induced  $\Delta Lk$  or  $\beta$ . To our surprise, O protein-induced  $\Delta Lk$  or  $\beta$  is independent of the phase match and the distance between these DNA-binding sites (Table 2). Again, the DNA-binding constant was determined to be  $\sim 2.0\text{--}3.0\times 10^8 \text{M}^{-1}$ , which is consistent with the EMSA results (Figure 3) and the published results (2). However, some variations in the DNA-binding constants are observed in these assays.

We also determined GalR-induced  $\Delta Lk$  or  $\beta$ . GalR is a dimer and specifically binds to the  $O_E$  and  $O_I$  operators of *E. coli* galactose operon to inhibit transcription from two *gal* promoters *P1* and *P2* (28). As described in ‘Materials and Methods’ section, we made two plasmids pCB42 and pCB46 that carry 18 *gal*  $O_E$  operators (Figure 1 and Table 1). The difference of these two plasmids is the space between the neighboring  $O_E$  operators: pCB42, 20 bp and pCB46, 25 bp (the head-to-tail distance of the  $O_E$  operators are 36 and 41 bp, respectively). Since the  $O_E$  operator is a 16 bp DNA sequence, it is reasonable to assume that GalR binds to the neighboring  $O_E$  operators of pCB42 on the opposite side; in contrast, GalR binds to the neighboring  $O_E$  operators of pCB46 on the same side. In this case, we were able to examine whether the phase match affects the induced  $\Delta Lk$  or  $\beta$ . Our results are summarized in Figure 4 and Table 3. For the ligation method, the binding of GalR to both plasmids pCB42 and pCB46 induced about 3 (–) supercoils into the


**Table 2.**  $\lambda$  O protein-induced DNA-unwinding angle ( $\beta$ ),  $\Delta Lk$  and DNA-binding constant ( $K$ )

Plasmid	The space between each DNA-binding site (bp)	Copy number	$\beta$ ( $^\circ$ )	$\Delta Lk$	$K$ ( $\times 10^8 \text{M}^{-1}$ )
pCB5 <sup>a</sup>	4	16	$80.0 \pm 5.6$	$-0.222 \pm 0.016$	$3.0 \pm 1.5$
pCB5 <sup>b</sup>	4	16	$79.1 \pm 1.5$	$-0.220 \pm 0.004$	$3.6 \pm 0.5$
pCB18 <sup>a</sup>	5	15	$79.4 \pm 1.4$	$-0.221 \pm 0.004$	$1.9 \pm 0.4$
pCB28 <sup>a</sup>	10	20	$80.0 \pm 1.8$	$-0.222 \pm 0.005$	$3.9 \pm 0.7$
pCB32 <sup>a</sup>	15	20	$80.8 \pm 1.8$	$-0.224 \pm 0.005$	$1.5 \pm 0.3$
pCB34 <sup>a</sup>	20	24	$80.0 \pm 1.3$	$-0.222 \pm 0.004$	$2.3 \pm 0.3$
pCB37 <sup>a</sup>	25	16	$79.5 \pm 1.3$	$-0.221 \pm 0.004$	$1.9 \pm 0.4$

<sup>a</sup>The protein-induced  $\Delta Lk$ ,  $\beta$  and the DNA-binding constant was determined by the ligation method as described under ‘Materials and Methods’ section.

<sup>b</sup>The protein-induced  $\Delta Lk$ ,  $\beta$  and the DNA-binding constant was determined by the Topo I method as described under ‘Materials and Methods’ section.

DNA templates (Figure 4A, lanes 1–10). Control experiments indicated that GalR-induced  $\Delta Lk$  or  $\beta$  was specific for the GalR– $O_E$  protein–DNA interaction. GalR failed to induce DNA supercoiling when its inducer, D-galactose, was present (Figure 4B) or the plasmid template does not contain the gal operators (Figure S2B). These results were used to fit Equation 1 to yield the GalR-induced  $\beta$  ( $60.9^\circ \pm 1.5^\circ$ ) or  $\Delta Lk$  ( $-0.169 \pm 0.004$ ) and the



**Figure 4.** Simultaneous determination of GalR-induced DNA-unwinding angle ( $\beta$ ) or  $\Delta Lk$  and the DNA-binding constant ( $K$ ). GalR titration assays were performed as described in ‘Materials and Methods’ section using the ligation method. DNA topoisomers were resolved with electrophoresis in 1% agarose gel containing 0.5  $\mu\text{g/ml}$  chloroquine. (A) The GalR titration experiment. In addition to 0.055 nM of plasmid pCB42, the reaction mixtures for the DNA samples applied to lanes 1–10 also contain 0, 1.3, 1.5, 2.0, 2.5, 3.0, 4.0, 6.0, 8.0, 10.0 nM of GalR (as dimer), respectively. (B) GalR failed to unwind the DNA when D-galactose was present. The reaction mixture for the DNA sample applied to lane 1 contains neither GalR nor D-galactose. As indicated at the top of the image, the reaction mixtures for the DNA samples applied to lanes 2–4 contain GalR (lanes 2 and 4, 5 nM; lanes 3 and 5, 10 nM) and D-galactose (lanes 4 and 5, 1 mM). (C) Quantification analysis of the binding data from GalR titration experiments.  $\beta$  and  $K$  were determined according to Equation 1.  $\Delta Lk$  was calculated by Equation 2.

DNA-binding constant of  $3.2(\pm 1.5) \times 10^8 \text{ M}^{-1}$  (Figure 4C and Table 3). Interestingly, there is no difference for the  $\beta$  or  $\Delta Lk$  values between these two plasmids, indicating that phase match does not influence GalR-induced  $\Delta Lk$  or  $\beta$ . Similar results were obtained by using the Topo I method for GalR-induced  $\Delta Lk$  or  $\beta$  (Table 3).

#### $\Delta Lk$ or $\beta$ induced by DNA looping proteins AraC and LacI

Since DNA loops are topological structures, we decided to examine whether protein-mediated DNA looping affects protein-induced  $\Delta Lk$  or  $\beta$ . For this purpose, we chose AraC and LacI, two well-characterized DNA-binding proteins that cause DNA looping *in vivo* and *in vitro* (29). In the absence of L-arabinose, AraC, a homodimer, binds *in trans* to  $I_1$  and  $O_2$  half-sites of the L-arabinose operon to form a DNA loop and therefore inhibits transcription from the  $P_{BAD}$  promoter (8). In the presence of L-arabinose, AraC also tightly binds *in cis* to the neighboring  $I_1$  and  $I_2$  half-sites; however, it does not form a DNA loop. In this case, AraC stimulates transcription from the  $P_{BAD}$  promoter (8). LacI, a homotetramer, forms DNA loops through specifically binding to the *lac* operators, such as  $O_1$ ,  $O_2$  and  $O_3$ , to negatively control gene transcription of *E. coli lac* operon (30). To study AraC-induced  $\Delta Lk$  or  $\beta$ , we constructed two plasmids pYZX36 and pYZX42 (Figure 1). pYZX36 and pYZX42 contain 7 and 14 head-to-tail *araI* sites, respectively. The 14 *araI* sites of plasmid pYZX42 are located in two different locations with seven *araI* sites for each location. The distance between these two groups of *araI* sites is 1.2 kb on one side and 2.7 kb on the other side. In this situation, AraC should be able to induce DNA looping between these two groups of *araI* sites (31). As expected, AraC did not induce significant DNA supercoiling for plasmid pYZX36. Nevertheless, AraC greatly supercoiled plasmid pYZX42, and the binding of AraC to *araI* sites on plasmid pYZX42 introduced about 5 (–) supercoils into the DNA template (compare lane 1 to lane 2 of Figure 5A). Interestingly, as the AraC concentration was increased, the plasmid DNA template became less supercoiled (Figure 5A). Our interpretation is that AraC-mediated DNA looping is the cause for AraC-induced  $\Delta Lk$  or  $\beta$ . Since we only used 1 nM of the *araI* site for the assay, a low concentration of AraC,

**Table 3.** GalR protein-induced DNA-unwinding angle ( $\beta$ ),  $\Delta Lk$  and DNA-binding constant ( $K$ )

Plasmid	The space between the individual DNA-binding site (bp)	Copy number	$\beta$ ( $^\circ$ )	$\Delta Lk$	$K$ ( $\times 10^8 \text{ M}^{-1}$ )
pCB42 <sup>a</sup>	20	18	$60.9 \pm 1.5$	$-0.169 \pm 0.004$	$4.4 \pm 1.7$
pCB42 <sup>b</sup>	20	18	$102.6 \pm 0.9$	$-0.285 \pm 0.003$	$7.2 \pm 0.5$
pCB46 <sup>a</sup>	25	18	$60.0 \pm 0.9$	$-0.167 \pm 0.003$	$3.8 \pm 1.2$
pCB46 <sup>b</sup>	25	18	$66.5 \pm 3.8$	$-0.185 \pm 0.011$	$7.7 \pm 0.9$

<sup>a</sup>The protein-induced  $\Delta Lk$ ,  $\beta$  and the DNA-binding constant was determined by the ligation method as described under ‘Materials and Methods’ section.


<sup>b</sup>The protein-induced  $\Delta Lk$ ,  $\beta$  and the DNA-binding constant was determined by the Topo I method as described under ‘Materials and Methods’ section.



e.g. 2.5 or 5 nM, should favor the binding of the protein *in trans* to form a DNA loop. In this scenario, AraC greatly supercoiled the DNA template. In contrast, a high concentration of AraC was in favor of the binding of the dimeric AraC to a single half site without forming a DNA loop, i.e. the DNA loop was titrated out. In this case, AraC should not significantly supercoil the DNA template. Our explanation was confirmed by the results shown in Figure 5B in which AraC failed to induce  $\Delta Lk$  or  $\beta$  when L-arabinose was present in the reaction mixtures (compare lanes 2 and 3 with lanes 4 and 5 of Figure 5B). These results suggest that AraC-mediated DNA looping is the reason for AraC-induced  $\Delta Lk$  or  $\beta$  under our experimental conditions. Because AraC-mediated DNA looping is a cooperative process, we cannot use Equation 1 to derive AraC-induced  $\Delta Lk$  or  $\beta$  and the DNA-binding constant. However, with an assumption of AraC binding

to all 14 *araI* sites of pYZX42, we estimated the maximum  $\Delta Lk$  and  $\beta$  induced by AraC to be  $-0.333 \pm 0.039$  and  $120 \pm 14^\circ$ , respectively (Table 4). Similar results were also obtained using the Topo I assay. The  $\Delta Lk$  and  $\beta$  induced by AraC may be different if a different plasmid containing multiple AraC binding sites is used.

We also made a few plasmids to study LacI-induced  $\Delta Lk$  or  $\beta$ . Among them is pYZX46 that carries 19 *lac O*<sub>1</sub> operators (Figure 1 and Table 1). The 19 *lac O*<sub>1</sub> operators are divided into two different locations with each location containing 9 or 10 *lac O*<sub>1</sub> operators. The distance between these two locations is 224 bp (Figure 1). According to previous results (32), LacI should induce a DNA loop between these two groups of *lac O*<sub>1</sub> operators. In this case, it should also induce DNA supercoiling. Figure 6A and B show results of the DNA supercoiling assay (the ligation method) by LacI in which the DNA concentration was kept sufficiently low to minimize the nonspecific DNA-binding. As expected, LacI induced 2–3 (–) supercoils into the DNA templates (compare lane 1 to lanes 2–4 of Figure 6A and B), which is consistent with the published results (20). Interestingly, although *lac O* operators are required for LacI-induced  $\Delta Lk$  or  $\beta$  (Figure S2D), IPTG inhibited the LacI-induced DNA supercoiling only when the concentration of both LacI and *lac O*<sub>1</sub> operator was very low (compare lane 2 with lane 5 of Figure 6A). When LacI's concentration and/or *lac O*<sub>1</sub>'s concentration was increased, IPTG stimulated LacI-induced DNA supercoiling (compare lanes 6 and 7 with lane 2 of Figure 6A; also compare lanes 5–7 with lanes 2–4 of Figure 6B). We believe that the stimulation of LacI-induced DNA supercoiling by IPTG stems from the LacI-mediated DNA looping which has a different topology. As demonstrated before (20,32), LacI, even in the presence of IPTG, is still able to induce DNA looping. We demonstrated here that the topology of the DNA loops in the presence and absence of IPTG is different. We estimated the maximum  $\Delta Lk$  induced by LacI in the absence or presence of IPTG are, respectively,  $-0.08$  ( $\beta$ ,  $29^\circ$ ) and  $-0.21$  ( $\beta$ ,  $75.8^\circ$ ) (Table 4). We also



**Figure 5.** AraC-induced  $\Delta Lk$ . Experiments to determine AraC-induced  $\Delta Lk$  were performed as described under 'Materials and Methods' section. DNA topoisomers were resolved with electrophoresis in 1% agarose gel containing 0.5  $\mu\text{g/ml}$  chloroquine. (A) The AraC titration experiment. In addition to 0.0714 nM of plasmid pYZX42, the reaction mixtures for the DNA samples applied to lanes 1–11 also contain 0, 2.5, 5.0, 7.5, 10.0, 12.5, 15.0, 20.0, 25.0, 30.0, 35.0 nM of AraC (as dimer), respectively. (B) AraC failed to induce  $\Delta Lk$  of pYZX42 when L-arabinose was present. The reaction mixture for the DNA sample applied to lane 1 contains neither AraC nor L-arabinose. As indicated at the top of the image, the reaction mixtures for the DNA samples applied to lanes 2–5 contain AraC (lanes 2 and 4, 3 nM; lanes 3 and 5, 6 nM) and L-arabinose (lanes 4 and 5, 1 mM).

**Table 4.** Protein-induced DNA-unwinding angle ( $\beta$ ),  $\Delta Lk$ , DNA bending angle ( $\alpha$ ) and DNA-binding constant ( $K$ )


Protein	$\beta$ ( $^\circ$ )	$\Delta Lk$ ( $^\circ$ )	$\alpha$ ( $^\circ$ )	$K$ ( $\text{M}^{-1}$ ) <sup>a</sup>
$\lambda$ O	$80.0 \pm 5.6^b$	$-0.222 \pm 0.016$	$89 \pm 2$	$2.3 \pm 1.7 \times 10^8$
GalR	$60.9 \pm 1.5^b$	$-0.169 \pm 0.004$	$119 \pm 4$	$10.4 \pm 3.6 \times 10^8$
GalR + galactose	$0^b$	0	N/A	N/A
CRP + cAMP	$0^b$	0	$89 \pm 5$	$2.8 \pm 1.1 \times 10^9$
AraC <sup>d</sup>	$120.0 \pm 14.0^b$	$-0.333 \pm 0.039$	90	$5 \times 10^9$
AraC + arabinose <sup>d</sup>	$0^b$	0	90	$2 \times 10^{11}$
LacI	$29.0 \pm 5.8^b$	$-0.081 \pm 0.016$	$73 \pm 4$	$1.2 \pm 0.4 \times 10^{10}$
LacI + IPTG	$75.8 \pm 6.3^b$	$-0.211 \pm 0.018$	N/A	N/A
LacI	$107.5 \pm 5.5^c$	$-0.299 \pm 0.015$	$73 \pm 4$	$1.2 \pm 0.4 \times 10^{10}$
LacI + IPTG	$90.7 \pm 2.7^c$	$-0.252 \pm 0.008$	N/A	N/A

<sup>a</sup>The DNA-binding constants ( $K$ ) were determined by using EMSA as described under 'Materials and Methods' section. The values are the average of at least three independent determinations.

<sup>b</sup>The protein-induced  $\beta$  or  $\Delta Lk$  was determined by the ligation method as described under 'Materials and Methods' section.

<sup>c</sup>The protein-induced  $\beta$  or  $\Delta Lk$  was determined by the Topo I method as described under 'Materials and Methods' section.

<sup>d</sup>The DNA-bending angle of AraC in the absence or presence of L-arabinose was obtained from reference (36). The DNA-binding constant of AraC was obtained from reference (44).




**Figure 6.** The LacI-induced  $\Delta Lk$  in the presence or absence of IPTG. The experiments to determine LacI-induced  $\Delta Lk$  were performed according to the ligation method (A and B) or the Topo I method (C and D) as described under ‘Materials and Methods’ section. Plasmid pYZX46 that contains 19 *lac O*<sub>1</sub> operators was used as the DNA template. The concentration of *lac O*<sub>1</sub> operator and LacI (as tetramer) are shown at the top of the images. The reaction mixtures for the DNA samples applied to lanes 5–7 also contain 1 mM of IPTG.

performed DNA supercoiling assay using the Topo I method. Intriguingly, LacI induced 6 to 7 (–) supercoils into the DNA templates (compare lane 1 with lanes 3 and 4 of Figure 6C), which is 4 more (–) supercoils than those in the ligation method. The induced  $\Delta Lk$  or  $\beta$  was determined to be  $-0.30$  or  $107.5^\circ$ , which is significantly greater than those calculated from the ligation method (Table 4). These results suggest that LacI may ‘keep’ some superhelical energy in the LacI–DNA loop complex.

### $\Delta Lk$ or $\beta$ induced by CRP

Previously, it was shown that *E. coli* cAMP receptor protein (CRP) may cause a small  $\Delta Lk$  upon binding to CRP binding sites (33,34). We therefore decided to study CRP-induced  $\Delta Lk$  or  $\beta$ . Two plasmids pCB51 and pCB55, each containing 24 CRP binding sites, were used for this study (Table 1). As described in ‘Materials and Methods’ section, the difference between these two plasmids is the head-to-tail distance of the neighboring CRP binding sites. Figure 7A shows results of the ligation assay of CRP titrating into a solution containing



**Figure 7.** CRP-induced  $\Delta Lk$ , the DNA-bending angle ( $\alpha$ ), and the DNA-binding constant. (A) CRP titration experiments were performed according to the ligation method as described under ‘Materials and Methods’ section. In addition to 0.0417 nM of plasmid pCB51 and 20  $\mu$ M of cAMP, the reaction mixtures for the DNA samples applied to lanes 1–10 also contain 0, 1.3, 1.5, 2.0, 2.5, 3.0, 4.0, 6.0, 8.0, 10.0 nM of CRP (as dimer), respectively. DNA topoisomers were resolved with electrophoresis in 1% agarose gel containing 0.5  $\mu$ g/ml chloroquine. (B) DNA bending assays were carried out as described under ‘Materials and Methods’ section. After the binding of CRP to the permuted DNA fragments in the presence of 20  $\mu$ M of cAMP, an 8% polyacrylamide gel was used to separate the bound and free DNA fragments. The autoradiogram of <sup>32</sup>P-labeled DNA fragments was shown. Lanes 1–9 are the *lac P*<sub>1</sub> promoter’s CRP fragments produced by digestion of plasmid pBend2-CRP with restriction enzymes of MluI, BglII, NheI, SpeI, XhoI, DraI, EcoRV, NruI and BamHI, respectively. The DNA fragments in the bottom of the gels are free DNA, and those in the upper part are protein–DNA complexes. (C) The EMSA assays were performed as described under ‘Materials and Methods’ section. A 159-bp <sup>32</sup>P-labeled EcoRV fragment of pBEND2-CRP was incubated with increasing concentrations of CRP in 50  $\mu$ l of 1  $\times$  EMSA binding buffer containing 20 mM Tris–HCl (pH 8.0), 20  $\mu$ M cAMP, 200 mM NaCl, 0.5 mM EDTA, 1 mM DTT, 5 mM MgCl<sub>2</sub> and 5% glycerol. The autoradiogram was shown. The radioactivities were quantified with a PhosphorImager. Lane 1 is free DNA fragment. In addition to the <sup>32</sup>P-labeled DNA fragment, the reaction mixtures for the DNA samples applied to lanes 2–11 also contain 0.1, 0.25, 0.5, 1, 1.5, 2.0, 3.0, 5.0, 10.0 and 20.0 nM of CRP (as dimer), respectively. (D) Quantification analysis of the binding data from the EMSA experiments. The bound ratio of DNA was plotted against the protein concentration. The curve was generated by fitting the data to Equation 6 as described under ‘Materials and Methods’ section to yield the DNA-binding constant of  $2.1 \times 10^9 \text{ M}^{-1}$ .


0.0417 nM of plasmid pCB51 (the total concentration of the CRP binding site is 1 nM). To our surprise, CRP in the presence of cAMP did not induce DNA supercoiling under our experimental conditions. Control experiments demonstrated that CRP bound to and bent the CRP binding site (Figure 7B and C). The DNA-binding constant was determined to be  $2.8 \pm 1.1 \times 10^9 \text{ M}^{-1}$  (Figure 7D). These results indicated that CRP-induced  $\Delta Lk$  or  $\beta$  is close to 0 (Table 4). Similar results were also obtained for plasmid pCB55 and for both plasmids using the Topo I method.

### Protein-induced $\Delta Lk$ or $\beta$ does not correlate with protein-induced DNA bending by sequence-specific DNA-binding proteins

So far, all sequence-specific DNA-binding proteins used here sharply bend the DNA-binding sequences (Table 4). Since DNA bending is a geometric property which may alter DNA supercoiling (35), we decided to examine whether there is a correlation between the protein-induced DNA bending and  $\Delta Lk$  or  $\beta$ . Our results in Table 4 showed that the protein-induced  $\Delta Lk$  or  $\beta$  does not correlate with the protein-induced DNA bending. For instance,  $\lambda$  O protein bends each of the iterons  $\sim 90^\circ$ , and the  $\Delta Lk$  and  $\beta$  were determined to be  $-0.22$  and  $80^\circ$ , respectively. In contrast, CRP bends its binding site  $\sim 90^\circ$  as well; it does not have an induced  $\Delta Lk$  or  $\beta$ . Although AraC bends its binding site  $\sim 90^\circ$  in the presence or absence of L-arabinose (36), AraC-induced  $\Delta Lk$  or  $\beta$  is exclusively dependent on AraC-mediated DNA looping (Figure 5); in the presence of L-arabinose, it did not have an induced  $\Delta Lk$  or  $\beta$  (Table 4 and Figure 5). These results suggest that there is no relationship between the protein-induced  $\Delta Lk$  or  $\beta$  and the DNA bending. Our results also showed that there is no correlation between the DNA-binding constant and the protein-induced  $\Delta Lk$  or  $\beta$  (Table 4).

## DISCUSSION

White *et al.* (14,37) previously pointed out that closed circular DNA can be used as a probe for protein-induced structural changes. The primary quantity that can be experimentally determined is  $\Delta Lk$ . We showed here that it is feasible to simultaneously derive  $\Delta Lk$  and the DNA-binding constant from a simple titration experiment (Figures 2 and 4). Two proteins,  $\lambda$  O protein and GalR were chosen for these determinations. Both DNA-binding proteins bind to their recognition sites with moderate binding affinity (2,38) and, more importantly, without cooperativity (39). In this case, Equation 1 can be used to calculate  $\Delta Lk$  or  $\beta$  and the DNA-binding constant from experimental results shown in Figures 2 and 4. The DNA-binding constants of  $\lambda$  O protein and GalR were determined to be  $\sim 2.6 \times 10^8 \text{ M}^{-1}$  (Table 2) and  $\sim 5.8 \times 10^8 \text{ M}^{-1}$  (Table 3), respectively. These results are consistent with our EMSA results (Table 4) and the published results (2,38). These results further suggest that these two proteins bind to their recognition site non-cooperatively. Interestingly, the protein-induced  $\Delta Lk$  by both  $\lambda$  O protein ( $\sim -0.22$ ) and GalR ( $\sim -0.18$ ) is independent of the distance and phase match between each neighboring binding site (Tables 2 and 3), suggesting that the protein-induced  $\Delta Lk$  is a unique property for these sequence-specific DNA-binding proteins. White *et al.* also showed that  $\Delta Lk$  can be described by two geometrical terms: the surface linking number (SLk) and the winding number ( $\phi$ (14,37)). In this case,  $\Delta Lk = \Delta SLk + \Delta \phi$ . For  $\lambda$  O protein binding to its recognition site, i.e.  $\lambda$  DNA replication origin, we believe that  $\Delta Lk$  mainly results from the wrapping of the DNA



**Figure 8.** Proposed models to explain the protein-induced  $\Delta Lk$  by sequence-specific DNA-binding proteins. Models (A to B) show that certain site-specific DNA-wrapping proteins, such as  $\lambda$  O protein, bind and wrap the DNA recognition sites to induce the  $\Delta Lk$  of the DNA template. Models (C to D) demonstrate that DNA-looping proteins, such as AraC and LacI, form a topological nucleoprotein complex and therefore induce the  $\Delta Lk$  of the DNA template. Blue circle and red cylinder, respectively, represent the DNA recognition sequence for a site-specific DNA-binding protein and the site-specific DNA protein.

around  $\lambda$  O protein (Figure 8A and B), which forms a unique nucleoprotein structure, the O-some (6). Our unpublished atomic force microscope results strongly support this interpretation. For GalR binding to its recognition site, i.e. the  $O_E$  operator, we also believe that DNA wrapping is the main cause for the induced  $\Delta Lk$ , although further studies are needed to support this interpretation.

As shown in Figures 5 and 6, the protein-mediated DNA looping by AraC or LacI can supercoil the DNA-binding sites. For AraC, the DNA looping is exclusively responsible for the induced DNA supercoiling since AraC in the presence of L-arabinose cannot supercoil the DNA-binding sites (Figure 5B). Intriguingly, LacI-induced DNA supercoiling is dependent on the initial topological status of the DNA template in the protein-induced DNA supercoiling assays (Figure 6 and Table 4). The LacI-induced  $\Delta Lk$  in the Topo I assay (the initial  $\sigma$  of the DNA template is  $\sim -0.06$ ) is much greater than the  $\Delta Lk$  in the ligation assay (the initial  $\sigma$  of the DNA template is near 0). These results suggest that LacI is able to keep superhelical energy within the LacI- $O_1$  nucleoprotein complexes in a thermodynamically stable status (Supplementary Figure S3). Models in Figure 8C and D present a possible case to explain how the DNA looping proteins bring the two groups of the DNA-

binding sites together to fold into a topological nucleoprotein complex. It is feasible to form a well-defined nucleoprotein complex, including the topological folding of DNA and the DNA regulatory proteins, which allow multiple protein–protein interactions *in vivo*. These nucleoprotein complexes may also be able to serve as DNA supercoiling barriers or dividers (23) to block the localized supercoiling diffusion to assist DNA transactions, such as replication, transcription and recombination.

In this article, we also demonstrated that CRP did not induce a  $\Delta Lk$  upon binding to its recognition site although CRP bent the site sharply [Figure 7B; (17)]. According to the crystal structure of the CRP–DNA complexes (40), CRP wraps around the DNA-binding site, which should result in a surface linking number change ( $\Delta SLk$ ). However, it is possible that this  $\Delta SLk$  is compensated by changes in the local DNA winding in the CRP binding sites (the  $\Delta SLk$ - $\Delta\phi$  or twist-writhe compensation). In this situation, the DNA helical axis lies in a plane and the overall  $\Delta Lk$  is equal to zero. This explanation resembles the interpretation of the so-called ‘linking number paradox’ for the topological changes in a minichromosome (15). It has been known for a long time that 146 bp of DNA wrap around the histone octamer core 1.8 turns in a left-handed superhelix. In this case,  $\Delta SLk = -1.8$ . Initially, it was mis-expected that  $\Delta Lk = \Delta SLk$ , which resulted in the ‘linking number paradox’ in literature (14,15). As mentioned above,  $\Delta Lk = \Delta SLk + \Delta\phi$ , where  $\Delta\phi = -0.8$  and significantly compensated the  $\Delta SLk$  to yield  $\Delta Lk = -1$  that is corresponding to  $\sim -1$  per nucleosome in the topoisomerase relaxation assays (41). Another example of the  $\Delta SLk$ - $\Delta\phi$  compensation is the binding of the TATA box binding protein (TBP) to its recognition site which resulted in very little negative supercoiling in the topoisomerase relaxation assay (42). The crystal structures of the TBP–DNA complexes showed that the negative  $\Delta\phi$  from DNA unwinding is canceled by the positive  $\Delta SLk$  gained from ‘wrapping’ TBP around DNA (42,43).

In summary, we have constructed a series of DNA plasmids that carry many tandem copies of a DNA-binding site for sequence-specific DNA-binding proteins, such as  $\lambda$  O protein, GalR, CRP, AraC and LacI. Using these plasmid DNA templates, we determined the protein-induced  $\Delta Lk$  and demonstrated that the protein-induced  $\Delta Lk$  is an intrinsic property of these DNA-binding proteins. We also demonstrated that the protein-mediated DNA looping by AraC and LacI can introduce a  $\Delta Lk$  to the DNA-binding sites. Moreover, our results showed that the protein-induced  $\Delta Lk$  does not correlate with the protein-induced DNA bending and the DNA-binding constant.

## SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

## ACKNOWLEDGEMENTS

We would like to thank many colleagues, mentioned in the text, who supplied us with the sequence-specific DNA-binding proteins. We also thank Drs Roger McMacken, Sankar Adhya, Watson Lees and Zhongwei Li for critically reading the article before submission and for helpful discussion.

## FUNDING

National Institutes of Health (1SC1HD063059-01A1 to F.L.). Funding for open access charge: National Institutes of Health (1SC1HD063059-01A1).

*Conflict of interest statement.* None declared.

## REFERENCES

- Kaguni, J.M. (2006) DnaA: controlling the initiation of bacterial DNA replication and more. *Annu. Rev. Microbiol.*, **60**, 351–375.
- Zahn, K. and Blattner, F.R. (1985) Binding and bending of the lambda replication origin by the phage O protein. *EMBO J.*, **4**, 3605–3616.
- Mott, M.L. and Berger, J.M. (2007) DNA replication initiation: mechanisms and regulation in bacteria. *Nat. Rev. Microbiol.*, **5**, 343–354.
- Leonard, A.C. and Grimwade, J.E. (2005) Building a bacterial orisome: emergence of new regulatory features for replication origin unwinding. *Mol. Microbiol.*, **55**, 978–985.
- Alfano, C. and McMacken, R. (1989) Ordered assembly of nucleoprotein structures at the bacteriophage lambda replication origin during the initiation of DNA replication. *J. Biol. Chem.*, **264**, 10699–10708.
- Dodson, M., Roberts, J., McMacken, R. and Echols, H. (1985) Specialized nucleoprotein structures at the origin of replication of bacteriophage lambda: complexes with lambda O protein and with lambda O, lambda P, and Escherichia coli DnaB proteins. *Proc. Natl Acad. Sci. USA*, **82**, 4678–4682.
- Cases, I. and de Lorenzo, V. (2005) Promoters in the environment: transcriptional regulation in its natural context. *Nat. Rev. Microbiol.*, **3**, 105–118.
- Schleif, R. (2000) Regulation of the L-arabinose operon of Escherichia coli. *Trends Genet.*, **16**, 559–565.
- Matthews, K.S. and Nichols, J.C. (1998) Lactose repressor protein: functional properties and structure. *Prog. Nucleic Acid Res. Mol. Biol.*, **58**, 127–164.
- Kercher, M.A., Lu, P. and Lewis, M. (1997) Lac repressor-operator complex. *Curr. Opin. Struct. Biol.*, **7**, 76–85.
- Geanakopoulos, M., Vasmatazis, G., Zhurkin, V.B. and Adhya, S. (2001) Gal repressosome contains an antiparallel DNA loop. *Nat. Struct. Biol.*, **8**, 432–436.
- Kolb, A., Busby, S., Buc, H., Garges, S. and Adhya, S. (1993) Transcriptional regulation by cAMP and its receptor protein. *Annu. Rev. Biochem.*, **62**, 749–795.
- Adhya, S., Geanakopoulos, M., Lewis, D.E., Roy, S. and Aki, T. (1998) Transcription regulation by repressosome and by RNA polymerase contact. *Cold Spring Harb. Symp. Quant. Biol.*, **63**, 1–9.
- White, J.H., Gallo, R.M. and Bauer, W.R. (1992) Closed circular DNA as a probe for protein-induced structural changes. *Trends Biochem. Sci.*, **17**, 7–12.
- Bates, A.D. and Maxwell, A. (2005) *DNA Topology*. Oxford University Press, Oxford, UK.
- Maher, L.J. III (1998) Mechanisms of DNA bending. *Curr. Opin. Chem. Biol.*, **2**, 688–694.
- Kim, J., Zwiebe, C., Wu, C. and Adhya, S. (1989) Bending of DNA by gene-regulatory proteins: construction and use of a DNA bending vector. *Gene*, **85**, 15–23.

18. van der Vliet, P.C. and Verrijzer, C.P. (1993) Bending of DNA by transcription factors. *Bioessays*, **15**, 25–32.
19. Kim, R. and Kim, S.H. (1983) Direct measurement of DNA unwinding angle in specific interaction between lac operator and repressor. *Cold Spring Harb. Symp. Quant. Biol.*, **47(Pt 1)**, 451–454.
20. Kramer, H., Amouyal, M., Nordheim, A. and Muller-Hill, B. (1988) DNA supercoiling changes the spacing requirement of two lac operators for DNA loop formation with lac repressor. *EMBO J.*, **7**, 547–556.
21. Sadler, J.R., Tecklenburg, M. and Betz, J.L. (1980) Plasmids containing many tandem copies of a synthetic lactose operator. *Gene*, **8**, 279–300.
22. Chen, J. and Matthews, K.S. (1992) Deletion of lactose repressor carboxyl-terminal domain affects tetramer formation. *J. Biol. Chem.*, **267**, 13843–13850.
23. Leng, F. and McMacken, R. (2002) Potent stimulation of transcription-coupled DNA supercoiling by sequence-specific DNA-binding proteins. *Proc. Natl Acad. Sci. USA*, **99**, 9139–9144.
24. Wu, H.M. and Crothers, D.M. (1984) The locus of sequence-directed and protein-induced DNA bending. *Nature*, **308**, 509–513.
25. Thompson, J.F. and Landy, A. (1988) Empirical estimation of protein-induced DNA bending angles: applications to lambda site-specific recombination complexes. *Nucleic Acids Res.*, **16**, 9687–9705.
26. Scherer, G. (1978) Nucleotide sequence of the O gene and of the origin of replication in bacteriophage lambda DNA. *Nucleic Acids Res.*, **5**, 3141–3156.
27. Dodson, M., Echols, H., Wickner, S., Alfano, C., Mensa-Wilmot, K., Gomes, B., LeBowitz, J., Roberts, J.D. and McMacken, R. (1986) Specialized nucleoprotein structures at the origin of replication of bacteriophage lambda: localized unwinding of duplex DNA by a six-protein reaction. *Proc. Natl Acad. Sci. USA*, **83**, 7638–7642.
28. Lewis, D.E. and Adhya, S. (2002) In vitro repression of the gal promoters by GalR and HU depends on the proper helical phasing of the two operators. *J. Biol. Chem.*, **277**, 2498–2504.
29. Schleif, R. (1992) DNA looping. *Annu. Rev. Biochem.*, **61**, 199–223.
30. Oehler, S., Eismann, E.R., Kramer, H. and Muller-Hill, B. (1990) The three operators of the lac operon cooperate in repression. *EMBO J.*, **9**, 973–979.
31. Lobell, R.B. and Schleif, R.F. (1990) DNA looping and unlooping by AraC protein. *Science*, **250**, 528–532.
32. Kramer, H., Niemoller, M., Amouyal, M., Revet, B., von Wilcken-Bergmann, B. and Muller-Hill, B. (1987) lac repressor forms loops with linear DNA carrying two suitably spaced lac operators. *EMBO J.*, **6**, 1481–1491.
33. Kolb, A. and Buc, H. (1982) Is DNA unwound by the cyclic AMP receptor protein? *Nucleic Acids Res.*, **10**, 473–485.
34. Douc-Rasy, S., Kolb, A. and Prunell, A. (1989) Protein-induced unwinding of DNA: measurement by gel electrophoresis of complexes with DNA minicircles. Application to restriction endonuclease EcoRI, catabolite gene activator protein and lac repressor. *Nucleic Acids Res.*, **17**, 5173–5189.
35. Cozzarelli, N.R., Boles, T.C. and White, J.H. (1990) Primer on the topology and geometry of DNA supercoiling. In Cozzarelli, N.R. and Wang, J.C. (eds), *DNA Topology and Its Biological Effects*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, pp. 139–184.
36. Saviola, B., Seabold, R.R. and Schleif, R.F. (1998) DNA bending by AraC: a negative mutant. *J. Bacteriol.*, **180**, 4227–4232.
37. White, J.H., Cozzarelli, N.R. and Bauer, W.R. (1988) Helical repeat and linking number of surface-wrapped DNA. *Science*, **241**, 323–327.
38. Senechal, D.F. and Brenowitz, M. (1991) Determination of binding constants for cooperative site-specific protein-DNA interactions using the gel mobility-shift assay. *J. Biol. Chem.*, **266**, 13661–13671.
39. Brenowitz, M., Jamison, E., Majumdar, A. and Adhya, S. (1990) Interaction of the Escherichia coli Gal repressor protein with its DNA operators in vitro. *Biochemistry*, **29**, 3374–3383.
40. Schultz, S.C., Shields, G.C. and Steitz, T.A. (1991) Crystal structure of a CAP-DNA complex: the DNA is bent by 90 degrees. *Science*, **253**, 1001–1007.
41. Simpson, R.T., Thoma, F. and Brubaker, J.M. (1985) Chromatin reconstituted from tandemly repeated cloned DNA fragments and core histones: a model system for study of higher order structure. *Cell*, **42**, 799–808.
42. Kim, Y., Geiger, J.H., Hahn, S. and Sigler, P.B. (1993) Crystal structure of a yeast TBP/TATA-box complex. *Nature*, **365**, 512–520.
43. Kim, J.L. and Burley, S.K. (1994) 1.9 Å resolution refined structure of TBP recognizing the minor groove of TATAAAAAG. *Nat. Struct. Biol.*, **1**, 638–653.
44. Carra, J.H. and Schleif, R.F. (1993) Variation of half-site organization and DNA looping by AraC protein. *EMBO J.*, **12**, 35–44.