

12-22-2015

Alcohol and Cannabinoids Differentially Affect HIV Infection and Function of Human Monocyte-Derived Dendritic Cells (MDDC)

Marisela Agudelo

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida; Institute of NeuroImmune Pharmacology, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida, marisela.agudelo@fiu.edu

Gloria Figueroa

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, gloria.figueroa@fiu.edu

Adriana Yndart-Arias

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida; Institute of NeuroImmune Pharmacology, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida; Center for Personalized Nanomedicine, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida, adriana.yndartarias@fiu.edu

Gianna Casteleiro

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, gianna.casteleiro@fiu.edu

Karla Muñoz

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, karla.munoz1@fiu.edu

Recommended Citation

Agudelo M, Figueroa G, Yndart A, Casteleiro G, Muñoz K, Samikkannu T, Atluri V and Nair M P (2015) Alcohol and Cannabinoids Differentially Affect HIV Infection and Function of Human Monocyte-Derived Dendritic Cells (MDDC). *Front.Microbiol.* 6:1452. doi:10.3389/fmicb.2015.01452

This work is brought to you for free and open access by the Herbert Wertheim College of Medicine at FIU Digital Commons. It has been accepted for inclusion in HWCOC Faculty Publications by an authorized administrator of FIU Digital Commons. For more information, please contact dcc@fiu.edu.

See next page for additional authors



This work is licensed under a [Creative Commons Attribution 4.0 License](#).
Follow this and additional works at: https://digitalcommons.fiu.edu/com_facpub

 Part of the [Medicine and Health Sciences Commons](#)

Authors

Marisela Agudelo, Gloria Figueroa, Adriana Yndart-Arias, Gianna Casteleiro, Karla Muñoz, Thangavel Samikkannu, Venkata Atluri, and Madhavan P. Nair



Alcohol and Cannabinoids Differentially Affect HIV Infection and Function of Human Monocyte-Derived Dendritic Cells (MDDC)

Marisela Agudelo*, Gloria Figueroa, Adriana Yndart, Gianna Casteleiro, Karla Muñoz, Thangavel Samikkannu, Venkata Atluri and Madhavan P. Nair

Department of Immunology, Herbert Wertheim College of Medicine, Florida International University, Miami, FL, USA

OPEN ACCESS

Edited by:

Francois Villinger,
Emory University School of Medicine,
USA

Reviewed by:

Yong-Hui Zheng,
Michigan State University, USA
Anil Kumar,
University of Missouri–Kansas City,
USA

*Correspondence:

Marisela Agudelo
magudelo@fiu.edu

Specialty section:

This article was submitted to
Virology,
a section of the journal
Frontiers in Microbiology

Received: 16 September 2015

Accepted: 04 December 2015

Published: 22 December 2015

Citation:

Agudelo M, Figueroa G, Yndart A, Casteleiro G, Muñoz K, Samikkannu T, Atluri V and Nair MP (2015) Alcohol and Cannabinoids Differentially Affect HIV Infection and Function of Human Monocyte-Derived Dendritic Cells (MDDC). *Front. Microbiol.* 6:1452. doi: 10.3389/fmicb.2015.01452

During human immunodeficiency virus (HIV) infection, alcohol has been known to induce inflammation while cannabinoids have been shown to have an anti-inflammatory role. For instance cannabinoids have been shown to reduce susceptibility to HIV-1 infection and attenuate HIV replication in macrophages. Recently, we demonstrated that alcohol induces cannabinoid receptors and regulates cytokine production by monocyte-derived dendritic cells (MDDC). However, the ability of alcohol and cannabinoids to alter MDDC function during HIV infection has not been clearly elucidated yet. In order to study the potential impact of alcohol and cannabinoids on differentiated MDDC infected with HIV, monocytes were cultured for 7 days with GM-CSF and IL-4, differentiated MDDC were infected with HIV-1Ba-L and treated with EtOH (0.1 and 0.2%), THC (5 and 10 μ M), or JWH-015 (5 and 10 μ M) for 4–7 days. HIV infection of MDDC was confirmed by p24 and Long Terminal Repeats (LTR) estimation. MDDC endocytosis assay and cytokine array profiles were measured to investigate the effects of HIV and substances of abuse on MDDC function. Our results show the HIV + EtOH treated MDDC had the highest levels of p24 production and expression when compared with the HIV positive controls and the cannabinoid treated cells. Although both cannabinoids, THC and JWH-015 had lower levels of p24 production and expression, the HIV + JWH-015 treated MDDC had the lowest levels of p24 when compared to the HIV + THC treated cells. In addition, MDDC endocytic function and cytokine production were also differentially altered after alcohol and cannabinoid treatments. Our results show a differential effect of alcohol and cannabinoids, which may provide insights into the divergent inflammatory role of alcohol and cannabinoids to modulate MDDC function in the context of HIV infection.

Keywords: alcohol, cannabinoids, THC, JWH-015, MDDC, HIV infection

INTRODUCTION

Since the discovery of the Human Immunodeficiency Virus (HIV), the acquired immunodeficiency syndrome (AIDS) epidemic has been consistently associated with substance abuse and in some cases substance abuse treatment has been proposed as AIDS prevention (Metzger et al., 1998; Volkow, 2012). Recently, the common problems associated with HIV and substance abuse

comorbidities have been extensively discussed as demonstrated by the review of studies in humans and animal models (Chang et al., 2014; Molina et al., 2015). In the context of alcohol abuse and HIV comorbidity, a higher prevalence (42%) of alcohol problems in HIV-infected patients compared to the prevalence in non-HIV patients has been reported using the CAGE questionnaire (Samet et al., 2004). In the context of marijuana abuse and HIV comorbidity, people living with HIV have been reported to have a higher prevalence of cannabis use compared to that of the general population (Braitstein et al., 2001; Furler et al., 2004). For instance, approximately 25% of HIV/AIDS patients have been reported to use marijuana to ameliorate nausea, reduce pain, or as an appetite stimulant (Prentiss et al., 2004); however, the effects of cannabinoids on immune function of these immunocompromised patients remain poorly understood.

Overall, alcohol and drugs of abuse may modulate host-pathogen interactions including immune consequences of HIV infection. To date, it is well known that both the innate and adaptive immune systems are a complex network of cells and cytokines with the primary function of preventing infection. Although the immune system seems to be very resilient, there is evidence of the negative effects of environmental factors such as substances of abuse on immunity (Friedman et al., 2006). Previous studies using animal models have demonstrated a correlation between substance-induced immune dysfunction and greater susceptibility to infections such as in the case of chronic alcohol consumption and susceptibility to SIV infection (Kumar et al., 2005; Molina et al., 2006) or marijuana effects on immune function and ability to decrease host resistance to infections (Klein et al., 2003; Cabral and Staab, 2005; Cabral, 2006). Although substances of abuse have been shown to alter immune functions *in vitro* and in animal models, there is a lack of studies in humans that correlate immunosuppressive effects with increased incidence of infections, including infection with HIV as previously reviewed (Cabral, 2006; Molina et al., 2010a). Therefore, studies on the effects of substances of abuse such as alcohol and marijuana and their immunomodulatory mechanisms of action on HIV infection and disease progression are increasingly on demand since it is evident that the use of recreational substances such as alcohol and marijuana is common in this population and the consequences of substance abuse on HIV infection remain unclear. As previously reviewed most of the literature has highlighted the independent role of alcohol or marijuana on HIV (Chang et al., 2014; Nair and Agudelo, 2014; Molina et al., 2015; Nair et al., 2015); however, a side by side comparison of the *in vitro* effects of alcohol and the role of cannabinoid compounds such as tetrahydrocannabinol (THC) and the synthetic cannabinoid, JWH-015, on HIV infection of monocyte-derived dendritic cells (MDDC) has not been explored.

In the current study, we report, for the first time, the differential effects of substances such as alcohol, THC, and JWH-015, on dendritic cell function after differentiation and following HIV infection *in vitro*. It is hypothesized that alcohol and HIV

can exert their effects on MDDC by altering MDDC functions and these effects can be inversely regulated by THC and by the CB₂ agonist, JWH-015.

MATERIALS AND METHODS

Differentiation of Monocyte-Derived Dendritic Cells (MDDC)

For all the *in vitro* studies, leukopacks were commercially obtained from the community blood bank (One Blood, Miami, FL, USA) from at least three different blood donors. MDDC were prepared from peripheral blood mononuclear cells (PBMC) as previously described by us (Nair et al., 2005, 2009; Agudelo et al., 2013). Briefly, after the separation of adherent and non-adherent cells, monocyte were further purified using the EasySep™ Human Monocyte Enrichment Kit (Stemcell Technologies, catalog #19059), and allowed to differentiate into MDDC by culturing them with complete RPMI media containing 100 U/ml of GM-CSF and 100 U/ml IL-4 (R&D systems, Minneapolis, MN, USA).

Infection and Treatment of MDDC

After allowing the monocytes to differentiate for 7 days into MDDC using cytokines IL-4 and GM-CSF, the MDDCs were infected and treated. Cells were incubated with polybrene (2 μl/ml) for 30 min prior to infection with HIV-1Ba-L (National Institutes of Health AIDS Research and Reference Reagent Program; catalog no. 510) for 2 h at a concentration of 20 ng/10 million cells. Cells were then washed twice to remove unabsorbed virus and treated with 0.1% (~20 mM) or 0.2% (~40 mM) EtOH (Sigma-Aldrich, St. Louis, MO, USA; catalog #E7023), which are equivalent to the physiological blood alcohol concentrations (BAC) of 50 mg/dL, 100 mg/dL, and 200 mg/dL, respectively, and are close to the legal limit for driving under intoxication of 0.08% (80 mg/dL). The cannabinoid group was treated with 5–10 μM of Δ⁹-THC (Sigma-Aldrich, catalog #T4764) or 5–10 μM of JWH-015 (Tocris Bioscience, Ellisville, MO, USA, catalog #1341), which are concentrations within the range of previous reports using monocytes and macrophages exposed to cannabinoids (Williams et al., 2014; Roth et al., 2015). After HIV infection and treatments, MDDC were kept in culture for up to 7 days. Half of media was replenished every 48 h. Experimental design is provided in **Figure 1**.

p24 Secretion by ELISA

Concentrations of HIV-1 p24 antigen in HIV-infected MDDC culture supernatants were determined according to the manufacturer's instructions using commercially available retrotek HIV-1 p24 antigen ELISA kit (Cat. #0801111; Zeptometrix, Buffalo, NY, USA). ELISA was performed at days 4 and 7 post-infection and post treatment of MDDC to measure secreted p24 levels.

Intracellular p24 Levels by Flow Cytometry

For intracellular p24 staining, the cells were harvested and counted; equal amounts of cells (1×10^6) were aliquoted in 12 mm \times 75 mm polystyrene falcon tubes, catalog #352058 (BD Biosciences, San Jose, CA, USA), blocked with human serum (Chemicon International, Temecula, CA, USA), fixed, and permeabilized with Cytofix/Cytoperm solution (BD Bioscience). The p24 protein was detected with the anti-p24 FITC-labeled antibody (KC57-FITC; Beckman Coulter, Indianapolis, IN, USA). Cells were acquired on an Accuri C6 instrument (BD Accuri, Ann Arbor, MI, USA) and analyzed with FlowJo software (Tree Star, INC, Ashland, OR, USA). A total of 10,000 events were collected for each sample. Cells were gated based on isotype control, anti-IgG1-FITC (Fisher Scientific, Pittsburgh, PA, USA, cat. #1675864). Cells positive for p24 are shown as histogram overlay with shifted mean fluorescence intensity (MFI) compared to controls. Percentages of p24 positive cells and MFI were calculated with FlowJo software.

HIV Long Terminal Repeats (LTR) Expression by Quantitative Real Time PCR

Expression levels of HIV LTR were analyzed using real-time, quantitative PCR. Reverse transcription was performed with the 2720 Thermal cycler (Applied Biosystems) using aliquots of total RNA extracted from samples. The cDNA samples were diluted to 20 ng/ μ l and gene-specific primers were used: forward and reverse LTR HIV-1 primers and Taqman LTR-HIV (Cat. #VC00021 and VC00023, Sigma-Aldrich). To normalize the fold changes in gene expression, two internal control genes were used, GAPDH Taqman primer (Cat. #4331182, Life technologies) and 18S rRNA (Cat. #4333760F, Applied Biosystems). Fold changes were calculated based on uninfected control. All real-time PCR reactions were performed using the MX3005P detection system (Agilent Technologies) and the amplifications were done using the Brilliant II QPCR Master Mix (Cat. #600804, Agilent Technologies). The thermal cycling conditions were composed of an initial denaturation step at 95°C for 10 min, 45 cycles at 95°C for 30 s and 55°C for 1 min. The experiments were carried out in duplicates for each data point. The relative quantification in gene expression was determined using the $2^{-\Delta\Delta C_t}$ method (Livak and Schmittgen, 2001).

Endocytosis Assay

After MDDC differentiation, cells were treated with alcohol, THC or JWH-015 for 24 h. Then, the cells were cultured with FITC-dextran (1 mg/ml) in PBS + 1% FBS for 1 h at 37°C. The assay was terminated by adding ice-cold PBS. Cell pellets were washed three times with PBS and the MDDC immediately analyzed for accumulation of intracellular fluorescence in a Biotek plate reader (Winooski, VT, USA) at 490 nm excitation and 520 nm emission. In addition, MDDC infected with HIV and treated with alcohol, THC, or JWH-015, were also analyzed to assess their endocytic capacity.

Cytokine Arrays

Supernatants were collected at 7 days post-infection from MDDC infected with HIV and treated with either EtOH, THC, or JWH015. The expression of 48 inflammatory cytokines was analyzed as per manufacture recommendations with the RayBiotech inflammation arrays (catalog #AAH-INF-3-8, RayBiotech, Norcross, GA, USA). Data are expressed as fold change of cytokines secreted by HIV-infected and treated MDDC compared to the cytokines secreted by HIV-infected MDDC. Chemiluminescence signals were detected by a film developer and analyzed by densitometry using Image J software. Data were further analyzed using the RAYBIO Analysis Tool.

Statistical Analysis

Results are presented as mean \pm SEM, and *p*-values <0.05 are considered significant. Data were analyzed using Prism software (GraphPad Software, La Jolla, CA, USA), and statistical significance was determined by performing unpaired two-tailed Student's *t*-test or ANOVA followed by the Dunnett's post-test. All *in vitro* studies were performed with cells from at least three different donors with multiple replicates (at least two).

RESULTS

Alcohol and Cannabinoids Differentially Modulate p24 Production by MDDC

To determine if alcohol, THC, and JWH-015 modulate HIV-1 replication, MDDC were infected by the macrophage tropic replication competent strain, HIV-1Ba-L, followed by treatment with alcohol, THC, or JWH-015 using the experimental design (described in Materials and Methods, **Figure 1**). Although our ELISA results have demonstrated an increase in p24 levels over time after HIV infection (0–7 days), only the alcohol treated cells showed a significant dose dependent increase in p24 levels at 7 days post-infection (**Figure 2A**) while both concentrations





FIGURE 1 | Experimental design. Monocytes were allowed to differentiate into MDDC. MDDC were infected with HIV-1 for 2 h followed by treatment with alcohol (0.1 and 0.2%), THC (5 and 10 μ M) or JWH-015 (5 and 10 μ M). After HIV infection and treatments, MDDC were kept in culture for up to 7 days. Half of media was replenished every 48 h.



(5 and 10 μ M) of THC treatment had no major effect on p24 levels when compared to the HIV infected control (Figure 2B). Surprisingly, treatment with 10 μ M of the synthetic cannabinoid, JWH-015, caused a decrease in p24 levels by 7 days (Figure 2C).

Alcohol and Cannabinoids Differentially Modulate Intracellular Levels of p24 and the Percentage of HIV Positive MDDC

Flow cytometry analysis of p24 protein levels revealed an increase in both mean fluorescent intensity (MFI; Figures 3A,B) and in the percentage of cells expressing intracellular levels of the viral p24 core antigen (Figure 4) after 7 days post-infection confirming the internalization of virus and production of p24 core antigen by MDDC. In addition, both concentrations of alcohol, 0.1 and 0.2%, (Figures 3C,D) induced more than twofold increase in MFI when compared to HIV control, while the cannabinoids caused a reduction in MFI (Figures 3E-H) when compared to HIV control. Although there were visible differences in the modulation of MFI (Figures 3B,D,F,H) and in the percentage of cells expressing intracellular p24 levels (Figure 4) by alcohol and cannabinoids (THC and JWH-015), the effects observed were not statistically significant.

HIV-Infection Rate of MDDC is Increased in the Presence of Alcohol and THC

Assessment of HIV LTR by real-time PCR, confirmed HIV integration and infectivity of MDDC. Overall, we observed a dose dependent increase in HIV LTR in alcohol-treated HIV-infected MDDC compared to the untreated HIV-infected cells; however, only the higher concentration of alcohol (0.2%) induced a significant expression of LTR (Figure 5A). Although THC also caused an induction in LTR when compared to control and alcohol treated cells, it was not statistically significant (Figure 5B). The synthetic cannabinoid, JWH-015 did not affect LTR expression when compared to HIV control, in contrast, there


was a decrease in LTR after treatment with 10 μ M of JWH-015 (Figure 5C).

Alcohol Treatment Induces Higher Levels of Endocytosis by Uninfected MDDC When Compare to Control and Cannabinoid Treated Cells

Receptor mediated endocytosis was measured prior to HIV infection and after HIV infection by the uptake of FITC-dextran. Our results show a significant increase of FITC-dextran intake by MDDC that had been treated with alcohol (0.1%), THC (5 μ M), or JWH-015 (5 μ M; Figure 6). Although all treatments increased endocytosis in uninfected cells, THC and JWH-015 treated cells showed significantly lower levels of endocytosis when compared to alcohol treated MDDC. The overall increase in endocytosis by the substance abuse treated cells was only observed prior to HIV infection since after the cells were infected with HIV, there was an overall decrease in endocytosis with no differences across the groups.

Alcohol and Cannabinoids Differentially Modulate MDDC Cytokine Profiles During HIV Infection

The production of cytokines was assessed after HIV infection and stimulation with alcohol, THC or JWH-015. A summary of the cytokines that were modulated (\geq threefold) during HIV infection and substance abuse treatment are shown in Figure 7 (A) 0.1% alcohol, (B) 5 μ M THC, and (C) 5 μ M JWH-015. Treatment of HIV-infected MDDC with 0.1% alcohol caused an induction of seven cytokines (G-CSF, IL-3, IL-7, M-CSF, TGF- β 1, s TNF RI, and s TNF RII), while treatment with 5 μ M THC induced the levels of five cytokines (IL-3, IL-7, M-CSF, TGF- β 1, s TNF RI), and treatment with 5 μ M JWH-015 upregulated only three cytokines (IL-11, M-CSF, and PDGF-BB). Cytokine downregulation was also observed across the groups,



0.1% alcohol treatment downregulated seven cytokines (Eotaxin, IL-1 β , IL-11, IL-16, IL-17, MIG, and PDGF-BB), while 5 μ M THC treatment downregulated six cytokines (Eotaxin, IL-1 β , IL-11, IL-12 p40, MIG, PDGF-BB), and treatment with 5 μ M JWH-015 caused downregulation of two cytokines (Eotaxin and TGF- β 1). Specific fold differences between treatments and HIV control are highlighted in **Table 1**.

DISCUSSION

Our study supports evidence of a major role of MDDC in the enhancement of HIV infection and integration *in vitro*. Although the MDDC role during HIV infection has been somehow controversial due to the discrepancy regarding HIV infection of MDDC and evidence showing that HIV spreads from virus-containing MDDC (not necessarily productively infected) to T cells via an infectious synapse (McDonald et al., 2003; Blanchet et al., 2011), MDDC are professional antigen

presenting cells and they do get productively infected with HIV as shown by previous reports (Scott-Algara et al., 2008) and our own current findings demonstrating an increase in p24 intracellular levels, p24 secretion, and LTR. Despite MDDC questionable ability to get productively infected with HIV, they play a major role during HIV-1 *trans* infection of CD4⁺ T cells as previously reviewed (Agudelo et al., 2010; Rinaldo, 2013). Moreover, there is evidence of the ability of circulating DC to purge HIV-1 from latency and re-initiate virus replication in proliferating primary T lymphocytes (van der Sluis et al., 2013).

To the best of our knowledge, this is the first report that performs a side by side comparison of the *in vitro* effects of alcohol and cannabinoids in MDDC infected with HIV. We analyzed the effects of alcohol, THC, and JWH-015 on HIV infection and integration by measuring p24 production, intracellular p24 levels, and LTR. In addition, the immunomodulatory effects of these substances on MDDC function under HIV infection were also evaluated as measured by endocytosis




FIGURE 4 | Alcohol and cannabinoids differentially modulate the percentage of HIV positive MDDC. MDDC were infected with HIV and treated with alcohol, THC or JWH-015 for up to 7 days prior to assessing intracellular p24 levels by flow cytometry. Bar graphs correspond to the percentage of MDDC positive for p24 antigen. A total of 10,000 events were collected for each sample. Cells were gated based on isotype control. Data are expressed as % of p24 positive cells of three individual experiments.

human PBMC, which may be due to alcohol-induced functional impairment of various subsets of lymphocytes in the peripheral blood (Bagasra et al., 1993). In addition, substantial evidence from animal (Kumar et al., 2005; Molina et al., 2006; Sarkar and Chang, 2013) and human studies (Míguez-Burbano et al., 2009; Baum et al., 2010; Molina et al., 2014) has demonstrated that alcohol has an immune-modulatory effect during HIV disease progression and has the ability to disrupt HAART (Míguez-Burbano et al., 2009). More specific associations between alcohol consumption and lower levels of CD4 T cells have been reported among HIV-infected alcoholic patients (Samet et al., 2007; Baum et al., 2010). However, additional studies are necessary to elucidate the mechanisms of alcohol-induced immune-modulation in the context of HIV infection.

In contrast to alcohol results, when the HIV-infected MDDC were treated with the main psychoactive component of marijuana, THC, there were no changes in p24 secretion (Figure 2B). However, there was an evident decrease in p24 intracellular levels (Figures 3E,F) and in the percentage of cells expressing p24 (Figure 4) implying a reduction in HIV infection. Surprisingly, when LTR levels were examined, integration seemed to be elevated since THC acted similar to alcohol by inducing LTR (Figure 5B) levels; although, the induction had high variability between samples and was not significant. In contrast, a previous report using THC immediately prior to or continuously following HIV-1 exposure failed to alter infection (Williams et al., 2014). However, their study was performed with primary human monocytes and treatment was performed during differentiation of monocytes. Although in different cells and under different treatment conditions, some of their findings (Williams et al., 2014) are somehow consistent to our results showing that THC reduced the number of p24 positive cells (Figure 4) with little to no effect on virus production (Figure 2B). It is relevant to point

and cytokine production. Our results show *in vitro* alcohol treatment of MDDC increased p24 secretion (Figure 2A), intracellular p24 levels (Figures 3C,D and 4), percentage of cells expressing p24, and LTR (Figure 5A). Our findings are in accordance with previous reports showing alcohol enhances HIV-1 infection of human monocyte-derived cells; however, those reports were performed in cord blood monocyte-derived macrophages (Mastrogiannis et al., 2014) and in blood monocyte-derived macrophages (Wang et al., 2002). Furthermore, alcohol ingestion has been shown to increase HIV-1 replication in






FIGURE 5 | HIV-infection rate of MDDC is increased in the presence of alcohol and THC. MDDC were infected with HIV and treated with (A) alcohol, (B) THC, or (C) JWH-015 for up to 7 days prior to assessment of HIV LTR by real-time PCR. Data are expressed as Transcript Accumulation index (TAI). To normalize the fold changes in gene expression, two internal control genes were used, GAPDH and 18S rRNA. Fold changes were calculated based on uninfected control. Data are representative of three individual experiments performed at least in duplicates.



out that in our *in vitro* findings, there were no differences in p24 secretion when we compared THC treated HIV-infected MDDC with non-treated HIV-infected MDDC. Similarly, animal studies have shown that simian immunodeficiency virus (SIV)-infected female rhesus macaques exposed to chronic administration of THC prior to and following SIV infection exhibited no changes in markers of SIV disease, including viral load (Amedee et al., 2014). However, there were contrasting effects of chronic THC exposure in males versus females, with males having lower plasma and CSF viral loads (Molina et al., 2010b).

In an effort to elucidate the mechanism of THC suppression of HIV-1 infection, we also performed treatment with CB₂ agonist, JWH-015. HIV-infected MDDC treated with JWH-015 showed a decrease in p24 secretion by 7 days post-infection (Figure 2C), a decrease in p24 intracellular levels (Figures 3G,H) and a decrease in the percentage of p24 positive cells (Figure 4) when compared with the HIV-infected control or the HIV-infected alcohol treated cells. We also observed a decrease in LTR levels after treatment with 10 μM JWH-015 when compared with HIV-infected control and HIV-infected alcohol or cannabinoid treated cells. These findings may support a major role of CB₂ receptors in the HIV replication machinery, and these results are in accordance with previous findings with macrophages revealing a marked decrease in HIV-1 LTR activation by the CB₂ ligands, JWH133, GP1a, O-1966 and showing a significant decrease in RT activity when a CB₂ agonist was present (Ramirez et al., 2013). Taken together, these results using MDDC and previous findings with macrophages (Ramirez et al., 2013) indicate that CB₂ may limit HIV-1 infection in human monocyte-derived cells. Furthermore, a similar study with JWH-133, ACEA, and O-1602 indicated that the ability of THC to reduce macrophage susceptibility to HIV infection during monocyte differentiation was mediated primarily through CB₂ (Williams et al., 2014).



Besides modulation of HIV infection of MDDC, our results demonstrate the ability of alcohol and cannabinoids (THC and JWH-015) to affect DC function as shown by an increase in endocytosis in uninfected cells and an overall decrease in the endocytic capacity after HIV infection (Figure 6). These findings may be explained by the capacity of immature dendritic cells to be highly endocytic (Garrett et al., 2000) and the ability of HIV-1_{Bal} to induce partial maturation of dendritic cells (Mercier et al., 2013). Therefore, based on our results and previous literature (Mercier et al., 2013) the observed decrease in endocytosis after HIV infection and the attenuation of the substance abuse-induction of endocytosis might be due to the maturation status of the MDDC. We are the first ones to show an induction of endocytosis by healthy uninfected

TABLE 1 | Summary of the effects of alcohol and cannabinoids, THC and JWH-015 on the modulation of cytokine production by HIV-infected MDDC.

| Cytokine | EtOH 0.1% | THC 5 μ M | JWH-015 5 μ M |
|----------------|--------------|---------------|-------------------|
| EOTAXIN | -52.3 | -55.2 | -15.5 |
| GCSF | 3.0 | 2.8 | 0.6 |
| ICAM-1 | 1.5 | 1.6 | 1.1 |
| I-309 | 0.6 | 1.8 | 2.0 |
| IL-1 β | -20.5 | -4.0 | 2.1 |
| IL-3 | 9.1 | 6.9 | 2.3 |
| IL-6sR | 2.5 | 2.6 | 1.6 |
| IL-7 | 4.4 | 5.6 | 0.1 |
| IL-11 | -8.3 | -16.2 | 16.7 |
| IL-12 p40 | -1.4 | -5.5 | 1.4 |
| IL-12 p70 | 0.3 | 0.0 | 2.4 |
| IL-16 | -4.1 | -1.4 | 1.1 |
| IL-17 | -3.1 | -1.5 | 1.5 |
| M-CSF | 24.1 | 37.1 | 3.1 |
| MIG | -3.6 | -4.2 | 1.7 |
| MIP-1 α | 2.7 | 2.9 | 2.0 |
| MIP-1 δ | 1.1 | 3.0 | 1.7 |
| TGF- β 1 | 8.7 | 3.3 | -5.6 |
| TNF- α | -0.9 | 0.5 | 2.1 |
| TNF- β | -2.6 | -0.4 | 2.7 |
| s TNF RI | 8.3 | 4.9 | -0.7 |
| s TNF RII | 4.1 | 2.5 | 0.1 |
| PDGF-BB | -64.9 | -46.4 | 10.4 |

Cytokine array was conducted using supernatants from HIV-infected MDDC treated with alcohol (0.1%), THC (5 μ M), of JWH-015 (5 μ M). A summary of cytokines with more than threefold increase (light gray) or decrease (dark gray) are highlighted in the table. Data are expressed as fold change of cytokines secreted by HIV-infected and treated MDDC compared to the cytokines secreted by HIV-infected MDDC.

MDDC treated with alcohol or cannabinoids. Previous literature in the context of alcohol has demonstrated the ability of alcohol to modulate endocytosis; however, the reports are in rat hepatocytes, in which ethanol impaired receptor-mediated endocytosis (Casey et al., 1989; Tuma et al., 1991). In the context of cannabinoids, there is evidence demonstrating that exposure to THC impairs the capacity for receptor-mediated endocytosis by human dendritic cells (Roth et al., 2015); however, their immunoregulatory effects on human MDDC were observed at lower THC concentrations (0.5–3.2 μ M). Other findings in mice, have demonstrated that under certain conditions, THC enhances HIV antigen-specific immune responses (Chen et al., 2015).

In the past, alcohol and cannabinoids have been shown to alter several aspects of immune function including cytokine production and lymphocyte phenotype (Friedman et al., 1995, 2003; Szabo, 1999; Klein et al., 2003; Szabo et al., 2004; Crews et al., 2006; Molina et al., 2010a). According to the cytokine analysis performed (Figure 7 and Table 1), alcohol and cannabinoids induced differential cytokine production by HIV-infected MDDC and caused a major immune-modulatory effect, which may exacerbate HIV infection and pathogenesis. When compared to HIV control, alcohol (0.1%) and THC (5 μ M) treatments had similar effects and induced IL-3, IL-7, M-CSF,

TGF- β 1, sTNF RI, and sTNF RII while decreasing IL-1 β , IL-11, IL-12p40, TNF β , and PDGF-BB. However, the JWH-015 treatments had opposing effects as shown by an increase on IL-1 β , IL-11, IL-12p40, and IL-12 p70, MCSF, MIP1 α , TNF α , TNF β , and PDGF-BB while reducing only TGF- β 1 and sTNF RI. All three substances, alcohol (-52.3), THC (-55.2) and JWH-015 (-15.5), remarkably reduced the levels of eotaxin, which is a chemokine associated with lower susceptibility to infection in macaques (Promadej-Lanier et al., 2010). Another unique difference observed was on the effects on IL-11 production after alcohol (-8.3), THC (-16.2), and JWH-015 (16.7) treatments. This is relevant since IL-11 may reduce HIV-1-mediated immune activation while achieving elimination of virally infected cells as previously reported (Favors et al., 2012). Lastly, the effects on PDGF-BB were also different between alcohol (-64.9), THC (-46.4) versus JWH-015 (10.4) treatments. This is relevant within the HIV field since platelet-derived growth factor (PDGF) is one of the toxic mediators shown to be upregulated in the brains of macaques with SIV encephalitis and shown to play a major role in HIV neuro-pathogenesis (Potula et al., 2004; Bethel-Brown et al., 2011).

Overall, the differential immune-modulatory effects of alcohol and cannabinoids could be explained by the different mechanisms of action of these substances since the effects of marijuana on the immune system have been demonstrated to be receptor-mediated, occurring directly via specific receptors on immune cells (Friedman et al., 2003). For instance, THC exerts its effects through both CB₁ and CB₂ receptors (Matsuda et al., 1990; Munro et al., 1993), while JWH-015 acts as a CB₂ agonist (Huffman, 2000). Unlike cannabinoids and other addictive drugs of abuse, alcohol, does not appear to bind to a specific receptor (Peter et al., 2008). In summary alcohol and cannabinoids have been shown to induce neuro-immunomodulatory consequences during healthy conditions and during HIV infection and disease progression as highlighted in several recent reviews (Chang et al., 2014; Molina et al., 2015; Nair et al., 2015). Our current findings were able to highlight a divergent role of both substances in HIV infection and integration of human MDDC, and along with future studies, will allow us to elucidate the immunological consequences of alcohol and/or cannabinoids on the innate and adaptive immune system of an immune-compromised host. However, additional studies are still necessary to elucidate the molecular mechanisms involving alcohol and/or cannabinoid induced effects in the context of HIV infection.

AUTHOR CONTRIBUTIONS

MA contributed to the experimental design of the project, perform experiments, perform data analysis, and wrote the manuscript, GF and GC performed experiments, AY and KM provided technical assistance and advice with experimental troubleshooting, TS provided advice and guidance regarding HIV experiments, VA provided assistance with HIV experiments, LTR estimation and data analysis, MN provided financial support and guidance with HIV experimental design.

ACKNOWLEDGMENTS

This research was supported in part by the National Institute on Alcohol Abuse and Alcoholism, award K99/R00 AA021264, and the National Institute on Drug Abuse, award R01DA040537. Additional support as part of startup package has been received from the Department of Immunology, Institute of

Neuroimmune Pharmacology, Herbert Wertheim College of Medicine, and FIU-Office of Research and Economic Development. MBRS RISE fellow, GC, was supported by NIH/NIGMS R25 GM061347. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

- Agudelo, M., Saiyed, Z., Samikkannu, T., Khatavkar, P., Gandhi, N., Nair, N., et al. (2010). *Dendritic Cells: Types, Life Cycles and Biological Functions. Role of Dendritic Cells in HIV Infection*, Chap. 10. New York, NY: Nova Publishers.
- Agudelo, M., Yndart, A., Morrison, M., Figueroa, G., Muñoz, K., Samikkannu, T., et al. (2013). Differential expression and functional role of cannabinoid genes in alcohol users. *Drug Alcohol Depend.* 133, 789–793. doi: 10.1016/j.drugalcdep.2013.08.023
- Amedee, A. M., Nichols, W. A., LeCapitaine, N. J., Stouwe, C. V., Birke, L. L., Lacour, N., et al. (2014). Chronic Δ 9-tetrahydrocannabinol administration may not attenuate simian immunodeficiency virus disease progression in female rhesus macaques. *Res. Hum. Retrovir.* 30, 1216–1225. doi: 10.1089/aid.2014.0108
- Bagasra, O., Balla, A. K., Lischner, H. W., and Pomerantz, R. J. (1993). Alcohol intake increases human immunodeficiency virus type 1 replication in human peripheral blood mononuclear cells. *J. Infect. Dis.* 167, 789–797. doi: 10.1093/infdis/167.4.789
- Baum, M. K., Rafie, C., Lai, S., Sales, S., Page, J. B., and Campa, A. (2010). Alcohol use accelerates HIV disease progression. *Res. Hum. Retrovir.* 26, 511–518. doi: 10.1089/aid.2009.0211
- Bethel-Brown, C., Yao, H., Callen, S., Lee, Y. H., Dash, P. K., Kumar, A., et al. (2011). HIV-1 tat-mediated induction of platelet-derived growth factor in astrocytes: role of early growth response gene 1. *J. Immunol.* 186, 4119–4129. doi: 10.4049/jimmunol.1002235
- Blanchet, F., Moris, A., Mitchell, J. P., and Piguat, V. (2011). A look at HIV journey: from dendritic cells to infection spread in CD4+ T cells. *Curr. Opin. HIV AIDS* 6, 391–397. doi: 10.1097/COH.0b013e328349b0a0
- Braitstein, P., Kendall, T., Chan, K., Wood, E., Montaner, J. S. G., O'Shaughnessy, M. V., et al. (2001). Mary-Jane and her patients: sociodemographic and clinical characteristics of HIV-positive individuals using medical marijuana and antiretroviral agents. *AIDS* 15, 532–533. doi: 10.1097/00002030-200103090-00016
- Cabral, G. (2006). Drugs of abuse, immune modulation, and AIDS. *J. Neuroimmune Pharmacol.* 1, 280–295. doi: 10.1007/s11481-006-9023-5
- Cabral, G., and Staab, A. (2005). Effects on the immune system. *Handb. Exp. Pharmacol.* 168, 385–423. doi: 10.1007/3-540-26573-2_13
- Casey, C. A., Kragoskow, S. L., Sorrell, M. F., and Tuma, D. J. (1989). Ethanol-induced impairments in receptor-mediated endocytosis of asialoorosomucoid in isolated rat hepatocytes: time course of impairments and recovery after ethanol withdrawal. *Alcohol. Clin. Exp. Res.* 13, 258–263. doi: 10.1111/j.1530-0277.1989.tb00323.x
- Chang, S. L., Connaghan, K. P., Wei, Y., and Li, M. D. (2014). “Chapter thirteen - neurohiv and use of addictive substances,” in *International Review of Neurobiology*, Vol. 118, eds D. S. Changhai Cui and R. A. Harris (Waltham, MA: Academic Press), 403–40.
- Chen, W., Crawford, R., Kaplan, B. F., and Kaminski, N. (2015). Modulation of HIVGP120 antigen-specific immune responses in vivo by Δ 9-tetrahydrocannabinol. *J. Neuroimmune Pharmacol.* 10, 344–55. doi: 10.1007/s11481-015-9597-x
- Crews, F. T., Bechara, R., Brown, L. A., Guidot, D. M., Mandrekar, P., Oak, S., et al. (2006). Cytokines and alcohol. *Alcohol. Clin. Exp. Res.* 30, 720–730. doi: 10.1111/j.1530-0277.2006.00084.x
- Favors, S. E., Curd, L. M., and Gregg, R. K. (2012). Use of the anti-inflammatory cytokine interleukin-11 to reverse HIV-1gp120 repression of a natural killer cell line. *Cell Immunol.* 276, 1–5. doi: 10.1016/j.cellimm.2012.02.011
- Friedman, H., Klein, T. W., Newton, C., and Daaka, Y. (1995). Marijuana, receptors and immunomodulation. *Adv. Exp. Med. Biol.* 373, 103–113. doi: 10.1007/978-1-4615-1951-5_15
- Friedman, H., Newton, C., and Klein, T. W. (2003). Microbial infections, immunomodulation, and drugs of abuse. *Clin. Microbiol. Rev.* 16, 209–219. doi: 10.1128/CMR.16.2.209-219.2003
- Friedman, H., Pross, S., and Klein, T. W. (2006). Addictive drugs and their relationship with infectious diseases. *FEMS Immunol. Med. Microbiol.* 47, 330–342. doi: 10.1111/j.1574-695X.2006.00097.x
- Furler, M. D., Einarsen, T. R., Millson, M., Walmsley, S., and Bodayan, R. (2004). Medicinal and recreational marijuana use by patients infected with HIV. *AIDS Patient Care STDS* 8, 215–228. doi: 10.1089/108729104323038892
- Garrett, W. S., Chen, L.-M., Kroschewski, R., Ebersold, M., Turley, S., Trombetta, S., et al. (2000). Developmental control of endocytosis in dendritic cells by Cdc42. *Cell* 102, 325–334. doi: 10.1016/S0092-8674(00)00038-6
- Huffman, J. (2000). The search for selective ligands for the CB2 receptor. *Curr. Pharm. Des.* 6, 1323–1337. doi: 10.2174/1381612003399347
- Klein, T. W., Newton, C., Larsen, K., Lu, L., Perkins, I., Nong, L., et al. (2003). The cannabinoid system and immune modulation. *J. Leukoc. Biol.* 74, 486–496. doi: 10.1189/jlb.0303101
- Kumar, R., Perez-Casanova, A. E., Tirado, G., Noel, R. J., Torres, C., Rodriguez, I., et al. (2005). Increased viral replication in simian immunodeficiency virus/simian-HIV-infected macaques with self-administering model of chronic alcohol consumption. *J. Acquir. Immune Defic. Syndr.* 39, 386–390. doi: 10.1097/01.qai.0000164517.01293.84
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* 25, 402–408. doi: 10.1006/meth.2001.1262
- Mastrogiannis, D. S., Wang, X., Dai, M., Li, J., Wang, Y., Zhou, Y., et al. (2014). Alcohol enhances HIV infection of cord blood monocyte-derived macrophages. *Curr. HIV Res.* 12, 301–308. doi: 10.2174/1570162X12666140721124923
- Matsuda, L. A., Lolait, S. J., Brownstein, M. J., Young, A. C., and Bonner, T. I. (1990). Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 346, 561–564. doi: 10.1038/346561a0
- McDonald, D., Wu, L., Bohks, S. M., KewalRamani, V. N., Unutmaz, D., and Hope, T. J. (2003). Recruitment of HIV and its receptors to dendritic cell-T cell junctions. *Science* 300, 1295–1297. doi: 10.1126/science.1084238
- Mercier, S. K., Donaghy, H., Botting, R. A., Turville, S. G., Harman, A. N., Nasr, N., et al. (2013). The microvesicle component of HIV-1 inocula modulates dendritic cell infection and maturation and enhances adhesion to and activation of T lymphocytes. *PLoS Pathog* 9:e1003700. doi: 10.1371/journal.ppat.1003700
- Metzger, D. S., Navaline, H., and Woody, G. E. (1998). Drug abuse treatment as AIDS prevention. *Public Health Rep.* 113(Suppl. 1), 97–106.
- Míguez-Burbano, M. J., Lewis, J. E., Fishman, J., Asthana, D., and Malow, R. M. (2009). The influence of different types of alcoholic beverages on disrupting highly active antiretroviral treatment (HAART) outcome. *Alcohol. Alcohol.* 44, 366–371. doi: 10.1093/alcalc/agg024
- Molina, P., Amedee, A., Winsauer, P., Nelson, S., Bagby, G., and Simon, L. (2015). Behavioral, metabolic, and immune consequences of chronic alcohol or cannabinoids on HIV/AIDS: studies in the non-human primate SIV model. *J. Neuroimmune Pharmacol.* 10, 217–232. doi: 10.1007/s11481-015-9599-8
- Molina, P. E., Bagby, G. J., and Nelson, S. (2014). Biomedical consequences of alcohol use disorders in the HIV-infected host. *Curr. HIV Res.* 12, 265–275. doi: 10.2174/1570162X12666140721121849
- Molina, P. E., Happel, K. I., Zhang, P., Kolls, J. K., and Nelson, S. (2010a). Focus on: alcohol and the immune system. *Alcohol. Res. Health* 33, 97–108.

- Molina, P. E., Winsauer, P., Zhang, P., Walker, E., Birke, L., Amedee, A., et al. (2010b). Cannabinoid administration attenuates the progression of simian immunodeficiency virus. *AIDS Res. Hum. Retrovir.* 27, 585–592. doi: 10.1089/aid.2010.0218
- Molina, P. E., McNurlan, M., Rathmacher, J., Lang, C. H., Zambell, K. L., Purcell, J., et al. (2006). Chronic alcohol accentuates nutritional, metabolic, and immune alterations during asymptomatic simian immunodeficiency virus infection. *Alcohol. Clin. Exp. Res.* 30, 2065–2078. doi: 10.1111/j.1530-0277.2006.00252.x
- Munro, S., Thomas, K. L., and Abu-Shaar, M. (1993). Molecular characterization of a peripheral receptor for cannabinoids. *Nature* 365, 61–65. doi: 10.1038/365061a0
- Nair, M., Figueroa, G., Casteleiro, G., Muñoz, K., and Agudelo, M. (2015). Alcohol versus cannabinoids: a review of their opposite neuro-immunomodulatory effects and future therapeutic potentials. *J. Alcohol. Drug Depend.* 3:p1184. doi: 10.4172/2329-6488.1000184
- Nair, M., Reynolds, J., Mahajan, S., Schwartz, S., Aalinkeel, R., Bindukumar, B., et al. (2005). RNAi-directed inhibition of DC-SIGN by dendritic cells: prospects for HIV-1 therapy. *AAPS J.* 7, E572–E578. doi: 10.1208/aapsj070358
- Nair, M., Saiyed, Z., Nair, N., Gandhi, N., Rodriguez, J., Boukli, N., et al. (2009). Methamphetamine enhances hiv-1 infectivity in monocyte derived dendritic cells. *J. Neuroimmune Pharmacol.* 4, 129–139. doi: 10.1007/s11481-008-9128-0
- Nair, M. P., and Agudelo, M. (2014). Commentary on the interactions of HIV and alcohol in the central nervous system. *Alcohol. Clin. Exp. Res.* 38, 624–625. doi: 10.1111/acer.12326
- Peter, C., Sanjiv, V. B., and Paula, L. H. (2008). How adaptation of the brain to alcohol leads to dependence: a pharmacological perspective. *Alcohol. Res. Health* 31, 310–339.
- Potula, R., Dhillon, N., Sui, Y., Zien, C. A., Funa, K., Pinson, D., et al. (2004). Association of platelet-derived growth factor- β chain with simian human immunodeficiency virus encephalitis. *Am. J. Pathol.* 165, 815–824. doi: 10.1016/S0002-9440(10)63344-5
- Prentiss, D., Power, R., Balmas, G., Tzuang, G., and Israelski, D. M. (2004). Patterns of marijuana use among patients with HIV/AIDS followed in a public health care setting. *J. Acquir. Immune Defic. Syndr.* 35, 38–45. doi: 10.1097/00126334-200401010-00005
- Promadej-Lanier, N., Hanson, D. L., Srinivasan, P., Luo, W., Adams, D. R., Guenther, P. C., et al. (2010). Resistance to simian HIV infection is associated with high plasma interleukin-8, RANTES and Eotaxin in a macaque model of repeated virus challenges. *J. Acquir. Immune Defic. Syndr.* 53, 574–581. doi: 10.1097/qai.0b013e3181d3521f
- Ramirez, S. H., Reichenbach, N. L., Fan, S., Rom, S., Merkel, S. F., Wang, X., et al. (2013). Attenuation of HIV-1 replication in macrophages by cannabinoid receptor 2 agonists. *J. Leukoc. Biol.* 93, 801–810. doi: 10.1189/jlb.1012523
- Rinaldo, C. R. (2013). HIV-1 trans infection of CD4+ T cells by professional antigen presenting cells. *Scientifica* 2013:164203. doi: 10.1155/2013/164203
- Roth, M., Castaneda, J., and Kiertscher, S. (2015). Exposure to Δ^9 -tetrahydrocannabinol impairs the differentiation of human monocyte-derived dendritic cells and their capacity for T cell activation. *J. Neuroimmune Pharmacol.* 10, 333–343. doi: 10.1007/s11481-015-9587-z
- Samet, J. H., Cheng, D. M., Libman, H., Nunes, D. P., Alperen, J. K., and Saitz, R. (2007). Alcohol consumption and HIV disease progression. *J. Acquir. Immune Defic. Syndr.* 46, 194–199. doi: 10.1097/QAI.0b013e318142aabb
- Samet, J. H., Phillips, S. J., Horton, N. J., Traphagen, E. T., and Freedberg, K. A. (2004). Detecting alcohol problems in HIV-infected patients: use of the CAGE questionnaire. *AIDS Res. Hum. Retrovi.* 20, 151–155. doi: 10.1089/088922204773004860
- Sarkar, S., and Chang, S. L. (2013). Ethanol concentration-dependent alterations in gene expression during acute binge drinking in the HIV-1 transgenic rat. *Alcohol. Clin. Exp. Res.* 37, 1082–1090. doi: 10.1111/acer.12077
- Scott-Algara, D., Arnold, V., Didier, C., Kattan, T., Pirozzi, G., Barré-Sinoussi, F., et al. (2008). The CD85j+ NK cell subset potently controls HIV-1 replication in autologous dendritic cells. *PLoS ONE* 3:e1975. doi: 10.1371/journal.pone.0001975
- Szabo, G. (1999). Consequences of alcohol consumption on host defence. *Alcohol* 34, 830–841. doi: 10.1093/alcalc/34.6.830
- Szabo, G., Catalano, D., White, B., and Mandrekar, P. (2004). Acute alcohol consumption inhibits accessory cell function of monocytes and dendritic cells. *Alcohol. Clin. Exp. Res.* 28, 824–828. doi: 10.1097/01.ALC.0000127104.80398.9B
- Tuma, D. J., Casey, C. A., and Sorrell, M. F. (1991). Chronic ethanol-induced impairments in receptor-mediated endocytosis of insulin in rat hepatocytes. *Alcohol. Clin. Exp. Res.* 15, 808–813. doi: 10.1111/j.1530-0277.1991.tb00606.x
- van der Sluis, R. M., Montfort, T. V., Pollakis, G., Sanders, R. W., Speijer, D., Berkhout, B., et al. (2013). Dendritic cell-induced activation of latent HIV-1 provirus in actively proliferating primary T lymphocytes. *PLoS Pathog* 9:e1003259. doi: 10.1371/journal.ppat.1003259
- Volkow, N. D. (2012). *Research Report Series: Drug Abuse and HIV*. North Bethesda, MD: National Institute on Drug Abuse.
- Wang, X., Douglas, S. D., Metzger, D. S., Guo, C.-J., Li, Y., O'Brien, C. P., et al. (2002). Alcohol potentiates HIV-1 infection of human blood mononuclear phagocytes. *Alcohol. Clin. Exp. Res.* 26, 1880–1886. doi: 10.1111/j.1530-0277.2002.tb02496.x
- Williams, J., Appelberg, S., Goldberger, B., Klein, T., Sleasman, J., and Goodenow, M. (2014). Δ^9 -tetrahydrocannabinol treatment during human monocyte differentiation reduces macrophage susceptibility to HIV-1 infection. *J. Neuroimmune Pharmacol.* 9, 369–379. doi: 10.1007/s11481-014-9527-3

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Agudelo, Figueroa, Yndart, Casteleiro, Muñoz, Samikkannu, Atluri and Nair. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.