

1998

# Florida Keys National Marine Sanctuary Water Quality Monitoring Project: 1998 Annual Report

Ronald Jones

*Southeast Environmental Research Center*

Joseph N. Boyer

*Southeast Environmental Research Center, Florida International University, boyerj@fiu.edu*

Follow this and additional works at: <https://digitalcommons.fiu.edu/sercrp>



Part of the [Environmental Monitoring Commons](#), and the [Water Resource Management Commons](#)

---

## Recommended Citation

Jones, Ronald and Boyer, Joseph N., "Florida Keys National Marine Sanctuary Water Quality Monitoring Project: 1998 Annual Report" (1998). *SERC Research Reports*. 50.  
<https://digitalcommons.fiu.edu/sercrp/50>

This work is brought to you for free and open access by the Southeast Environmental Research Center at FIU Digital Commons. It has been accepted for inclusion in SERC Research Reports by an authorized administrator of FIU Digital Commons. For more information, please contact [dcc@fiu.edu](mailto:dcc@fiu.edu).

**Florida Keys National Marine Sanctuary**  
**Water Quality Monitoring Project**

1998 Annual Report

Principal Investigators

Ronald D. Jones

and

Joseph N. Boyer

Southeast Environmental Research Program  
Florida International University  
Miami, FL 33199

US EPA Contract #X994621-94-0

This is contribution #81 of the Southeast Environmental Research Program at Florida International University.

# **Table of Contents**

I. EXECUTIVE SUMMARY

II. PROJECT BACKGROUND

III. RESULTS AND DISCUSSION

IV. LIST OF APPENDICES

## I. Executive Summary

This report serves as a summary of our efforts to date in the execution of the water quality monitoring program for the FKNMS. Last year we added 4 sampling sites and made minor adjustments to six others to establish water quality sampling within some of the Sanctuary Preservation Areas and Ecological Reserves. We have received 11 requests for data by outside researchers working in the FKNMS of which one has resulted in a master's thesis. Two scientific manuscripts have been submitted for publication: one is a book chapter in *Linkages Between Ecosystems: the South Florida Hydroscape*, St. Lucie Press; the other is in special issue of *Estuarine, Coastal and Shelf Science* on visualization in coastal marine science. Another manuscript is being prepared in conjunction with the FKNMS seagrass monitoring program for submission next year. We also maintain a web site at <http://www.fiu.edu/~serp/jrpp/wqmn/datamaps/datamaps.html> where data from the FKNMS is integrated with the other parts of the SERP water quality network (Florida Bay, Whitewater Bay, Biscayne Bay, Ten Thousand Islands, and SW Florida Shelf) and displayed as downloadable contour maps.

The period of record for this report is March 95 - October 98 and includes data from 13 quarterly sampling events at 154 stations within the FKNMS including the Dry Tortugas National Park. Water quality variables measured at each station include the dissolved nutrients nitrate ( $\text{NO}_3^-$ ), nitrite ( $\text{NO}_2^-$ ), ammonium ( $\text{NH}_4^+$ ), total dissolved inorganic nitrogen (DIN), and soluble reactive phosphate (SRP as  $\text{PO}_4^-$ ). Total unfiltered concentrations of organic nitrogen (TON), organic carbon (TOC), phosphorus (TP), and silicate ( $\text{Si}(\text{OH})_4$ ) were also measured. Biological parameters monitored include chlorophyll *a* (Chl *a*) and alkaline phosphatase activity (APA). Field parameters from each station include salinity, temperature, dissolved oxygen (DO), turbidity (NTU), relative fluorescence (RFU) and light attenuation ( $K_d$ ).

Stratification of sites according to depth showed that temperature, DO, TOC, and TON were significantly higher at the surface while salinity,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ , TP, and turbidity were higher in bottom waters. In April a mass of colder, nutrient laden water from the Gulf of Mexico moved up onto the Tortugas reefs and fueled a large benthic macroalgae bloom. This event was observed throughout most of the eastern Gulf as far north as Pensacola. Bottom water temperatures at the two most SW stations dropped  $\sim 4^\circ\text{C}$ ,  $\text{NO}_3^-$  increased 3 orders of magnitude,

SRP and  $\text{Si(OH)}_4$  increased by a factor of 100, while TP, turbidity, and Chl *a* specific fluorescence doubled.

Stratification according to geographical region showed that the Tortugas (Segment 1) and the Upper Keys (Seg. 9) generally had lower nutrient concentrations than the Middle Keys (Seg. 7) or Lower Keys (Seg. 5). In the Lower Keys DIN and TP was elevated in the Backcountry (Seg. 4). The Sluiceway (Seg. 6) had lowest salinity and highest TOC, TON, and  $\text{Si(OH)}_4$  concentrations. The offshore Marquesas (Seg. 2) area exhibited highest phytoplankton biomass (Chl *a*) for any segment of the FKNMS. Declining inshore to offshore trends were observed for  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ ,  $\text{Si(OH)}_4$ , TOC, TON, and turbidity for all oceanside transects. TP concentrations in the Lower Keys transects decreased with distance offshore but increased along transects in the Upper Keys, mostly because of low concentrations alongshore. Stations grouped by shore type showed that those stations situated along channels/passes through the Keys had higher nutrient concentrations, phytoplankton biomass, and turbidity than those stations off land. The absolute differences were very small and were not likely to be biologically important.

The water quality of the FKNMS is put in perspective with data from 150 stations sampled in SW Florida Shelf, Florida Bay, Whitewater Bay, Ten Thousand Islands, and Biscayne Bay. It becomes clear from this analysis that the ambient water quality in the Lower Keys and Marquesas is most strongly influenced by water quality of the SW Florida Shelf, the Middle Keys by SW Florida Shelf and Florida Bay transport, the Backcountry by internal nutrient sources, and the Upper Keys from Florida Current intrusion.

## II. Project Background

This report includes water quality data collected on a quarterly basis from 150 stations within the Florida Keys National Marine Sanctuary (Fig. 1). The period of record for this report is from Mar. 95 - Oct. 98 and includes 13 sampling events. A summary of the stations, their latitudes and longitudes, and the period of record surveyed are enclosed in Table 1.

Water quality parameters monitored at each station include the dissolved nutrients nitrate+nitrite ( $\text{NO}_x^-$ ), nitrite ( $\text{NO}_2^-$ ), nitrate ( $\text{NO}_3^-$ ), ammonium ( $\text{NH}_4^+$ ), total inorganic nitrogen (DIN), and soluble reactive phosphate (SRP). Total concentrations of nitrogen (TN), organic nitrogen (TON), organic carbon (TOC), and phosphorus (TP) were also measured. Concentrations for each of these parameters are reported  $\mu\text{mol l}^{-1}$  ( $\mu\text{M}$ ). Biological parameters monitored include chlorophyll *a* (Chl *a*;  $\mu\text{g l}^{-1}$ ) and alkaline phosphatase activity (APA;  $\mu\text{M hr}^{-1}$ ). Field parameters determined by CTD casts for both the surface and bottom of the water column at each station include salinity (ppt), dissolved oxygen (DO;  $\text{mg l}^{-1}$ ), temperature ( $^{\circ}\text{C}$ ), light attenuation ( $K_d$ ;  $\text{m}^{-1}$ ), and in vivo chlorophyll *a* fluorescence (FSU). Turbidity as both optical backscatterance (OBS) and nephelometry (NTU) was also measured at each station. We continue to develop a curvilinear relationship between OBS and NTU for comparison with our pre-existing dataset from other regions of South Florida.

Summary statistics for all measured parameters split out by segment are shown in Table 2. This summary includes data from all sampling dates and stations for the period of record listed by median value (Median), minimum value (Min.), maximum value (Max.), and number of samples (*n*). Typical water quality data is skewed to the low end which results in a non-normal distribution. Therefore it is more appropriate to use the median as the measure of central tendency.




Table 1: Florida Keys National Marine Sanctuary Station Summary

Station Name	Station #	Latitude	Longitude	Period of Record	Survey #
Fowey Rocks	200	25 35.400	-80 06.000	Mar. 95 - July 98	1 - 13
Sands Key	201	25 30.600	-80 09.900	Mar. 95 - July 98	1 - 13
Bowles Bank	202	25 29.628	-80 09.006	Mar. 95 - July 98	1 - 13
Triumph Reef	203	25 28.832	-80 07.061	Mar. 95 - July 98	1 - 13
Elliott Key	204	25 25.849	-80 11.732	Mar. 95 - July 98	1 - 13
Margo Fish Shoai	205	25 25.679	-80 10.920	Mar. 95 - July 98	1 - 13
Ajax Reef	206	25 24.898	-80 08.580	Mar. 95 - July 98	1 - 13
Old Rhodes Key	207	25 21.866	-80 12.905	Mar. 95 - July 98	1 - 13
Old Rhodes Key Channel	208	25 21.551	-80 12.031	Mar. 95 - July 98	1 - 13
Channel Key	209	24 35.142	-81 43.363	Mar. 95 - July 98	1 - 13
Old Rhodes Key Reef	210	25 19.500	-80 09.816	Mar. 95 - July 98	1 - 13
Pennikamp G27	211	25 18.473	-80 15.699	Mar. 95 - July 98	1 - 13
Turtle Harbor	212	25 17.671	-80 14.725	Mar. 95 - July 98	1 - 13
Turtle Reef	213	25 15.787	-80 11.264	Mar. 95 - July 98	1 - 13
Port Elizabeth	214	25 14.772	-80 17.951	Mar. 95 - July 98	1 - 13
Carysfort Channel	215	25 14.200	-80 17.338	Mar. 95 - July 98	1 - 13
Carysfort Reef	216	25 13.200	-80 13.000	Mar. 95 - July 98	1 - 13
Rafflesnake Key	217	25 10.399	-80 20.391	Mar. 95 - July 98	1 - 13
White Bank	218	25 09.538	-80 17.673	Mar. 95 - July 98	1 - 13
The Elbow	219	25 08.500	-80 15.500	Mar. 95 - July 98	1 - 13
Radabob Key	220	25 06.813	-80 22.405	Mar. 95 - July 98	1 - 13
Radabob Key Channel	221	25 06.615	-80 21.653	Mar. 95 - July 98	1 - 13
Dixie Shoal	222	25 04.342	-80 18.766	Mar. 95 - July 98	1 - 13
Mosquito Bank	223	25 03.611	-80 25.387	Mar. 95 - July 98	1 - 13
Molasses Reef Channel	224	25 01.849	-80 23.716	Mar. 95 - July 98	1 - 13
Molasses Reef	225	25 01.000	-80 22.045	Mar. 95 - July 98	1 - 13
Tavernier Harbor	226	25 00.504	-80 30.08g	Mar. 95 - July 98	1 - 13
Triangles	227	25 00.100	-80 28.494	Mar. 95 - July 98	1 - 13
Conch Reef	228	2458.600	-8027.000	Mar. 95 -July 98	1 - 13
Plantation Point	229	24 57.007	-80 33.731	Mar. 95 - July 98	1 - 13
The Rocks	230	24 56.379	-80 33.097	Mar. 95 - July 98	1 - 13
Davis Reef	231	2455.500	-80 30.200	Mar. 95 -July 98	1 - 13
Upper Matecumbe Key	232	24 55.622	-80 35.911	Mar. 95 - July 98	1 - 13
Upper MateCumbe Chnl	233	24 54.111	-80 34.943	Mar. 95 - July 98	1 - 13
Fish Haven	234	24 51.766	-80 34.018	Mar. 95 - July 98	1 - 13
Indian Key	235	24 52.274	-80 40.119	Mar. 95 - July 98	1 - 13
Indian Key Channel	236	24 51.582	-80 39.744	Mar. 95 - July 98	1 - 13
Indian Key Offshore	237	24 49.959	-80 38.944	Mar. 95 - July 98	1 - 13
Matecumbe Harbor	238	24 50.065	-80 44.810	Mar. 95 - July 98	1 - 13
Lower Matecumbe Chnl	239	24 48.134	-80 42.972	Mar. 95 - July 98	1 - 13
Matecumbe Offshore	240	24 47.072	-80 42.342	Mar. 95 - July 98	1 - 13
Long Key	241	24 48.148	-80 48.534	Mar. 95 - July 98	1 - 13
Long Key Channel	242	24 47.490	-80 48.735	Mar. 95 - July 98	1 - 13
Tennessee Reef	243	24 44.700	-80 47.000	Mar. 95 - July 98	1 - 13
Long Key Pass Inshore	244	24 47.600	-80 51.800	Mar. 95 - July 98	1 - 13
Long Key Pass Channel	245	24 45.700	-80 51.700	Mar. 95 - July 98	1 - 13
Long Key Pass Offshore	246	24 42.790	-80 51.488	Mar. 95 - July 98	1 - 13
Key Colony Beach	247	24 42.791	-81 00.669	Mar. 95 - July 98	1 - 13
Coffins Patch Channel	248	24 41.500	-80 58.000	Mar. 95 - July 98	1 - 13
Coffins Patch Offshore	249	24 40.500	-80 57.500	Mar. 95 - July 98	1 - 13
Seven Mile Bridge	250	2441.516	-81 10.947	Mar. 95 - July 98	1 - 13
Seven Mile Br. Channel	251	24 38.701	-81 09.905	Mar. 95 - July 98	1 - 13
Seven Mile Br. Offshore	252	24 36.416	-81 09.265	Mar. 95 - July 98	1 - 13
Spanish Harbor Keys	253	2438.722	-81 18.434	Mar.95 - July 98	1 - 13
Bahia Honda Key	254	2439.475	-81 15.901	Mar. 95 - July 98	1 - 13



Station Name	Station#	Latitude	Longitude	Period of Record	Survey#
Bahia Honda Channel	255	24 37.301	-81 14.585	Mar. 95 - July 98	1 - 13
Bahia Honda Offshore	256	2435.098	-81 14.045	Mar. 95 - July 98	1 - 13
Long Beach	257	2438.000	-81 20.239	Mar. 95 - July 98	1 - 13
Big Pine Channel	258	2435.594	-81 19.453	Mar. 95 - July 98	1 - 13
Big Pine Shoal	259	2434.222	-81 19.300	Mar. 95 - July 98	1 - 13
Newfound Harbor Keys	260	24 37.530	-81 24.997	Mar. 95 - July 98	1 - 13
American Shoal Channel	261	24 35.000	-81 30.000	Mar. 95 - July 98	1 - 13
Looe Key Channel	262	24 35.171	-81 25.132	Mar. 95 - July 98	1 - 13
Looe Key	263	2432.906	-81 23.845	Mar. 95 - July 98	1 - 13
Conch Reef	264	24 56.806	-80 27.568	Mar. 95 - July 98	1 - 13
	265	24 49.000	-81 05.000	Mar. 95 - July 98	1 - 13
Tarpon Creek	266	24 36.663	-81 30.316	Mar. 95 - July 98	1 - 13
American Shoal	267	2431.805	-81 29.238	Mar. 95 - July 98	1 - 13
Saddlebunch Keys	268	24 35.269	-81 34.619	Mar. 95 - July 98	1 - 13
West Washerwoman	269	24 33.566	-81 33.573	Mar. 95 - July 98	1 - 13
Maryland Shoal	270	2431.297	-81 33.859	Mar. 95 - July 98	1 - 13
Boca Chica Key	271	24 33.400	-81 40.000	Mar. 95 - July 98	1 - 13
Eastern Sambo	272	24 31.000	-81 39.700	Mar. 95 - July 98	1 - 13
Eastern Sambo Offshore	273	2429.575	-81 39.423	Mar. 95 - July 98	1 - 13
Boca Chica Channel	274	24 32.500	-81 45.000	Mar. 95 - July 98	1 - 13
BocaChica Mid	275	2431.286	-81 44.653	Mar. 95 - July 98	1 - 13
Boca Chica Offshore	276	24 28.523	-81 44.245	Mar. 95 - July 98	1 - 13
Key West Cut A	277	24 31.057	-81 49.266	Mar. 95 - July 98	1 - 13
Western Head	278	24 29.917	-81 49.137	Mar. 95 - July 98	1 - 13
Main Ship Channel	279	2427.482	-81 48.778	Mar. 95 - July 98	1 - 13
Eastern Dry Rocks	280	24 27.218	-81 50.619	Mar. 95 - July 98	1 - 13
Middle Ground	281	2429.700	-81 53.500	Mar. 95 - July 98	1 - 13
Arsenic Bank	282	24 52.000	-80 53.100	Mar. 95 - July 98	1 - 13
	283	24 54.000	-81 00.000	Mar. 95 - July 98	1 - 13
Tripod Bank	284	24 52.600	-80 57.500	Mar. 95 - July 98	1 - 13
Channel Key Pass	285	24 48.599	-80 55.217	Mar. 95 - July 98	1 - 13
Toms Harbor Cut	286	24 47.000	-80 55.437	Mar. 95 - July 98	1 - 13
Bamboo Banks	287	2450.800	-81 00.500	Mar. 95 - July 98	1 - 13
	288	24 53.000	-81 05.000	Mar. 95 - July 98	1 - 13
Bamboo Key	289	2446.665	-81 01.980	Mar. 95 - July 98	1 - 13
Bluefish Bank	290	2451.760	-81 12.974	Mar. 95 - July 98	1 - 13
Bullard Bank	291	2449.009	-81 08.807	Mar. 95 - July 98	1 - 13
John Sawyer Bank	292	2445.787	-81 06.749	Mar. 95 - July 98	1 - 13
Bethel Bank	293	2444.252	-81 06.558	Mar. 95 - July 98	1 - 13
Red Bay Bank	294	2445.330	-81 10.331	Mar. 95 - July 98	1 - 13
Bullfrog Banks	295	2451.098	-81 21.074	Mar. 95 - July 98	1 - 13
W. Bahia Honda Key	296	2447.491	-81 17.345	Mar. 95 - July 98	1 - 13
Cocconut Key	297	2444.446	-81 14.000	Mar. 95 - July 98	1 - 13
Harbor Key Bank	298	24 50.451	-81 25.867	Mar. 95 - July 98	1 - 13
Bogie Channel	299	24 41.242	-81 20.509	Mar. 95 - July 98	1 - 13
Little Pine Key	300	2442.529	-81 18.554	Mar. 95 - July 98	1 - 13
Cutoe Key	301	24 46.355	-81 23.799	Mar. 95 - July 98	1 - 13
Content Passage	302	24 48.077	-81 28.936	Mar. 95 - July 98	1 - 13
Pine Channel	303	24 41.545	-81 23.519	Mar. 95 - July 98	1 - 13
Toptree Hammock Chan.	304	24 42.732	-81 27.094	Mar. 95 - July 98	1 - 13
Cudjoe Key	305	24 46.111	-81 33.300	Mar. 95 - July 98	1 - 13
Johnson Key Channel	306	2444.262	-81 35.548	Mar. 95 - July 98	1 - 13
Tarpon Belly Keys	307	2443.502	-81 31.231	Mar. 95 - July 98	1 - 13
Kemp Channel	308	2441.658	-81 29.292	Mar. 95 - July 98	1 - 13
Snipe Point	309	2441.863	-81 40.991	Mar. 95 - July 98	1 - 13
Snipe Keys	310	24 39.451	-81 37.959	Mar. 95 - July 98	1 - 13

Station Name	Station #	Latitude	Longitude	Period of Record	Survey #
Shark Key	311	24 37.057	-81 39.005	Mar. 95 - July 98	1 - 13
E. Harbor Key Channel	312	24 39.587	-81 45.806	Mar. 95 - July 98	1 - 13
Lower Harbor Keys	313	24 38.494	-81 43.926	Mar. 95 - July 98	1 - 13
Howe Key Channel	314	24 45.026	-81 26.147	Mar. 95 - July 98	1 - 13
Calda Channel	315	24 37.995	-81 49.696	Mar. 95 - July 98	1 - 13
Man of War Harbor	316	24 35.146	-81 48.144	Mar. 95 - July 98	1 - 13
Garrison Bight	317	24 34.500	-81 47.300	Mar. 95 - July 98	1 - 13
KY Northwest Channel	318	24 39.050	-81 54.280	Mar. 95 - July 98	1 - 13
N Boca Grande Channel	319	24 37.740	-82 04.280	Mar. 95 - July 98	1 - 13
Loggerhead Marker	320	24 39.100	-82 58.000	Mar. 95 - July 98	1 - 13
Loggerhead Channel	321	24 38.500	-82 56.400	Mar. 95 - July 98	1 - 13
Satan Shoal	322	24 27.500	-81 58.900	Mar. 95 - July 98	1 - 13
	323	24 42.300	-82 10.200	Mar. 95 - July 98	1 - 13
Ellis Rock	324	24 39.200	-82 09.500	Mar. 95 - July 98	1 - 13
SE Marquesas	325	24 32.500	-82 06.000	Mar. 95 - July 98	1 - 13
	326	24 42.500	-82 15.000	Mar. 95 - July 98	1 - 13
N Quicksands	327	24 37.000	-82 13.400	Mar. 95 - July 98	1 - 13
Marquesas Rock	328	24 27.240	-82 13.440	Mar. 95 - July 98	1 - 13
	329	24 43.000	-82 32.500	Mar. 95 - July 98	1 - 13
New Ground	330	24 39.920	-82 22.440	Mar. 95 - July 98	1 - 13
	331	24 30.410	-82 17.260	Mar. 95 - July 98	1 - 13
S Quicksands	332	24 32.130	-82 22.040	Mar. 95 - July 98	1 - 13
Half Moon Shoal	333	24 34.630	-82 29.480	Mar. 95 - July 98	1 - 13
	334	24 30.730	-82 27.020	Mar. 95 - July 98	1 - 13
	335	24 27.650	-82 22.990	Mar. 95 - July 98	1 - 13
	336	24 39.290	-82 38.780	Mar. 95 - July 98	1 - 13
Rebecca Shoal	337	24 35.480	-82 35.270	Mar. 95 - July 98	1 - 13
Garden Key	338	24 38.208	-82 52.620	Mar. 95 - July 98	1 - 13
	339	24 25.700	-82 30.650	Mar. 95 - July 98	1 - 13
	340	24 29.100	-82 39.200	Mar. 95 - July 98	1 - 13
Northwest Channel	341	24 41.900	-82 53.000	Mar. 95 - July 98	1 - 13
NE DTNP	342	24 42.360	-82 44.740	Mar. 95 - July 98	1 - 13
N DTNP	343	24 45.300	-82 51.300	Mar. 95 - July 98	1 - 13
Southwest Channel	344	24 33.800	-82 48.300	Mar. 95 - July 98	1 - 13
	345	24 22.740	-82 42.280	Mar. 95 - July 98	1 - 13
W DTNP	346	24 40.000	-83 00.500	Mar. 95 - July 98	1 - 13
Loggerhead Offshore	347	24 39.240	-82 59.800	Mar. 95 - July 98	1 - 13
Hospital Key	348	24 38.500	-82 51.300	Mar. 95 - July 98	1 - 13
Loggerhead Inshore	349	24 38.277	-82 55.868	Mar. 95 - July 98	1 - 13
	350	24 27.000	-82 52.000	Mar. 95 - July 98	1 - 13
Grecian Rocks	400	25 06.437	-80 18.382	Oct. 97 - July 98	10 - 13
Alligator Reef	401	24 51.200	-80 37.300	Oct. 97 - July 98	10 - 13
Sombrero Key	402	24 37.800	-81 06.700	Oct. 97 - July 98	10 - 13
Western Sambo	403	24 29.000	-81 42.000	Oct. 97 - July 98	10 - 13

Table 2: Summary Statistics for Surveys #1-13.

	Seg.	Surface				Bottom			
		Median	Min.	Max.	<i>n</i>	Median	Min.	Max.	<i>n</i>
NO <sub>3</sub>	1	0.070	0.000	0.710	119	0.110	0.000	3.660	139
	2	0.060	0.000	0.500	153	0.060	0.000	1.300	160
	4	0.190	0.000	4.420	273				
	5	0.150	0.000	1.550	300	0.130	0.000	1.480	253
	6	0.090	0.000	0.980	140				
	7	0.110	0.000	2.090	187	0.100	0.000	2.310	129
	9	0.080	0.000	1.660	335	0.090	0.000	0.630	219
NO <sub>2</sub>	1	0.030	0.000	0.120	193	0.040	0.000	0.810	195
	2	0.030	0.000	0.140	240	0.030	0.000	0.430	242
	4	0.060	0.010	0.350	299				
	5	0.050	0.000	0.160	392	0.050	0.000	0.200	308
	6	0.060	0.010	0.240	190				
	7	0.050	0.000	0.250	229	0.040	0.000	0.690	168
	9	0.040	0.000	0.190	406	0.040	0.000	0.170	279
NH <sub>4</sub>	1	0.275	0.020	1.890	194	0.260	0.010	1.140	195
	2	0.280	0.030	1.600	245	0.280	0.030	1.020	246
	4	0.480	0.130	2.420	299				
	5	0.310	0.000	1.750	391	0.310	0.010	2.380	308
	6	0.340	0.010	1.450	188				
	7	0.360	0.010	2.440	234	0.280	0.010	1.890	170
	9	0.280	0.040	2.450	407	0.285	0.020	1.880	280
TN	1	8.680	4.310	24.790	195	8.095	4.520	19.490	194
	2	9.530	4.190	25.920	246	9.370	4.910	16.900	246
	4	15.920	9.460	34.850	299				
	5	11.700	4.950	27.960	392	10.470	5.040	23.500	308
	6	16.100	6.990	32.630	190				
	7	12.630	5.490	46.350	232	10.960	4.780	27.400	170
	9	9.615	5.450	27.130	406	9.115	4.330	28.620	280
DIN	1	0.370	0.030	2.100	193	0.410	0.040	5.600	194
	2	0.360	0.060	1.690	245	0.370	0.060	1.790	246
	4	0.710	0.200	5.820	299				
	5	0.490	0.060	2.210	387	0.500	0.060	2.710	305
	6	0.470	0.100	1.940	190				
	7	0.530	0.060	3.670	230	0.425	0.060	2.930	168
	9	0.430	0.070	3.300	407	0.420	0.040	2.090	280
TON	1	8.320	3.640	24.120	194	7.600	2.250	18.830	193
	2	9.040	3.810	24.930	245	9.005	4.440	16.540	246
	4	15.040	8.630	33.050	299				
	5	11.160	4.480	27.280	387	9.985	4.010	22.930	304
	6	15.620	6.750	32.100	190				
	7	12.160	4.690	42.680	228	10.550	3.790	26.640	167
	9	9.200	4.980	24.850	406	8.620	3.900	28.200	279

	Seg.	Surface				Bottom				
		Median	Min.	Max.	<i>n</i>	Median	Min.	Max.	<i>n</i>	
TP	1	0.170	0.080	0.320	195	0.180	0.090	0.400	194	
	2	0.190	0.050	0.660	246	0.190	0.030	0.490	246	
	4	0.220	0.090	0.530	299					
	5	0.170	0.010	0.450	387	0.175	0.010	0.270	302	
	6	0.210	0.120	0.840	190					
	7	0.170	0.010	0.320	234	0.170	0.010	0.270	171	
	9	0.150	0.010	0.300	401	0.150	0.010	0.310	277	
	SRP	1	0.010	0.000	0.060	142	0.010	0.000	0.220	152
		2	0.010	0.000	0.090	192	0.010	0.000	0.070	185
4		0.010	0.000	0.170	247					
5		0.010	0.000	0.080	290	0.010	0.000	0.100	235	
6		0.010	0.000	0.140	140					
7		0.010	0.000	0.100	191	0.010	0.000	0.090	135	
9		0.010	0.000	0.120	319	0.010	0.000	0.090	220	
APA		1	0.030	0.010	0.200	135	0.030	0.010	0.080	134
		2	0.040	0.010	0.840	173	0.040	0.010	0.240	173
	4	0.070	0.010	0.490	296					
	5	0.050	0.010	0.370	389	0.040	0.010	0.430	306	
	6	0.080	0.010	0.520	190					
	7	0.060	0.010	0.430	225	0.050	0.010	0.490	163	
	9	0.055	0.010	0.450	384	0.050	0.010	0.200	263	
	Chl <i>a</i>	1	0.220	0.080	1.350	191				
		2	0.355	0.060	6.810	246				
4		0.290	0.070	1.460	299					
5		0.280	0.050	1.680	390					
6		0.250	0.070	1.970	189					
7		0.255	0.010	1.790	234					
9		0.200	0.040	1.430	407					
TOC		1	181.170	86.980	1054.790	193	170.880	89.380	883.100	193
		2	199.670	88.480	501.750	245	193.830	94.940	847.710	246
	4	244.000	136.000	1653.540	299					
	5	207.550	93.440	674.040	388	201.390	92.770	332.900	306	
	6	276.280	122.170	970.170	190					
	7	228.060	98.080	805.310	233	204.270	102.400	760.770	170	
	9	200.895	92.650	512.480	406	192.250	92.830	482.500	279	
	Si(OH) <sub>4</sub>	1	0.340	0.000	3.670	133	0.510	0.010	5.650	151
		2	0.580	0.000	4.990	193	0.750	0.000	6.920	187
4		1.810	0.010	20.020	241					
5		1.405	0.000	15.940	322	1.085	0.010	5.840	252	
6		5.320	0.080	127.110	161					
7		1.220	0.010	37.360	191	0.730	0.010	30.200	142	
9		0.410	0.000	12.990	303	0.375	0.000	11.360	204	

	Seg.	Surface				Bottom				
		Median	Min.	Max.	<i>n</i>	Median	Min.	Max.	<i>n</i>	
Turbidity	1	0.320	0.010	3.000	153	0.395	0.030	2.630	154	
	2	0.800	0.040	12.970	217	1.010	0.050	8.620	218	
	4	1.000	0.050	11.350	283					
	5	0.480	0.020	4.880	383	0.500	0.020	4.880	311	
	6	0.830	0.070	37.000	185					
	7	0.600	0.010	17.350	218	0.375	0.010	16.900	162	
	9	0.380	0.010	8.800	385	0.350	0.010	7.950	271	
	Salinity	1	36.200	32.300	36.600	192	36.300	34.000	37.000	193
		2	36.200	33.600	37.000	241	36.300	34.600	37.400	241
4		36.200	30.500	38.600	299	36.200	33.500	38.700	296	
5		36.200	33.600	38.600	365	36.200	33.400	38.600	362	
6		35.900	29.900	40.300	189	35.900	29.900	39.700	187	
7		36.100	5.500	38.400	216	36.100	5.100	38.900	218	
9		36.100	32.500	37.800	393	36.100	32.100	37.800	372	
Temp.		1	27.450	21.100	31.100	194	25.100	18.200	30.600	194
		2	27.500	19.700	32.300	242	26.700	19.000	32.200	242
	4	27.400	19.100	36.100	299	27.400	18.700	34.500	296	
	5	27.400	20.300	33.600	368	27.000	20.100	33.400	364	
	6	28.300	20.900	32.700	189	28.200	20.300	32.700	187	
	7	26.700	19.500	39.600	216	25.900	19.400	32.900	219	
	9	26.200	17.300	32.200	395	25.600	17.100	32.100	374	
	DO	1	6.100	4.600	10.400	190	6.250	4.500	7.300	180
		2	6.200	4.600	7.800	238	6.300	4.800	7.700	238
4		6.300	2.800	10.800	299	6.300	2.800	10.900	296	
5		6.000	3.000	10.800	361	6.100	3.000	9.300	357	
6		6.400	4.200	10.500	189	6.500	4.200	10.600	187	
7		6.200	4.400	8.300	214	6.200	4.400	8.400	213	
9		6.200	4.700	8.500	384	6.300	4.400	8.700	363	
K <sub>d</sub>		1	0.121	0.037	0.700	195				
		2	0.194	0.030	0.619	243				
	4	0.310	0.026	2.562	261					
	5	0.185	0.005	0.949	366					
	6	0.370	0.021	1.389	140					
	7	0.191	0.012	1.078	223					
	9	0.201	0.007	1.573	348					
	TN:TP	1	53.300	22.300	204.900	195				
		2	50.450	13.500	148.300	246				
4		72.300	25.600	199.300	299					
5		70.400	21.300	1107.700	387					
6		77.150	18.600	161.300	190					
7		75.500	28.800	1047.100	232					
9		65.200	26.000	1356.200	401					

	Seg.	Surface				Bottom				
		Median	Min.	Max.	<i>n</i>	Median	Min.	Max.	<i>n</i>	
DIN:SRP	1	39.500	3.500	451.000	142					
	2	29.600	2.100	676.000	191					
	4	74.000	3.200	997.300	247					
	5	50.800	5.300	612.000	285					
	6	48.750	2.300	707.000	140					
	7	63.500	1.400	624.000	187					
	9	52.800	3.500	935.300	319					
	DO <sub>sat</sub>	1	91.750	69.600	150.000	190	91.150	65.800	107.600	180
		2	92.300	68.300	114.900	238	92.700	72.000	113.500	238
4		92.500	43.500	169.900	299	92.950	43.500	171.400	296	
5		90.400	46.900	153.300	361	91.000	46.900	143.600	357	
6		94.800	65.000	148.200	189	95.300	62.700	149.600	187	
7		92.400	68.000	124.400	214	92.300	64.500	127.400	213	
9		93.600	72.300	126.100	384	93.900	67.700	128.700	363	

### **III. Results and Discussion**

We have submitted the following manuscript for publication in a book entitled *Linkages Between Ecosystems: the South Florida Hydroscape* edited by J. W. Porter and K. G. Porter. We print it in entirety but ask that it not be reproduced as it has not yet completed the peer review process. We welcome any questions or comments.

A view from the bridge: External and internal forces affecting the  
ambient water quality of the Florida Keys National Marine Sanctuary  
(FKNMS)

*Joseph N. Boyer<sup>1</sup> and Ronald D. Jones<sup>1,2</sup>*

<sup>1</sup>Southeast Environmental Research Program and <sup>2</sup>Department of Biological Sciences,  
Florida International University, Miami, FL 33199

boyerj@fiu.edu

305-348-4076 (office)

305-348-4096 (fax)

Running Head: Water Quality in the FKNMS

BOYER, J. N., AND R. D. JONES. (submitted) A view from the bridge: External and internal  
forces affecting the ambient water quality of the Florida Keys National Marine Sanctuary. *In*  
J. W. Porter and K. G. Porter (eds.), *Linkages Between Ecosystems: the South Florida  
Hydroscape*. St. Lucie Press.



### *Acknowledgments*

We thank all the field and laboratory technicians involved with this project including: J. Absten, O. Beceiro, N. Black, S. Bolanos, C. Covas, D. Diaz, T. Frankovich, B. Gilhooly, S. Kaczynski, E. Kotler, C. Menendez, S. Perez, P. Sterling, F. Tam, and especially Pete Lorenzo. We want to thank the captains and crew of the R/V Bellows and R/V Suncoaster of the Florida Institute of Oceanography for their professional support of the monitoring program. We also appreciate the laboratory and lodging support supplied by the Florida Program of the National Underwater Research Center, UNC-Wilmington and Keys Marine Laboratory. This project was possible due to continued funding by the US-EPA under Agreement #X994621-94-0. Funding other portions of the SERP Water Quality Monitoring Network was provided by the South Florida Water Management District and Everglades National Park (SFWMD/NPS Coop. Agreement C-7919 and NPS/SERP Coop. Agreement 5280-2-9017). This is contribution #XX of the Southeast Environmental Research Program at Florida International University.

This publication is dedicated to the memory of Cristina Menendez.

## Abstract

Preliminary results of a spatially extensive water quality monitoring program in the FKNMS are reported. Quarterly sampling at 150 fixed stations began in March 1995 and continue at present and consists of 12 sampling events. Stratification of sites according to depth showed that temperature, dissolved oxygen, total organic carbon, and total organic nitrogen were significantly higher at the surface while salinity, nitrate, nitrite, ammonium, total phosphorus, and turbidity were greater in bottom waters. Stratification according to geographical region showed that the Upper Keys generally had lower nutrient concentrations than the Middle or Lower Keys. In the Lower Keys inorganic nitrogen and total phosphorus was elevated in the Backcountry area. The offshore Marquesas/Quicksands area exhibited the highest phytoplankton biomass (chlorophyll *a*) for any segment of the FKNMS. Declining inshore to offshore trends were observed for nitrate, ammonia, silicate, total organic carbon, total organic nitrogen, and turbidity for all oceanside transects. Total phosphorus concentrations in the Lower Keys transects decreased with distance offshore but increased along transects in the Upper Keys, mostly because of low concentrations alongshore. Stations stratified as being off land or channel/pass showed that those stations situated along channels/passes through the Keys had higher nutrient concentrations, phytoplankton biomass, and turbidity than those stations off land. The differences were statistically significant but the absolute differences were very small and not likely to be biologically important. The water quality of the FKNMS is put in perspective with data from 150 stations sampled in SW Florida Shelf, Florida Bay, Whitewater Bay, Ten Thousand Islands, and Biscayne Bay. It becomes clear from this analysis that the ambient water quality in the Lower Keys and Marquesas is most strongly influenced by water quality of the SW Florida Shelf, the Middle Keys by SW Florida Shelf and Florida Bay transport, the Backcountry by internal nutrient sources, and the Upper Keys from Florida Current intrusion.

## **Introduction**

The Florida Keys are a archipelago of sub-tropical islands of Pleistocene origin which extend in a NE to SW direction from Miami to Key West and out to the Dry Tortugas (Fig. 1). In 1990, President Bush signed into law the Florida Keys National Sanctuary and Protection Act (HR5909) which designated a boundary encompassing >2,800 square nautical miles of islands, coastal waters, and coral reef tract as the Florida Keys National Marine Sanctuary (FKNMS). The Comprehensive Management Plan (NOAA 1995) required the FKNMS to have a Water Quality Protection Plan (WQPP) thereafter developed by EPA and the State of Florida (EPA 1995). The contract for the water quality monitoring component of the WQPP was subsequently awarded to the Southeast Environmental Research Program at Florida International University and the field sampling program began in March 1995.

The waters of the FKNMS are characterized by complex water circulation patterns over both spatial and temporal scales with much of this variability due to seasonal influence in regional circulation regimes. The FKNMS is directly influenced by the Florida Current, the Gulf of Mexico Loop Current, inshore currents of the SW Florida Shelf (Shelf), and by tidal exchange with both Florida Bay and Biscayne Bay. Advection from these external sources has significant effects on the physical, chemical, and biological composition of waters within the FKNMS, as does internal nutrient loading and freshwater runoff from the Keys themselves. Water quality of the FKNMS may be directly affected both by external nutrient transport and internal nutrient loading sources. Therefore, the geographical boundary of the FKNMS must not be thought of as enclosing a distinct ecosystem but rather as being one of political/regulatory definition.

Ongoing quarterly sampling of >200 stations in the FKNMS and Shelf, as well as monthly sampling of 100 stations in Florida Bay, Biscayne Bay, and the mangrove estuaries of the SW coast, has provided us with a unique opportunity to explore the spatial component of water quality variability. By stratifying the sampling stations according to depth, regional geography, distance from shore, proximity to tidal passes, and influence of Shelf waters we report some preliminary conclusions as to the relative importance of external vs. internal factors on the ambient water quality within the FKNMS.

## Methods

### *Site Characteristics and Sampling Design*

A spatial framework for FKNMS water quality management was proposed on the basis of geographical variation of regional circulation (Klein and Orlando 1994). The final implementation plan (EPA 1995) partitioned the FKNMS into 9 segments which was collapsed to 7 for routine sampling (Fig. 1). Station locations were developed using a stratified random design along onshore/offshore transects in Seg. 5, 7, and 9 or within EMAP grid cells in Seg. 1, 2, 4, and 6.

Segment 1 (Tortugas) includes the Dry Tortugas National Park and surrounding waters and is most influenced by the Loop Current and Dry Tortugas Gyre. Segment 2 (Marquesas) includes the Marquesas Keys and a shallow sandy area between the Marquesas and Tortugas called the Quicksands; Segment 4 (Backcountry) contains the shallow, hard-bottomed waters of the gulfside Lower Keys. Segments 2 and 4 are both influenced by water moving south from the Shelf. Segment 6 can be considered as part of western Florida Bay. This area is referred to as the Sluiceway as it heavily influenced by transport from Florida Bay and the Shelf (Smith 1994). Segment 5 (Lower Keys), 7 (Middle Keys), and 9 (Upper Keys) include the inshore, Hawk Channel and reef tract of the Atlantic side of the Florida Keys. The Lower Keys are most influenced by cyclonic gyres spun off of the Florida Current, the Middle Keys by exchange with Florida Bay, while the Upper Keys are influenced by the Florida Current frontal eddies and to a certain extent by exchange with Biscayne Bay. All three oceanside segments are also influenced by wind and tidally driven lateral Hawk Channel transport (Pitts 1997).

### *Field Sampling*

The period of record of this study was from March 1995 to April 1998 and included 12 quarterly sampling events. For each event, field measurements and grab samples were collected from 150 fixed stations within the FKNMS boundary (Fig. 1). Depth profiles of temperature ( $^{\circ}\text{C}$ ), salinity (ppt), dissolved oxygen (DO,  $\text{mg l}^{-1}$ ), photosynthetically available radiation (PAR,  $\mu\text{E m}^{-2}$ ), in situ chlorophyll *a* specific fluorescence (FSU), optical backscatterance turbidity (OBS), depth as measured by pressure transducer (m), and density ( $\sigma_t$ ,  $\text{kg m}^{-3}$ ) were measured by CTD casts (Seabird SBE 19). The CTD was equipped with internal RAM and operated in stand alone mode at a sampling rate of 0.5 sec. The vertical light attenuation coefficient ( $K_d$ ,  $\text{m}^{-1}$ ) was

calculated at 0.5 m intervals from PAR and depth using the standard exponential equation (Kirk 1994) and averaged over the station depth. This was necessary due to periodic occurrence of optically distinct layers within the water column. During these events,  $K_d$  was reported for the upper layer.

In the Backcountry area (Seg. 4, Fig. 1) where it was too shallow to use a CTD, surface salinity and temperature were measured using a combination salinity-conductivity-temperature probe (Orion model 140). DO was measured using an oxygen electrode (Orion model 840) corrected for salinity and temperature. PAR was measured using a Li-Cor irradiance meter equipped with two  $4\pi$  spherical sensors (LI-193SB) separated by 0.5 m in depth and oriented at  $90^\circ$  to each other. The light meter measured instantaneous difference between sensors which was then used to calculate  $K_d$  from in-air surface irradiance.

Water was collected from ~0.25 m below the surface and at ~1 m from the bottom with a teflon-lined Niskin bottle (General Oceanics) except in the Backcountry where it was collected directly into sample bottles. Duplicate, unfiltered water samples were collected using 3x sample rinsed 120 ml HDPE bottles for analysis of total constituents. Duplicate water samples for dissolved nutrients were collected using 3x sample rinsed 150 ml syringes which were then filtered by hand through 25 mm glass fiber filters (Whatman GF/F) into 3x sample rinsed 60 ml HDPE bottles. The wet filters used for chlorophyll *a* (Chl *a*) analysis, were placed in 1.8 ml plastic centrifuge tubes to which 1.5 ml of 90 % acetone/water was added (Strickland and Parsons 1972).

Unfiltered samples were kept at ambient temperature in the dark during transport to the laboratory. During shipboard collection in the Tortugas/Marquesas and overnight stays in the Keys, unfiltered samples were analyzed for alkaline phosphatase activity (APA) and turbidity (see *Analytical*) prior to refrigeration. Filtered samples and filters were kept on ice in the dark during transport. During shipboard collection in the Tortugas/Marquesas and overnight stays in the lower Keys, filtrates and filters were frozen until analysis.

### *Laboratory Analysis*

Unfiltered water samples were analyzed for total organic carbon (TOC), total nitrogen (TN), total phosphorus (TP), silicate ( $\text{Si}(\text{OH})_4$ ), alkaline phosphatase activity (APA), and turbidity (NTU). TOC was measured by direct injection onto hot platinum catalyst in a Shimadzu TOC-5000 after first acidifying to  $\text{pH} < 2$  and purging with  $\text{CO}_2$ -free air. TN was measured using an ANTEK 7000N Nitrogen Analyzer using  $\text{O}_2$  as carrier gas to promote complete recovery of the nitrogen in the water samples (Frankovich and Jones 1998). TP was determined using a dry ashing, acid hydrolysis technique (Solórzano and Sharp 1980).  $\text{Si}(\text{OH})_4$  was measured using the molybdosilicate method (Strickland and Parsons 1972). The APA assay measures the activity of alkaline phosphatase, an enzyme used by bacteria to mineralize orthophosphate from organic compounds. The assay is performed by adding a known concentration of an organic phosphate compound (methylfluorescein phosphate) to an unfiltered water sample. AP in the water sample cleaves the orthophosphate, leaving methylfluorescein, a highly fluorescent compound. Fluorescence at initial and after 2 hr incubation were measured using a Gilford Fluoro IV Spectrofluorometer (excitation = 430 nm, emission = 507 nm) and subtracted to give APA in  $\mu\text{M h}^{-1}$  (Jones 1996). Turbidity was measured using an HF Scientific model DRT-15C turbidimeter and reported in NTU.

Filtrates were analyzed for soluble reactive phosphorus (SRP), nitrate+nitrite ( $\text{NO}_x^-$ ), nitrite ( $\text{NO}_2^-$ ), and total ammonia ( $\text{NH}_4^+$ ) on a four channel autoanalyzer (Alpkem model RFA 300). Filters for Chl *a* content ( $\mu\text{g l}^{-1}$ ) were allowed to extract for a minimum of 2 days at  $-20^\circ\text{C}$  before analysis. Extracts were analyzed using a Gilford Fluoro IV Spectrofluorometer (excitation = 435 nm, emission = 667 nm). All analyses were completed within 1 month after collection in accordance to SERP laboratory quality control guidelines.

Some parameters were not measured directly, but were calculated by difference. Nitrate ( $\text{NO}_3^-$ ) was calculated as  $\text{NO}_x^- - \text{NO}_2^-$ , dissolved inorganic nitrogen (DIN) as  $\text{NO}_x^- + \text{NH}_4^+$ , and total organic nitrogen (TON) defined as  $\text{TN} - \text{DIN}$ . All concentrations are reported as  $\mu\text{M}$  unless noted. All elemental ratios discussed were calculated on a molar basis. Percent DO saturation ( $\text{DO}_{\text{sat}}$  as %) was calculated using the equations of Garcia and Gordon (1992).

### *Statistical Analysis*

Stations were grouped four different ways for statistical analysis: by surface or bottom samples, surface by segment, surface by transect distance, and surface by shore type. These groupings were subjectively defined using best available knowledge in an effort to provide information as to source, transport, and fate of water quality components. For the first grouping, stations were selected as being >3 m depth where both surface and bottom samples were collected and stratified by depth. The second grouping included surface samples stratified by segment (Fig. 1) in accordance with the implementation plan (EPA 1995). The third grouping consisted of those surface stations situated on ocean-side transects being aggregated according to their distance from shore: Alongshore, Hawk Channel, or Reef Tract. In addition, we initiated a transect of stations in the Tortugas off Loggerhead Key to serve as a reference. Since sampling at these locations in the Tortugas were only recently set up to address this question, the data is more sparse. Also there are only two “channel” stations in the Tortugas which makes the data more susceptible to outlier conditions.

One of the concerns of this program is to determine the contribution of water movement through the passes of the Keys to the water quality of the reef. To this end we decided to characterize the last grouping of transects as to shore type: those that are adjacent to land off Biscayne National Park off Old Rhodes Key, Elliot Key and the Safety Valve (BISC), those that abut land in Key Largo, Middle, and Lower Keys (LAND), and those transects which are aligned along an open channel or pass through the Keys (PASS). These grouping strategies may be changed when enough data is collected (ca. 5-7 yr) to be analyzed using a statistically objective, multivariate approach as has been done previously for Florida Bay and Ten Thousand Islands (Boyer et al. 1997; Boyer and Jones 1998).

Typical water quality variables are usually skewed to the left resulting in non-normal distributions. Therefore it is more appropriate to use the median as the measure of central tendency. Data distributions of selected water quality variables are reported as box-and-whiskers plots. The box-and-whisker plot is a powerful statistic as it shows the median, range, the data distribution as well as serving as a graphical, nonparametric ANOVA. The center horizontal line of the box is the median of the data, the top and bottom of the box are the 25<sup>th</sup> and 75<sup>th</sup> percentiles (quartiles), and the ends of the whiskers are the 5<sup>th</sup> and 95<sup>th</sup> percentiles. The notch in

the box is the 95% confidence interval of the median. When notches between boxes do not overlap, the medians are considered significantly different. Outliers (<5<sup>th</sup> and >95<sup>th</sup> percentiles) were excluded from the graphs to reduce visual compression. Differences in variables were also tested between groups using the Wilcoxon Ranked Sign test (comparable to *t*-test) and among groups by the Kruskal-Wallis test (ANOVA) with significance set at  $P < 0.05$ .

In an effort to elucidate the contribution of external factors to the water quality of the FKNMS and to visualize gradients in water quality over the region, we combined data from other portions of our water quality monitoring network: Florida Bay, Biscayne Bay, Whitewater Bay, Ten Thousand Islands, and the Shelf (Fig. 1). Data from these 153 additional stations was collected during the same month as the FKNMS surveys and analyzed by SERP personnel using similar methodology and quality control as previously described. Spatial contour maps of median water quality variables (of 12 surveys) were generated in SURFER (Golden Software) using the Kriging algorithm.

## Results

### *Overall Water Quality*

Summary statistics for all surface water quality variables including data from all 12 sampling events and all stations for the period of record (Table 1) are shown as median value, minimum value, maximum value, and number of samples measured. Overall, the region was warm and euhaline with a median temperature of 26.7°C and salinity of 36.2 ppt. The median DO was 6.2 mg l<sup>-1</sup>, or ~93 % DO<sub>sat</sub>. On this coarse scale, the FKNMS exhibited very good water quality with median NO<sub>3</sub><sup>-</sup>, NH<sub>4</sub><sup>+</sup>, and TP concentrations of 0.10, 0.32, and 0.18 μM, respectively. NH<sub>4</sub><sup>+</sup> was the dominant DIN species in almost all of the samples (~70 %). DIN comprised a small fraction (4 %) of the TN pool with TON making up the bulk (11.4 μM). SRP concentrations were very low (median 0.01 μM) and comprised only 6 % of the TP pool. Chl *a* concentrations were very low overall, 0.25 μg l<sup>-1</sup>, but ranged from 0.01 to 6.8 μg l<sup>-1</sup>. Median turbidity was low (0.6 NTU) as reflected in a low K<sub>d</sub> (0.206 m<sup>-1</sup>). This resulted in a median photic depth (to 1 % incident PAR) of ~22 m. Molar ratios of N to P suggested a general P limitation of the water column; median TN:TP = 67 and median DIN:SRP = 49.



### *Stations Grouped by Depth*

Some general differences were observed for those stations >3 m depth where both surface and bottom concentrations were measured. Temperature, DO, TOC, and TON were significantly higher at the surface while salinity,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{NH}_4^+$ , TP, and turbidity were significantly higher in bottom waters. There were no significant differences in SRP, APA, or  $\text{Si}(\text{OH})_4$  with depth.

### *Stations Grouped By Segment*

$\text{NO}_3^-$  was highest in the Backcountry (0.18  $\mu\text{M}$ ) followed by the Lower and Middle Keys. Interestingly,  $\text{NO}_3^-$  concentrations in the Sluiceway (0.08  $\mu\text{M}$ ) were not significantly different than the Upper Keys, Tortugas, or Marquesas. Median  $\text{NO}_2^-$  concentrations in the Tortugas and Marquesas (0.03  $\mu\text{M}$ ) were significantly lower than for the other Keys segments.  $\text{NH}_4^+$  was highest in the Backcountry (0.46  $\mu\text{M}$ ) and lowest in the Tortugas, Marquesas, and Upper Keys (~0.25  $\mu\text{M}$ ). The Middle Keys had the most variability in DIN for any of the ocean-side segments.

TP was highest in the Backcountry and Sluiceway (~0.21  $\mu\text{M}$ ) and lowest in the Upper Keys (0.15  $\mu\text{M}$ ) with the remaining segments being intermediate (Fig. 2). SRP was very low (~0.01  $\mu\text{M}$ ) for all areas but was slightly elevated and most variable in the Marquesas and Backcountry. Median  $\text{Si}(\text{OH})_4$  concentrations were highest in the Sluiceway (4.9  $\mu\text{M}$ ), lowest in the Tortugas, Marquesas, and Upper Keys (~0.45  $\mu\text{M}$ ), and intermediate in the Backcountry, Lower, and Middle Keys (~1.4  $\mu\text{M}$ ). Consistently higher Chl *a* concentrations were observed in the Marquesas (0.36  $\mu\text{g l}^{-1}$ ) than for any other area of the FKNMS (Fig. 2). Lowest Chl *a* concentrations were found in the Upper Keys (0.20  $\mu\text{g l}^{-1}$ ).

The organic C and N pools as well as APA showed remarkable similarity in relative concentration among segments (Fig. 3). Highest TOC (~260  $\mu\text{M}$ ), TON (~15.5  $\mu\text{M}$ ), and APA (~0.08  $\mu\text{M h}^{-1}$ ) were observed in the Backcountry and Sluiceway which declined SW towards the Tortugas and NE towards the Upper Keys. Median  $\text{DO}_{\text{sat}}$  was relatively similar among segments but was significantly higher and more variable in the Backcountry and Sluiceway (~95 %). This result would not have been evident had we only reported DO in  $\text{mg l}^{-1}$  as it was not significant across segments.

Salinity was comparable for most segments but was slightly lower in the Sluiceway (Fig. 3). Salinity in the Sluiceway and Backcountry were highly variable and precluded any statistical discrimination from the other segments. Turbidity was highest in the Backcountry (1.0 NTU), Sluiceway (0.85 NTU), and Marquesas (0.83 NTU) with lowest turbidity occurring in the Tortugas segment (0.35 NTU, Fig. 3). The shallow Quicksands area in the Marquesas probably accounted for the elevated turbidity in this segment. The Middle Keys showed high variability in turbidity although the overall median was low.

The Tortugas, Marquesas, and Lower Keys had significantly higher water temperature ( $\sim 27.3^{\circ}\text{C}$ ) than the other segments (Fig. 4). Light attenuation showed a similar pattern as turbidity with highest  $K_d$  in Sluiceway ( $0.36\text{ m}^{-1}$ ) and Backcountry stations ( $0.31\text{ m}^{-1}$ ) and lowest  $K_d$  in the Tortugas ( $0.12\text{ m}^{-1}$ ). This works out to respective median photic depths of 13, 15, and 38 m. Median TN:TP ratios in the Tortugas (55) and Marquesas (52) were significantly lower than in the other segments (Fig. 4). Much of this difference was due to decreased TON concentrations in these areas rather than increased in TP. Lowest DIN:SRP ratios were found in the Marquesas (29) followed by the Tortugas (39). Decreased DIN as well as elevated SRP in the Marquesas relative to the Tortugas was responsible for these differences.

#### *Stations Grouped By Distance along Transect*

Median concentrations of  $\text{NO}_3^-$  in the Middle, and Lower Keys were significantly higher in Alongshore stations than those of Hawk Channel and Reef Tract (Fig. 5). Alongshore  $\text{NO}_3^-$  in the Upper Keys and Tortugas ( $\sim 0.1\text{ }\mu\text{M}$ ) was not nearly as high as found in the Middle and Lower Keys ( $\sim 2.5\text{ }\mu\text{M}$ ).  $\text{NO}_3^-$  concentrations on the Reef Tract and offshore in the Tortugas, Upper, and Middle Keys were comparable ( $\sim 0.05\text{ }\mu\text{M}$ ) and were all significantly lower than for the Lower Keys.  $\text{NH}_4^+$  concentrations followed similar trends as  $\text{NO}_3^-$  being higher in Alongshore stations in the Middle and Lower Keys and declined with distance offshore (Fig. 5). Alongshore  $\text{NH}_4^+$  was highest in the Middle Keys ( $\sim 0.5\text{ }\mu\text{M}$ ). No significant differences in  $\text{NH}_4^+$  was seen among Hawk Channel, Reef Tract, and Tortugas groups ( $\sim 0.3\text{ }\mu\text{M}$ ).

Alongshore TP concentrations ( $\sim 0.18\text{ }\mu\text{M}$ ) were significantly higher than the Reef Tract only in the Lower Keys (Fig. 5). The Middle Keys and Tortugas showed no offshore trend while the Upper Keys showed a slight increasing trend in TP from shore to reef. TP concentrations in the

Tortugas, Lower, and Middle Keys were comparable ( $\sim 0.18 \mu\text{M}$ ) while the Upper Keys were lowest overall ( $\sim 0.16 \mu\text{M}$ ). The major trends in TP were mirrored by SRP but were not statistically significant. Median  $\text{Si(OH)}_4$  concentrations dropped dramatically with distance offshore in the Middle Keys (Fig. 5). In the Lower Keys,  $\text{Si(OH)}_4$  was significantly lower only in the Reef Tract stations. There was no difference in  $\text{Si(OH)}_4$  concentrations in the Upper Keys or Tortugas transects. Alongshore  $\text{Si(OH)}_4$  concentrations were highest in the Middle Keys ( $\sim 3 \mu\text{M}$ ) while Reef Tract concentrations were highest in the Lower Keys ( $\sim 0.5 \mu\text{M}$ ).

There was no significant trend in Chl *a* with distance from land in the Lower Keys (Fig. 6), although there was a slight decline in the Middle Keys and a small increase in the Upper Keys. Chl *a* in the Offshore Tortugas sites was significantly lower than Alongshore and Channel sites and was comparable to levels in the Upper Keys ( $0.2 \mu\text{g l}^{-1}$ ). TOC in the Lower, Middle, and Upper Keys was elevated Alongshore and declined sequentially through Hawk Channel to the Reef Tract (Fig. 6). There was no significant difference in TOC within Tortugas groups ( $\sim 170 \mu\text{M}$ ) and was similar to Reef Tract concentrations in the Keys. Highest TOC in Alongshore stations occurred in the Middle Keys ( $\sim 250 \mu\text{M}$ ). TON concentrations exhibited similar patterns as TOC (data not shown).

Turbidity in all segments declined significantly with distance from land (Fig. 6). All Reef Tract and Offshore Tortugas sites had comparably low turbidity levels ( $\sim 0.2 \text{ NTU}$ ). Highest Alongshore turbidity was found in the Middle Keys ( $1.3 \text{ NTU}$ ). No significant differences in Alongshore turbidity in the Tortugas, Lower, and Upper Keys were observed ( $\sim 0.6 \text{ NTU}$ ). Trends in median salinity with distance offshore were small; trends in salinity variability were large (Fig. 6). Salinity from Alongshore to Reef Tract increased significantly in the Upper Keys whereas in the Lower Keys, salinity actually decreased offshore. No significant change in salinity was observed along Middle Keys, Marquesas, or Tortugas transects. In all segments, Alongshore salinities were much more variable than those of Reef Tract and Offshore. Reef Tract and Tortugas Offshore salinities were not significantly different, therefore, Alongshore salinity in the Lower Keys was higher than local seawater values while Alongshore salinity in the Upper Keys was depressed relative to local seawater values.

### *Stations Grouped By Shore Type*

Ocean-side transects showed marked differences in water quality when grouped by shore type (Fig. 7). Transects situated on open channels (Pass) through the Keys were elevated in  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , TP,  $\text{Si(OH)}_4$ , Chl *a*, TOC, and turbidity relative to those against the island chain (Land). Both salinity and temperature were significantly lower in Pass transects than for Land. Although these differences were statistically significant, the absolute differences were very small being only fractional. We also found that these effects diminished rapidly with distance offshore (data not shown). Interestingly, those transects located along Biscayne National Park (Old Rhodes Key, Elliot Key, the Safety Valve) were lowest of all for  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , TP,  $\text{Si(OH)}_4$ , and Chl *a*.

### **Discussion**

Distinguishing internal from external sources of nutrients in the FKNMS is a difficult task. The finer discrimination of internal sources into natural and anthropogenic inputs is even more difficult but most important as anthropogenic inputs may be regulated and possibly controlled by management activities. Advective transport of nutrients through the FKNMS was not measured by the existing fixed sampling plan. However, nutrient distribution patterns may be compared to the regional circulation regimes in an effort to visualize the contribution of external sources and advective transport to internal water quality of the FKNMS.

Circulation in coastal South Florida is dominated by regional currents such as the Loop Current, Florida Current, and Tortugas Gyre and by local transport via Hawk Channel and along-shore Shelf movements (Klein and Orlando 1994). Regional currents may influence water quality over large areas by the advection of external surface water masses into and through the FKNMS (Lee et al. 1994) and by the intrusion of deep offshore ocean waters onto the reef tract as internal bores (Leichter et al. 1996). Local currents become more important in the mixing and transport of freshwater and nutrients from terrestrial sources (Smith 1994; Pitts 1997).

Spatial patterns of salinity in coastal South Florida show these major sources of freshwater to have more than just local impacts (Fig. 8). In Biscayne Bay, freshwater is released through the canal system operated by SFWMD; the impact is clearly seen to affect northern Key Largo by causing a depression in median salinity coupled with high variability in alongshore sites (Fig. 6 and 7). Freshwater entering NE Florida Bay via overland flow from Taylor Slough and C-111 basin in

ENP can be seen to mix in a SW direction. The extent of influence of freshwater from Florida Bay on alongshore salinity in the Keys is less than that of Biscayne Bay but it is more episodic.

Transport of low salinity water from Florida Bay does not affect the Middle Keys sites enough to depress the median salinity in this region but is manifested as increased variability (Fig. 6). On the west coast, the large influence of the Shark River Slough, which drains the bulk of the Everglades and exits through the Whitewater Bay - Ten Thousand Islands mangrove complex, is clearly seen as impacting the Shelf waters (Fig. 8). The mixing of Shelf waters with the Gulf of Mexico produces a salinity gradient in a SW direction which extends out to Key West. This freshwater source does not seem to impact the Backcountry because of its shallow nature but instead follows a trajectory of entering western Florida Bay and exiting out through the channels in the Middle Keys (Smith 1994). This net transport of lower salinity water from mainland to reef in open channels through the Keys is observed more so as an increase in the range and variability of salinity than as a large depression in salinity (Fig. 7).

In addition to surface currents there is evidence that internal tidal bores regularly impact the Key Largo reef tract (Leichter et al. 1996). Internal bores are episodes of higher density, deep water intrusion onto the shallower shelf or reef tract. Depending on their energy, internal tidal bores can promote stratification of the water column or cause complete vertical mixing as a breaking internal wave of subthermocline water. To determine the extent of stratification we calculated the difference between surface and bottom density,  $\Delta\sigma_t$ , where positive values denoted greater density of bottom water relative to the surface. The resulting graph of  $\Delta\sigma_t$  (Fig. 9), shows that the SW area of the Tortugas segment tends to experience the greatest frequency of stratification events. The decreased temperature and increased salinity in bottom waters from intrusion of deeper denser oceanic waters to this region may also account for increases in  $\text{NO}_3^-$ , TP, and SRP in these bottom waters as well. For example, in April 1998 a mass of colder, nutrient laden water from the Gulf of Mexico moved up onto the Tortugas reefs and fueled a large benthic macroalgae bloom (J. Porter, pers. comm.). This event was observed throughout most of the eastern Gulf as far north as Pensacola (NOAA ???). At the two most SW stations (Table 2), temperatures dropped  $\sim 4^\circ\text{C}$ ,  $\text{NO}_3^-$  increased 3 orders of magnitude, SRP and  $\text{Si(OH)}_4$  increased by a factor of 100, while TP, turbidity, and in vivo Chl *a* specific fluorescence (measured via CTD) all doubled. As there was only a small increase in  $\text{NH}_4^+$  during

this event we believe the general case of elevated  $\text{NH}_4^+$  and turbidity found in bottom waters throughout the FKNMS is most probably due to benthic flux and resuspension and not to subthermocline advection.

Surface  $\text{Si(OH)}_4$  concentrations exhibited a pattern similar to salinity (Fig. 10). The source of  $\text{Si(OH)}_4$  in this geologic area of carbonate rock and sediments is from siliceous periphyton (diatoms) growing in the Shark River Slough, Taylor Slough, and C-111 basin watersheds. Unlike the Mississippi River plume with Chl *a* concentrations of  $76 \mu\text{g l}^{-1}$  (Nelson and Dortch 1996), phytoplankton biomass on the Shelf ( $1\text{-}2 \mu\text{g l}^{-1}$  Chl *a*) was not sufficient to account for the depletion of  $\text{Si(OH)}_4$  in this area. Therefore,  $\text{Si(OH)}_4$  concentrations on the Shelf were rapidly depleted by mixing alone allowing  $\text{Si(OH)}_4$  to be used as a semi-conservative tracer of freshwater in this system (Ryther et al. 1967; Moore et al. 1986). Unlike Florida Bay and the west coast, there was very little  $\text{Si(OH)}_4$  loading to southern Biscayne Bay, mostly because the source of freshwater to this system is from canals which drain agricultural and urban areas of Dade County.

In the Lower and Middle Keys, it is clear that the source of  $\text{Si(OH)}_4$  to the nearshore Atlantic waters is through the Sluiceway and Backcountry.  $\text{Si(OH)}_4$  concentrations near the coast were elevated relative to the reef tract with much higher concentrations occurring in the Lower and Middle Keys than the Upper Keys (Fig. 5). There is an interesting peak in  $\text{Si(OH)}_4$  concentration in an area of the Sluiceway which is densely covered with the seagrass, *Syringodium* (Fourqurean, in press). We are unsure as to the source but postulate that it may be due to benthic flux.

Visualization of spatial patterns of  $\text{NO}_3^-$  concentration over South Florida waters provide an extended view of source gradients over the region (Fig. 11). Biscayne Bay, Florida Bay, and the Shark River area of the west coast exhibited high  $\text{NO}_3^-$  concentrations relative to the FKNMS and Shelf. Elevated  $\text{NO}_3^-$  in Biscayne Bay is the result of loading from both the canal drainage system and from inshore groundwater (Alleman et al. 1995; Meeder et al. 1997). The source of  $\text{NO}_3^-$  to Florida Bay is the Taylor Slough and C-111 basin (Boyer and Jones, in press; Rudnick et al., in press) while the Shark River Slough impacts the west coast mangrove rivers and out onto the Shelf (Rudnick et al., in press). The oceanside transects off Biscayne Bay in Seg. 9 exhibited the lowest  $\text{NO}_3^-$  alongshore compared to the Middle and Lower Keys. A similar pattern was observed in a previous transect survey from these areas (Szmant and Forrester 1996). They also showed an inshore elevation of  $\text{NO}_3^-$  relative to Hawk Channel and the reef tract which is also

demonstrated in our analysis (Fig. 5). Interestingly,  $\text{NO}_3^-$  concentrations in all stations in the Tortugas transect were similar to those of reef tract sites in the mainland Keys; there was no inshore elevation of  $\text{NO}_3^-$  on the transect off uninhabited Loggerhead Key.

Figure 10 also shows that a distinct intensification of  $\text{NO}_3^-$  occurs in the Backcountry region. Part of this increase may be due to local sources of  $\text{NO}_3^-$ , i.e. septic systems and stormwater runoff around Big Pine Key (Lapointe and Clark 1992). However, there is another area, the Snipe Keys, that exhibits high  $\text{NO}_3^-$  which is uninhabited by man. This rules out the premise of septic systems being the only source of  $\text{NO}_3^-$  in this area. It is important to note that the Backcountry area is very shallow (~0.5 m) and hydraulically isolated from the Shelf and Atlantic which results in its having a relatively long water residence time. Elevated  $\text{NO}_3^-$  concentrations may be partially due to simple evaporative concentration as is seen in salinity (Fig. 8).

Total dissolved  $\text{NH}_4^+$  concentrations were distributed in a similar manner as  $\text{NO}_3^-$  with highest concentrations occurring in Florida Bay, the Ten Thousand Islands, and the Backcountry (data not shown).  $\text{NH}_4^+$  concentrations were very low in Biscayne Bay because it is not a major component of loading from the canal drainage system.  $\text{NH}_4^+$  also showed similarities with  $\text{NO}_3^-$  in its spatial distribution, being lowest in the Upper Keys and highest inshore relative to offshore. There was no alongshore elevation of  $\text{NH}_4^+$  concentrations in the Tortugas where levels were similar to those of reef tract sites in the mainland Keys. That the least developed portion of the Upper Keys in Biscayne National Park and uninhabited Loggerhead Key (Tortugas) exhibited lowest  $\text{NO}_3^-$  and  $\text{NH}_4^+$  concentrations is evidence of a local anthropogenic source for both of these variables along the ocean side of the Upper, Middle, and Lower Keys. This pattern of decline implies an onshore source which is diluted with distance from land by low nutrient Atlantic Ocean waters.

Elevated DIN concentrations in the Backcountry, on the other hand, are not so easily explained. We postulate that the high concentrations found there are due to a combination of anthropogenic loading, physical entrapment, and benthic  $\text{N}_2$  fixation. The relative contribution of these potential sources is unknown. Lapointe and Matzie (1996) have shown that stormwater and septic systems are responsible for increased DIN loading in and around Big Pine Key. The effect of increased water residence time in DIN concentration is probably small. Salinities in this area were only 1-2 ppt higher than local seawater which resulted in a concentration effect of only 5-

6%. Benthic  $N_2$  fixation may potentially be very important in the N budget of the Backcountry. Measured rates of  $N_2$  fixation in a *Thalassia* bed in Biscayne Bay, having very similar physical and chemical conditions, were  $540 \mu\text{mol N m}^{-2} \text{d}^{-1}$  (Capone and Taylor 1980). Without the plant community N demand, one day of  $N_2$  fixation has the potential to generate a water column concentration of  $>1 \mu\text{M NH}_4^+$  (0.5 m deep). Much of this  $\text{NH}_4^+$  is probably nitrified and may help account for the elevated  $\text{NO}_3^-$  concentrations observed in this area as well. Clearly,  $N_2$  fixation may be a significant component of the N budget in the Backcountry and that it may be exported as DIN to the FKNMS in general.

Spatial patterns in TP in South Florida coastal waters were strongly driven by the west coast outputs (Fig 12). A declining gradient in TP extended from the inshore waters of Whitewater Bay - Ten Thousand Islands mangrove complex out onto the Shelf and Tortugas. A declining gradient also extended from north central Florida Bay to the Middle Keys. Brand (1997) has postulated that groundwater from a subterranean Miocene quartz sand channel, "the river of sand", containing high levels of phosphorus is the source of TP in this region. However, no evidence of this source exists to date and the data from Florida Bay does not indicate a subterranean source either (Boyer and Jones unpublished data). A very small TP gradient was seen NE Florida Bay signifying that Taylor Slough and the C-111 basin contribute little TP to the system. Finally, there was no evidence of a significant terrestrial source of TP to Biscayne Bay.

In the Keys, there was evidence of elevated TP in alongshore stations of the Middle and Lower Keys but the differences were very small (Fig. 5). The Upper Keys actually showed higher TP concentrations on the reef tract than inshore implying an offshore source. Interestingly, the Tortugas area had higher TP concentrations than the Upper Keys as a result of Shelf water advection.

In South Florida coastal waters, very little of TP is found in the inorganic form (SRP -  $\text{PO}_4^-$ ); most is organic P (TOP). The distribution of SRP on the west coast and Shelf was similar to that of TP with the general gradient from the west coast to Tortugas remaining (Fig 13). However, the SRP distribution was distinctly different from that of TP in Florida Bay, Whitewater Bay, and Biscayne Bay. In central Florida Bay the N-S gradient previously observed for TP was highly diminished for SRP indicating that almost all the TP in central Florida Bay was in the form of TOP. It is unlikely that the source of TOP to this region is from overland flow or groundwater as



this is also the region that expresses highest salinity. Alternately, we hypothesize that the presence of the Flamingo channel, running parallel to the southern coastline of Cape Sable, acts as a tidal conduit for episodic advection of inshore Shelf water to enter north central Florida Bay. Subsequent trapping and evaporation then may act to concentrate TOP in this region. The second difference in P distributions was that there was a significant SRP gradient present in NE Florida Bay that was not observed for TP. The sources of SRP to this area are the Taylor Slough and C-111 basin (W. Walker per. communication; Boyer and Jones, in press; Rudnick et al., in press).

Whitewater Bay displayed an east-west gradient in SRP concentrations which increased with salinity leading us to conclude that the freshwater inputs from the Everglades were not a source of SRP to this area. Finally, there was evidence of a significant onshore-offshore SRP gradient in southern Biscayne Bay; most probably as a direct result of canal loading and groundwater seepage to this region (A. Lietz personal communication; Meeder et al. 1997).

Concentrations of TOC (Fig. 14) and TON (not shown) were remarkably similar in pattern of distribution across the South Florida coastal hydroscape. The decreasing gradient from west coast to Tortugas was very similar to that of TP. A steep gradient with distance from land was observed in Biscayne Bay. Both these gradients were due to terrestrial loading. On the west coast, the source of TOC and TON was from the mangrove forests. Our data from this area shows that concentrations of TOC and TON increased from Everglades headwaters through the mangrove zone and then decrease with distance offshore. In Biscayne Bay, much of the TOC and TON is from agricultural land use. The high concentrations of TOC and TON found in Florida Bay were due to a combination of terrestrial loading (Boyer and Jones, in press), in situ production by seagrass and phytoplankton, and evaporative concentration (Fourqurean et al. 1993).

Advection of Shelf and Florida Bay waters through the Sluiceway and passes accounted for this region and the inshore area of the Middle Keys as having highest TOC and TON of the FKNMS (Fig. 7). Strong offshore gradients in TOC and TON existed for all mainland Keys segments (Fig. 6) but not for the Tortugas transect. Part of this difference may be explained by the absence of mangroves in the single Tortugas transect. The higher concentrations of TOC and TON in the inshore waters of the Keys then implies a terrestrial source rather than simply benthic

production and sediment resuspension. Main Keys reef tract concentrations of TOC and TON were similar to those found in the Tortugas.

Much emphasis has been placed on assessing the impact of episodic phytoplankton blooms in Florida Bay on the offshore reef tract environment. Spatial patterns of Chl *a* concentrations (Fig. 15) showed that NW Florida Bay, Whitewater Bay, and the Ten Thousand Islands exhibited high levels of Chl *a* relative to Biscayne Bay, Shelf, and FKNMS. The highest Chl *a* concentrations were found in west coast mangrove estuaries (up to 45  $\mu\text{g l}^{-1}$  in Alligator Bay, TTI). Chl *a* is also routinely high ( $\sim 2 \mu\text{g l}^{-1}$ ) in NW Florida Bay along the channel connecting the Shelf to Flamingo, ENP. It is interesting that Chl *a* concentrations are higher in the Marquesas (0.36  $\mu\text{g l}^{-1}$ ) than in other areas of the FKNMS (Fig. 2). When examined in context with the whole South Florida ecosystem, it is obvious that the Marquesas zone should be considered a continuum of the Shelf rather than a separate management entity. This shallow sandy area (often called the Quicksands) acts as a physical mixing zone between the Shelf and the Atlantic Ocean and is a highly productive area for other biota as well as it encompasses the historical Tortugas shrimping grounds. A Chl *a* concentration of 1  $\mu\text{g l}^{-1}$  in the water column of a reef tract is considered a problem as it indicates potential of eutrophication. On the other hand, a similar Chl *a* level in the Quicksands indicates a productive shrimp fishery.

The oceanside transects in the Upper Keys (Seg. 9) exhibited the lowest overall Chl *a* concentrations of any zone in the FKNMS. Ocean transects showed a slight increase in Chl *a* on the reef tract in this area (Fig. 6). Transects off the Middle and Lower Keys showed that a drop in Chl *a* occurred only in the reef tract sites; there was no linear decline with distance from shore (data not shown). alongshore compared to the Middle and Lower Keys. Interestingly, Chl *a* concentrations in the Tortugas transect showed a similar pattern as the mainland Keys. Inshore and Hawk Channel Chl *a* concentrations among Middle Keys, Lower Keys and Tortugas sites were not significantly different. As inshore Chl *a* concentrations in the Tortugas were similar to those in the Middle and Lower Keys, we see no evidence of phytoplankton bloom transport from Florida Bay under this type of sampling design. There was however some slight evidence of increased Chl *a* in those stations situated along the major passes in the Keys relative to those abutting land (Fig. 7). The differences between these two groupings were very small (0.25 vs. 0.20  $\mu\text{g l}^{-1}$ ).

Along with P concentration, turbidity is probably the second most important determinant of local ecosystem health. The fine, low density carbonate sediments in this area are easily resuspended, rapidly transported, and have high light scattering potential per gram of material. High water column turbidity and transport directly affects filter feeding organisms by clogging their feeding apparatus and by increasing local sedimentation rate. Sustained high turbidity of the water column indirectly affects benthic community structure by decreasing light penetration, promoting seagrasses extinction. Large scale observations of turbidity clearly show patterns of onshore-offshore gradients which extend out onto the Shelf to the Marquesas (Fig. 16). In the last seven years, turbidities in Florida Bay have increased dramatically in the NE and central regions (Boyer et al. 1998) potentially as a consequence of destabilization of the sediment from seagrass die-off (Robblee et al. 1991).

Strong turbidity gradients were observed for all Keys transects (Fig. 6) but reef tract levels were remarkably similar regardless of inshore levels. High alongshore turbidity is most probably due to the shallow water column being easily resuspended by wind and wave action. Inshore stations in the Middle Keys had higher turbidity than other segments. Transects aligned with major passes had slightly greater turbidity than those against land but the difference was not statistically significant (Fig. 7). Light extinction ( $K_d$ ) was highest alongshore and improved with distance from land (data not shown). This trend was expected as light extinction is directly related to the turbidity of the water.

Using the DIN:SRP ratio is a relatively simple method of determining phytoplankton nutrient limitation status of the water column (Redfield 1967). Most of the South Florida hydroscape was shown to have DIN:SRP values  $\gg 16:1$ , indicating the potential for phytoplankton to be limited by P at these sites (Fig. 17). The bulk of Florida Bay and both southern and northern Biscayne Bay were severely P limited, mostly as a result of high DIN concentrations. All of the FKNMS is routinely P limited using this metric. Interestingly, the Marquesas/Quicksands area was the least P limited of all zones and exhibited a significant regression between SRP and Chl *a*. Only in the northern Ten Thousand Islands and Shelf did N become the limiting nutrient. The south-north shift from P to N limitation observed in the west coast estuaries has been ascribed to changes in landuse and bedrock geochemistry of the watersheds (Boyer and Jones 1998). The west coast south of 25.4 N latitude is influenced by overland freshwater flow from the Everglades and Shark

River Slough having very low P concentrations relative to N. Above 25.7 N latitude the bedrock geology of the watershed changes from carbonate to silicate based and landuse changes from relatively undeveloped wetland (Big Cypress Basin) to a highly urban/agricultural mix (Naples, FL).

The large scale of this monitoring program has allowed us to assemble a much more holistic view of broad physical/chemical/biological interactions occurring over the South Florida hydroscape. Much information has been gained by inference from this type of data collection program: major nutrient sources have been confirmed, relative differences in geographical determinants of water quality have been demonstrated, and large scale transport via circulation pathways have been elucidated. In addition we have shown the importance of looking "outside the box" for questions asked within. Rather than thinking of water quality monitoring as being a static, non-scientific pursuit it should be viewed as a tool for answering management questions and developing new scientific hypotheses. One of the more important management questions to be answered is "Is the water quality better or worse than it used to be?" As it stands, this monitoring program based on quarterly sample intervals may require up to 10 years before small trends may be detected because of seasonal variability and background noise.

## **References**

- ALLEMAN, R. W., ET AL. 1995. Biscayne Bay surface water improvement and management. Technical supporting document. South Florida Water Management District.
- BOYER, J. N., J. W. FOURQUREAN, AND R. D. JONES. 1997. Spatial characterization of water quality in Florida Bay and Whitewater Bay by multivariate analysis: Zones of similar influence (ZSI). *Estuaries* **20**: 743-758.

- BOYER, J. N., J. W. FOURQUREAN, AND R. D. JONES. 1998. Seasonal and long term trends in the water quality of Florida Bay (1989 - 1997). *Estuaries* (in press).
- BOYER, J. N., AND R. D. JONES. 1998. Influence of coastal morphology and watershed characteristics on the water quality of mangrove estuaries in the Ten Thousand Islands - Whitewater Bay complex. Proceedings of the 1998 Florida Bay Science Conference. University of Florida Sea Grant.
- BOYER, J. N., AND R. D. JONES. (in press) Effects of freshwater inputs and loading of phosphorus and nitrogen on the water quality of Eastern Florida Bay. *In* K. R. Reddy, G. A. O'Connor, and C. L. Schelske [eds.], Phosphorus biogeochemistry in sub-tropical ecosystems: Florida as a case example. CRC/Lewis Publishers.
- BRAND, L. 1998. The role of groundwater in the Florida Bay ecosystem. Proceedings of the 1998 Florida Bay Science Conference. University of Florida Sea Grant.
- CAPONE, D. G., AND B. F. TAYLOR. 1980. Microbial nitrogen cycling in a seagrass community, p. 153-161. *In* V. S. Kennedy [ed.], *Estuarine perspectives*. Academic.
- DEPARTMENT OF COMMERCE. 1995. Florida Keys National Marine Sanctuary draft management plan / environmental impact statement. National Oceanic and Atmospheric Administration, Sanctuary Programs Division. 245 p.
- ENVIRONMENTAL PROTECTION AGENCY. 1995. Water quality protection program for the Florida Keys National Marine Sanctuary: Phase III report. Final report submitted to the Environmental Protection Agency under Work Assignment 1, Contract No. 68-C2-0134. Battelle Ocean Sciences, Duxbury, MA and Continental Shelf Associates, Inc., Jupiter FL.
- FOURQUREAN, J. W. (submitted). *In* J. W. Porter and K. G. Porter (eds.), *Linkages Between Ecosystems: the South Florida Hydroscape*. St. Lucie Press.
- FOURQUREAN, J. W., R. D. JONES, AND J. C. ZIEMAN. 1993. Processes influencing water column nutrient characteristics and phosphorus limitation of phytoplankton biomass in Florida Bay, FL, USA: inferences from spatial distributions. *Estuar., Coast. Shelf Sci.* **36**: 295-314.
- FRANKOVICH, T. A., AND R. D. JONES. 1998. A rapid, precise, and sensitive method for the determination of total nitrogen in natural waters. *Mar. Chem.* **60**: 227-234.

- FROELICH, P. N., JR., D. K. ATWOOD, AND G. S. GIESE. 1978. Influence of Amazon River discharge on surface salinity and dissolved silicate concentrations in the Caribbean Sea. *Deep Sea Res.* **25**: 735-744.
- GARCIA, H. E., AND L. I. GORDON. 1992. Oxygen solubility in seawater: Better fitting equations. *Limnol. Oceanogr.* **37**: 1307-1312.
- JONES, R. D. 1996. Phosphorus cycling, p. 343-348. *In* G. J. Hurst, G. A. Knudsen, M. J. McInerney, L. D. Stetzenbach and M. V. Walter [eds.], *Manual of environmental microbiology*. ASM Press.
- KIRK, J. T. O. 1994. *Light and photosynthesis in aquatic systems*, 2nd Ed. Cambridge
- KLEIN, C. J., AND S. P. ORLANDO JR. 1994. A spatial framework for water-quality management in the Florida Keys National Marine Sanctuary. *Bull. Mar. Sci.* **54**: 1036-1044.
- LAPOINTE, B. E., AND M. W. CLARK. 1992. Nutrient inputs from the watershed and coastal eutrophication in the Florida Keys. *Estuaries* **15**: 465-476.
- LAPOINTE, B. E., AND W. R. MATZIE. 1996. Effects of stormwater nutrient discharges on eutrophication processes in nearshore waters of the Florida Keys. *Estuaries* **19**: 422-435.
- LEE, T. N., M. E. CLARKE, E. WILLIAMS, A. F. SZMANT, AND T. BERGER. 1994. Evolution of the Tortugas gyre and its influence on recruitment in the Florida Keys. *Bull. Mar. Sci.* **54**: 621-646.
- LEICHTER, J. J., S. R. WING, S. L. MILLER, AND M. W. DENNY. 1996. Pulsed delivery of subthermocline water to Conch Reef (Florida Keys) by internal tidal bores. *Limnol. Oceanogr.* **41**: 1490-1501.
- MEEDER, J. F., J. ALVORD, M. BYRNS, M. S. ROSS, AND A. RENSHAW. 1997. Distribution of benthic nearshore communities and their relationship to groundwater nutrient loading. Final report to Biscayne National Park.
- MOORE, W. S., J. L. SARMIENTO, AND R. M. KEY. 1986. Tracing the Amazon component of surface Atlantic water using  $^{228}\text{Ra}$ , salinity, and silica. *J. Geophys. Res.* **91**: 2574-2580.

- NELSON, D. M., AND Q. DORTCH. 1996. Silicic acid depletion and silicon limitation in the plume of the Mississippi River: evidence from kinetic studies in spring and summer. *Mar. Ecol. Prog. Ser.* **136**: 163-178.
- NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION. 1995. Florida Keys National Marine Sanctuary Draft Management Plan/Environmental Impact Statement.
- PITTS, P. A. 1997. An investigation of tidal and nontidal current patterns in Western Hawk Channel, Florida Keys. *Cont. Shelf Res.* **17**: 1679-1687.
- ROBBLEE, M. B., T. B. BARBER, P. R. CARLSON JR., M. J. DURAKO, J. W. FOURQUREAN, L. M. MUEHLSTEIN, D. PORTER, L. A. YABRO, R. T. ZIEMAN, AND J. C. ZIEMAN. 1991. Mass mortality of the tropical seagrass *Thalassia testudinum* in Florida Bay (USA). *Mar. Ecol. Prog. Ser.* **71**: 297-299.
- RUDNICK, D., Z. CHEN, D. CHILDERS, T. FONTAINE, AND J. N. BOYER. (in press) Phosphorus and nitrogen inputs to Florida Bay: the importance of the Everglades watershed. *Estuaries*.
- RYTHER, J. H., D. W. MENZE, AND N. CORWIN. 1967. Influence of the Amazon River outflow on the ecology of the western tropical Atlantic, I. Hydrography and nutrient chemistry. *J. Mar. Res.* **25**: 69-83.
- SOLORZANO, L., AND J. H. SHARP. 1980. Determination of total dissolved phosphorus and particulate phosphorus in natural waters. *Limnol. Oceanogr.* **25**: 754-758.
- SMITH, N. P. 1994. Long-term Gulf-to-Atlantic transport through tidal channels in the Florida Keys. *Bull. Mar. Sci.* **54**: 602-609.
- STRICKLAND, J. D. H., AND T. R. PARSONS. 1972. A practical handbook of seawater analysis. *Bull. Fish. Res. Bd. Can.* **167**: 107-112.
- SZMANT, A. M., AND A. FORRESTER. 1996. Water column and sediment nitrogen and phosphorus distribution patterns in the Florida Keys, USA. *Coral Reefs* **15**: 21-41.

## List of Tables

Table 1. Summary statistics for each water quality variable in the FKNMS. Data are summarized as median (Median), minimum value (Min.), maximum value (Max.), and number of samples ( $n$ ).

Table 2. Physical and chemical differences between surface and bottom waters on 25 April 1998 stratification event at 2 sites in the SW Tortugas (Segment 1) of the FKNMS.



Table 1

	Median	Min.	Max.	n
NO <sub>3</sub> <sup>-</sup>	0.08	0.00	1.61	862
NO <sub>2</sub> <sup>-</sup>	0.04	0.00	0.22	1180
NH <sub>4</sub> <sup>+</sup>	0.27	0.01	2.44	1191
TON	9.65	3.64	42.67	1180
TP	0.17	0.01	0.66	1183
SRP	0.01	0.00	0.12	915
APA	0.04	0.01	0.84	1029
Chl <i>a</i>	0.25	0.01	6.81	1188
TOC	198.70	86.98	1054.79	1186
Si(OH) <sub>4</sub>	0.59	0.00	37.36	902
Turbidity	0.46	0.01	12.97	1083
Salinity	36.20	31.80	38.40	1140
Temperature	26.45	17.50	39.60	1146
DO	6.20	4.60	10.40	1123
DO <sub>sat</sub>	92.50	68.28	150.03	1123
K <sub>d</sub>	0.16	0.01	1.12	1138
TN:TP	62.10	15.89	1356.21	1181
DIN:SRP	43.83	1.40	935.33	906
%I <sub>0</sub>	17.21	0.00	97.03	1138

Table 2

	Station #345		Station #350	
	Surface	Bottom	Surface	Bottom
Salinity	36.00	36.00	36.10	36.00
Temperature	24.00	19.70	23.60	20.20
NO <sub>3</sub> <sup>-</sup>	0.05	3.66	0.00	3.23
NO <sub>2</sub> <sup>-</sup>	0.01	0.81	0.04	0.35
NH <sub>4</sub> <sup>+</sup>	0.94	1.14	0.32	0.57
TP	0.23	0.39	0.19	0.40
SRP	0.02	0.22	0.01	0.20
Si(OH) <sub>4</sub>	0.09	2.07	0.29	2.26
TOC	170.10	155.48	171.42	168.35
TON	4.86	2.25	5.45	4.01
Chl <i>a</i> (fluor)	0.94	1.62	1.10	1.82
Turbidity	0.17	0.50	0.43	1.11
DO <sub>sat</sub>	94.28	80.91	93.90	78.74

## List of Figures

Fig. 1. The SERP Water Quality Monitoring Network showing the FKNMS boundary, segments, and distribution of sampling stations within the FKNMS, Florida Bay, Biscayne Bay, Whitewater Bay, Ten Thousand Islands, and Southwest Florida Shelf.

Fig. 2. Box-and-whisker plots of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , TP, SRP,  $\text{Si(OH)}_4$ , and Chl *a* stratified by FKNMS segment: 1 = Tortugas, 2 = Marquesas, 4 = Backcountry, 5 = Lower Keys, 6 = Sluiceway, 7 = Middle Keys, and 9 = Upper Keys.

Fig. 3. Box-and-whisker plots of TOC, TON, APA,  $\text{DO}_{\text{sat}}$ , salinity, and turbidity by FKNMS segment.

Fig. 4. Box-and-whisker plots of temperature,  $K_d$ , TN:TP, and DIN:SRP by FKNMS segment.

Fig. 5. Box-and-whisker plots of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , TP,  $\text{Si(OH)}_4$  by FKNMS segment and location on transect from land. In the Tortugas segment AS = alongshore, CH = channel, and OS = offshore.

Fig. 6. Box-and-whisker plots of Chl *a*, TOC, turbidity, and salinity by FKNMS segment and location on transect from land.

Fig. 7. Box-and-whisker plots of  $\text{NO}_3^-$ ,  $\text{NH}_4^+$ , TP,  $\text{Si(OH)}_4$ , Chl *a*, TOC, salinity, and turbidity for ocean-side Keys transects stratified as being located along Biscayne Bay National Park (BISC), against land in Key Largo, Middle and Lower Keys (LAND), or in line with major passes in the Keys (PASS).

Fig. 8. Contour map of median salinity from fixed stations (+) in South Florida coastal waters reported as the median for 12 sampling events.

Fig. 9. Contour map of median delta sigma-t ( $\Delta\sigma_t$  in  $\text{kg m}^{-3}$ ) as the difference in density between surface and bottom waters where positive values mean the bottom is more dense than surface.

Fig. 10. Contour map of median  $\text{Si(OH)}_4$  concentrations ( $\mu\text{M}$ ).

Fig. 11. Contour map of median  $\text{NO}_3^-$  concentrations ( $\mu\text{M}$ ).

Fig. 12. Contour map of median TP concentrations ( $\mu\text{M}$ ).


Fig. 13. Contour map of median SRP concentrations ( $\mu\text{M}$ ).







Fig. 14. Contour map of median TOC concentrations ( $\mu\text{M}$ ).





Fig. 15. Contour map of median Chl *a* concentrations ( $\mu\text{g l}^{-1}$ ).


Fig. 16. Contour map of median turbidity (NTU).

Fig. 17. Contour map of median DIN:SRP ratio.






























# Salinity Median for 1995-8




# delta Sigma-t Median for 1995-8




# Silicate Median for 1995-8




# Nitrate Median for 1995-8



# Total Phosphorus Median for 1995-8




# Soluble Reactive Phosphorus Median for 1995-8







# Total Organic Carbon Median for 1995-8




# Chlorophyll a Median for 1995-8



# Turbidity Median for 1995-8



# DIN:SRP Ratio Median for 1995-8



## **IV. List of Appendices**

Appendix 1. Color contour maps of selected water quality variables by sampling event.

These maps encompass all 354 stations of the SERP Water Quality Monitoring Network which includes the FKNMS, Biscayne Bay, Florida Bay, Whitewater Bay, Ten Thousand Islands, and Southwest Florida Shelf. The data was collected over a period of a month so care should be taken in interpreting these maps as they are not truly synoptic.

Appendix 2. All data from the 4 sampling events in 1998 reported in record format by survey, date, and station number.

## **Appendix 1 - Contour Maps**