2010

Size-based variation in intertissue comparisons of stable carbon and nitrogen isotopic signatures of bull sharks (*Carcharhinus leucas*) and tiger sharks (*Galeocerdo cuvier*)

Philip Matich

Department of Biological Sciences and Marine Sciences Program, Florida International University, pmati001@fiu.edu

Michael R. Heithaus

Department of Biological Sciences and Marine Sciences Program, Florida International University, heithaus@fiu.edu

Craig A. Layman

Marine Sciences Program, Florida International University

Follow this and additional works at: https://digitalcommons.fiu.edu/fce_lter_journal_articles

Recommended Citation

Matich, P., M.R. Heithaus, C.A. Layman. 2010. Size-based variation in intertissue comparisons of stable carbon and nitrogen isotopic signatures of bull sharks (*Carcharhinus leucas*) and tiger sharks (*Galeocerdo cuvier*). *Canadian Journal of Fisheries and Aquatic Sciences* 67: 877-885.

This material is based upon work supported by the National Science Foundation through the Florida Coastal Everglades Long-Term Ecological Research program under Cooperative Agreements #DBI-0620409 and #DEB-9910514. Any opinions, findings, conclusions, or recommendations expressed in the material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

This work is brought to you for free and open access by the FCE LTER at FIU Digital Commons. It has been accepted for inclusion in FCE LTER Journal Articles by an authorized administrator of FIU Digital Commons. For more information, please contact dcc@fiu.edu, jkrefft@fiu.edu.

**Size-based variation in inter-tissue comparisons of stable carbon and nitrogen
isotopic signatures of bull sharks and tiger sharks**

Philip Matich*, Michael R. Heithaus, and Craig A. Layman

Marine Sciences Program

Florida International University

3000 NE 151st

North Miami, FL 33181

Running head: isotope variation in shark tissues

*To whom correspondence should be addressed, pmati001@fiu.edu, (305) 919-5602 voice,
(305) 919-4030 fax

Abstract

Stable isotopes are an important tool for understanding the trophic roles of elasmobranchs. However, whether different tissues provide consistent stable isotope values within an individual are largely unknown. To address this, the relationships among carbon and nitrogen isotope values were quantified for blood, muscle, and fin from juvenile bull sharks (*Carcharhinus leucas*), and blood and fin from large tiger sharks (*Galeocerdo cuvier*) collected in two different ecosystems. We also investigated the relationship between shark size and the magnitude of differences in isotopic values between tissues. Isotope values were significantly positively correlated for all paired tissue comparisons, but R^2 values were much higher for $\delta^{13}\text{C}$ than $\delta^{15}\text{N}$. Paired differences between isotopic values of tissues were relatively small, but varied significantly with shark total length, suggesting shark size can be an important factor influencing the magnitude of differences in isotope values of different tissues. For studies of juvenile sharks, care should be taken in using slow turnover tissues like muscle and fin, because they may retain a maternal signature for an extended time. While correlations were relatively strong, results suggest correction factors should be generated for the desired study species, and may only allow course-scale comparisons between studies using different tissue types.

Key words:

Apex predator, *Carcharhinus leucas*, estuary, food webs, foraging ecology, *Galeocerdo cuvier*, stable isotopes

Introduction

Elasmobranchs (sharks, skates, and rays) play crucial roles in marine ecosystems (Heithaus et al. 2008), but gaps in our knowledge of their trophic interactions hinder understanding of marine community dynamics and ecosystem function. Current studies of trophic interactions of elasmobranchs, especially sharks, are particularly important because populations of many species are declining rapidly worldwide (e.g. Dulvy et al. 2008). These declines already may be causing drastic shifts in food web structure and function (Heithaus et al. 2008).

Most studies of elasmobranch trophic interactions have employed stomach content analysis (see Weatherbee and Cortes 2004 for a review). Although stomach content analysis allows identification of specific prey taxa, it has drawbacks, including the need for large sample sizes and often destructive sampling. Sharks also often have empty stomachs (Weatherbee and Cortes 2004), further limiting information that can be gleaned from this approach. Stable isotope analysis provides an alternative, or complementary, method for gaining insights into the trophic interactions of sharks (e.g. Fisk et al. 2002, Domi et al. 2005, MacNeil et al. 2005), especially because samples can be collected without sacrificing individuals. This method is based on the principle that a consumer's tissues isotopically resemble those of its food (Post 2002), and thus present an extended dietary record (Bearhop et al. 2004). However, stable isotopes are incorporated into different body tissues at different rates, which can affect interpretation of data (Martinez del Rio et al. 2009).

Our understanding of the dynamics of stable isotope values in elasmobranchs lags behind that of other taxa. For example, isotopic turnover rates in tissues of elasmobranchs have only been reported for two species ($\delta^{15}\text{N}$ in captive *Potamotrygon motoro*; MacNeil et al. 2006; $\delta^{15}\text{N}$

and $\delta^{13}\text{C}$ in captive *Carcharhinus plumbeus*; Logan and Lutcavage 2010), compared to numerous studies investigating isotopic turnover rates in mammals (e.g. MacAvoy et al. 2006, Miller et al. 2008), birds (e.g. Hobson and Clark 1992, Haramis et al. 2007), and bony fishes (e.g. Jardine et al. 2004, Perga and Gerdeaux 2005, McIntyre and Flecker 2006). In addition to understanding turnover rates, it is important to understand the variability of isotopic values for various tissue types within an individual in order to make full use of stable isotopic data and compare information among studies (e.g. Pinnegar and Polunin 1999, Vander Zanden and Rasmussen 2001, Sweeting et al. 2005).

The purpose of this study was to (1) compare the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of muscle, blood, and dorsal fin tissues from juvenile bull sharks (*Carcharhinus leucas*) and blood and dorsal fin tissues of large (juvenile and adult) tiger sharks (*Galeocerdo cuvier*) to determine if resulting intra-specific values from one tissue are comparable to those of other tissues for each species, and (2) gain insights into how differences among tissues within individuals may vary with shark size. Understanding if stable isotope analysis provides relatively consistent dietary data across tissue types, and if this consistency is similar across size-classes, may allow for less invasive sampling of tissues, and provide insight into ecological drivers of dietary variation.

Methods

Muscle, whole blood (“blood” hereafter), and dorsal fin (“fin”) tissues were collected from 81 juvenile bull sharks (70-162 cm total length) captured on 500m longlines within the Shark River estuary of Everglades National Park, Florida, USA (see Heithaus et al. 2009 for specific details of the study area and capture methods). We used a biopsy punch to collect a 0.5 cm³ muscle tissue biopsy *ca.* 5 cm lateral to the first dorsal fin, scissors to collect a 0.5 cm³ tissue

clip from the dorsal fin, and an 18 gauge needle to collect 2 ml of blood from the caudal vein. Tissues were placed on ice and frozen upon return to the laboratory. Skin was removed from muscle samples before laboratory preparations. All samples were dried and homogenized. Blood and fin clips were collected from 46 tiger sharks (159-396 cm TL) captured on drumlines during long-term studies in the hypersaline seagrass ecosystem of Shark Bay, Western Australia (see Wirsing et al. 2006 for study site and sampling details). Sample collection, storage, and processing protocols were identical to those for bull sharks.

All samples were analyzed at the Florida International University Stable Isotope Facility (43 *C. leucas* blood samples, 50 *C. leucas* muscle samples, and 26 *C. leucas* fin samples) or the Yale Earth System Center for Stable Isotopic Studies (34 *C. leucas* blood samples, 27 *C. leucas* muscle samples, 19 *C. leucas* fin samples, 46 *G. cuvier* blood samples, and 46 *G. cuvier* fin samples). Lipids were not extracted from any tissues, and C:N ratios indicated that corrections for lipid content were not necessary (Post et al. 2007). To verify analytical consistency, we randomly selected samples to be analyzed at both Florida International University and Yale University, for which the variation between resulting $\delta^{13}\text{C}$ $\delta^{15}\text{N}$ values were $0.13\text{‰} \pm 0.20\text{SE}$.

We used least squares regression analysis to determine (1) the relationships between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for all paired tissues of bull sharks (i.e. blood-muscle, blood-fin, muscle-fin) and tiger sharks (i.e. blood-fin), and (2) the relationship between shark length and paired differences between tissues. Each paired difference was calculated by taking the absolute difference between the $\delta^{13}\text{C}$ or $\delta^{15}\text{N}$ values of two tissue types for each shark (e.g. if muscle = -13.1‰ and blood = -13.8‰ , then the paired difference = 0.7‰). Cook's test was used to identify outliers, each tissue comparison regression model slope was tested to determine if it deviated significantly from a slope of one, and paired difference models were tested as linear and polynomial models to

identify the best fitting model. Because isotope assimilation into body tissues experiences a lag time based on the turnover rate of the specific tissue type (reviewed by Martinez del Rio et al. 2009), and sharks can experience ontogenetic shifts in diet (reviewed by Weatherbee and Cortes 2004), in some cases polynomial models may produce the best fit for determining the relationship between isotope values and shark size.

Results

Comparisons of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values revealed highly significant positive correlations for all tissue pairs in bull sharks. The slopes of all three bull shark $\delta^{13}\text{C}$ comparisons did not differ from 1:1 and all R^2 values were >0.71 (Fig. 1a, c, e). Blood was on average $0.57\text{‰} \pm 0.055$ SE more depleted (i.e. more negative) than muscle and on average $2.8\text{‰} \pm 0.10$ SE more depleted than fin, and muscle was on average $2.1\text{‰} \pm 0.092$ SE more depleted than fin (Fig. 1a, c, e). Relationships between $\delta^{15}\text{N}$ values were significant, but weaker than those of $\delta^{13}\text{C}$, with R^2 values between 0.15-0.43 (Fig. 1b, d, f). Only the relationship between muscle and fin deviated from a slope of one (slope = 0.6, $t_{41} = -7.8$, $p = <0.001$). Mean differences for bull shark blood and muscle $\delta^{15}\text{N}$ was $0.80\text{‰} \pm 0.064$ SE, blood and fin was $0.65\text{‰} \pm 0.16$ SE, and muscle and fin was $0.20\text{‰} \pm 0.15$ SE (Fig. 1b, d, f). The ranges of $\delta^{13}\text{C}$ values were relatively wide for all bull shark tissue types, while the ranges of $\delta^{15}\text{N}$ values were relatively narrow (Table 1).

Relationships between tissue types were similar in tiger sharks. Correlations for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ of blood and fin were positive and significant, but the relationship was tighter for $\delta^{13}\text{C}$ ($R^2 = 0.62$) than for $\delta^{15}\text{N}$ ($R^2 = 0.32$) (Fig. 1g, h). The slope for $\delta^{13}\text{C}$ was not significantly different from one, but the slope for $\delta^{15}\text{N}$ was (slope = 0.63, $t_{40} = -10.0$, $p = <0.001$). For tiger sharks, the $\delta^{13}\text{C}$ of blood was on average $1.2\text{‰} \pm 0.26$ SE more depleted than fin while the mean difference

in $\delta^{15}\text{N}$ was only $0.09\text{‰} \pm 0.21$ SE (Fig. 1g, h). Similar to the bull sharks, the ranges of $\delta^{13}\text{C}$ values were relatively wider than those of $\delta^{15}\text{N}$ values (Table 1). **[Insert Figure 1 and Table 1]**

Based on the tight relationships in isotopic values of tissues, it is not surprising most tissue types showed similar relationships between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ and shark total length. For both $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in bull sharks, all tissues declined until 110-130 cm TL, and then increased (Fig. 2a-f). All relationships between isotope values and shark total length were significant ($p < 0.05$) for bull sharks. For tiger sharks, $\delta^{13}\text{C}$ of fin and blood slightly increased with size until 250-300 cm TL, and then declined (Fig. 2g and i), while $\delta^{15}\text{N}$ declined with size until 250-300 cm TL, and then increased (Fig. 2h and j). Only the relationship between blood $\delta^{13}\text{C}$ values and tiger shark total length was significant. **[Insert Figure 2]**

The difference in $\delta^{13}\text{C}$ values between tissue types for bull sharks was influenced by shark total length for all pairings. In all cases for bull sharks, paired differences in $\delta^{13}\text{C}$ values were highest for the smallest individuals and decreased with size. This relationship was strongest for fin and blood ($R^2 = 0.64$), and weakest for fin and muscle ($R^2 = 0.21$; Fig. 3a, c, e). The paired difference between muscle and blood dropped rapidly until ~110cm TL, when the direction of the difference became less predictable. The difference between fin and blood dropped linearly and approached zero at approximately 165cm TL, and the difference between fin and muscle showed a relatively weak relationship with shark length. Paired differences for $\delta^{15}\text{N}$ of bull sharks showed a different pattern. There was no significant relationship between shark size and tissue difference in $\delta^{15}\text{N}$ of fin and muscle, while somewhat weak, but significant, nonlinear relationships were found for comparisons between blood and muscle ($R^2 = 0.18$), and blood and fin ($R^2 = 0.39$; Fig. 3b, d, f). The difference in $\delta^{15}\text{N}$ for these comparisons was

relatively low at small total lengths, increased slightly with size, and then declined in the largest individuals.

For tiger sharks, there was a significant but relatively weak ($R^2 = 0.27$), positive effect of shark size on differences in $\delta^{13}\text{C}$ of fin and blood, and shark size explained no variation in differences between $\delta^{15}\text{N}$ of fin and blood (Fig. 3g, h). **[Insert Figure 3]**

Discussion

Our study of two shark species at different life history stages, and from two different environments, has important implications for using stable isotope data in studies of elasmobranchs. Variability in stable isotope values within and among individuals can be driven by many ecological factors, including environmental conditions, metabolic processes, food quality, or changes in behavior, among many other factors (reviewed by Martinez del Rio et al. 2009). Yet, patterns of variability in stable isotope values among individuals can provide important insights into the trophic ecology of individuals within a population, as well as into differences among population and species.

Body size appears to be one factor that explained the regression slopes for some of the inter-tissue paired differences for our sample populations (Fig. 3). The paired differences in $\delta^{13}\text{C}$ of bull shark tissues were greatest in smaller individuals and decreased with size, indicating that isotopic values of different tissues were more similar for larger individuals. Prior to birth, bull sharks are directly connected to their mothers by an umbilical cord, which serves as a pathway through which nutrients and energy are transferred between mother and fetus. Based on the presence of open umbilical scars, bull sharks in the coastal Everglades are born between 65-75 cm TL. Because of their connection to their mothers, pups should have $\delta^{13}\text{C}$ values similar to

181 their mothers (coastal predators; $\delta^{13}\text{C}$ $\sim$ -15‰ in our study area; Chasar et al. 2005), as seen in
182 cetaceans (e.g. bottlenose dolphins, *Tursiops truncatus*, Knoff et al. 2008; sea lions, *Zalophus*
183 *californianus*, Porras-Peters et al. 2008). After birth, juvenile sharks spend several years in low-
184 salinity estuaries and nearshore waters (e.g. Wiley and Simpfendorfer 2007, Heithaus et al.
185 2009), and therefore $\delta^{13}\text{C}$ values should begin to diverge from their mothers as they adopt a more
186 $\delta^{13}\text{C}$ -depleted estuarine diet (consumer taxa $\delta^{13}\text{C}$ is typically $< -25\text{‰}$ in the Shark River;
187 Williams and Trexler 2006, M. Heithaus *unpublished data*; see also Fig 2). The change in $\delta^{13}\text{C}$
188 values should occur earlier in tissues that turnover more rapidly. For example, differences
189 between blood and both fin and muscle in the smallest bull sharks suggests that fin tissue largely
190 maintains the maternal signature, likely due to a slower turnover rate. In contrast, blood reflects
191 the young sharks' diet within two years of birth, likely due to a faster turnover rate in this tissue
192 type (MacNeil et al. 2006).

193 The regression model for the paired difference of $\delta^{13}\text{C}$ for muscle and blood appears to
194 reach equilibrium around 110 cm TL and two years of age (based on growth rates in Branstetter
195 and Stiles 1987 and estimated sizes at birth; Heithaus et al. 2009). This may indicate the time
196 period for which muscle $\delta^{13}\text{C}$ values are no longer influenced by the maternal diet for juveniles,
197 and accurately portray that individual's diet over its lifetime. Deviations in isotope values of
198 larger individuals may reflect other underlying ecological patterns, for example seasonal shifts in
199 diet, which may be displayed more rapidly in blood values than in muscle or fin (P. Matich et al.
200 *unpublished data*). In contrast to bull sharks, differences in $\delta^{13}\text{C}$ among blood and fin clips
201 increased with size in tiger sharks. This likely reflects a difference in the feeding ecology of the
202 two species, and the increasing difference in $\delta^{13}\text{C}$ of blood and fin may reflect a shift in the diets
203 of tiger sharks as they grow (e.g. Lowe et al. 1996, Simpfendorfer et al. 2001).

Size-based differences among tissues in stable isotope values are important to consider when investigating the ecological drivers of dietary variation within populations. $\delta^{13}\text{C}$ values (Fig. 2a, c, e) support the hypothesis that the maternal influence on isotopic values of juvenile bull sharks is evident for several years, but individual variability in isotopic values makes it difficult to draw conclusions about the precise timing of tissue values equilibrating. Especially for $\delta^{13}\text{C}$ of both species, the range of isotope values was relatively wide, even for sharks of a given size, suggesting that other factors, like habitat use (e.g. Darimont et al. 2009, Quevedo et al. 2009), body condition (e.g. Tinker et al. 2008, Tucker et al. 2009), and/or seasonal shifts (e.g. Inger et al. 2006, Cherel et al. 2007) may affect the diet patterns for individuals of these two populations.

The strong positive correlations between tissues in $\delta^{13}\text{C}$ for both bull sharks and tiger sharks (Fig. 1) suggest that for a species, multiple tissues may be compared after applying a correction factor. A strict 1:1 substitution of values among tissues is not recommended, and we suggest correction factors should be generated for individual populations because ecological differences may lead to variability in isotopic assimilation across individuals of the same taxa (Post 2002). Using correction factors generated for a species in one ecosystem may differ from those generated for the same species collected from a different ecosystem, and therefore it is currently most appropriate to generate correction factors on a per-population basis.

Tissue comparisons may allow for gaps within data sets to be filled and to increase the number of individuals that can be directly compared. Individuals for which isotope values of a particular tissue are not available may have correction factors applied to estimate isotopic value(s) of the uncollected tissue. Yet, it is important to consider potential factors that limit the use of correction factors. Species that experience ontogenetic shifts in diet may experience

variability in inter-tissue relationships between isotope values (e.g. Quillfeldt et al. 2008, Tierney et al. 2008, Young et al. 2010), and therefore correction factors may be more accurate for certain age/size-classes of animals. For example, the difference between tissues for bull sharks (paired differences; Fig. 3) were largest (7‰ fin-blood) for the smallest individuals sampled, and tended to decrease and approach equilibrium (1:1 relationship) as bull shark total length increased. This suggests that correction factors may be more useful for larger individuals, which generally had smaller differences in isotope values for different tissues. Therefore, care must be taken when using correction factors and variability in factors that affect trophic role (such as body size) must be taken into consideration prior to using estimated isotope values produced by correction factors for diet analysis.

Relationships among tissues in $\delta^{15}\text{N}$ were relatively weak, raising doubts as to whether tissues can be compared reliably. The relatively small range in $\delta^{15}\text{N}$ for both species (3.3‰ and 3.4‰ for tiger sharks and bull sharks, respectively), however, could be responsible for these patterns, and the question of interest may determine the magnitude of potential error when substituting $\delta^{15}\text{N}$ values for different tissue types when using correction factors. The paired differences in $\delta^{15}\text{N}$ for bull sharks ($R^2 = 0.04$ to 0.39) and tiger sharks ($R^2 < 0.01$) were relatively weak, suggesting that combining data sets with multiple tissue types may be problematic for $\delta^{15}\text{N}$. Because we found the $\delta^{15}\text{N}$ relationships to be relatively weak, we suggest that further ecological and physiological studies are needed to elucidate the factor(s) affecting inter-tissue differences in $\delta^{15}\text{N}$.

Published turnover rates for elasmobranch tissues (MacNeil et al. 2006), combined with the long duration before convergence of $\delta^{13}\text{C}$ values of blood and muscle of bull sharks in our study, suggest that using stable isotopes from these tissues are most appropriate for elucidating

long-term dietary patterns. Such long-term information may be useful for investigating questions such as the degree of specialization within populations, how changes in environmental factors may influence consumer diets, and what ecological factors influence inter-population variation in feeding behaviors. Other taxa exhibit considerably faster turnover rates for blood (e.g. ~52 days ($\delta^{13}\text{C}$) and ~46 days ($\delta^{15}\text{N}$) for mice (*Mus musculus*) MacAvoy et al. 2006), muscle (e.g. 4-5 months ($\delta^{15}\text{N}$) for whitefish (*Coregonus lavaretus*) Perga and Gerdeaux 2005), and fin (e.g. ~37 days ($\delta^{15}\text{N}$) for armored catfish (*Ancistrus triradiatus*) McIntyre and Flecker 2006) tissues, allowing for more fine-scale diet studies. Therefore, stomach content analysis remains an important complimentary method for studying elasmobranch trophic ecology, especially when investigating short-term variability in diets.

Our understanding and application of stable isotopes in elasmobranchs is still in its infancy. Sharks and rays are important top and mesopredators in multiple ecosystems (Heithaus et al. 2010). With many populations jeopardized worldwide, stable isotope analysis provides an important tool for studying their trophic ecology non-lethally. Yet, further studies in the field and laboratory, and across a variety of taxa, environments, and life history stages, are needed to better understand how stable isotopes can be best applied and interpreted for studies of their trophic ecology.

Acknowledgements

Funding for this project was provided by the National Science Foundation (DBI0620409, DEB9910514, OCE0526065, OCE0746164) and Florida International University's Marine Sciences Program. We thank the many volunteers who assisted with shark fishing and processing stable isotope samples, especially Derek Burkholder, Richard Chang, Bryan Delius,

Meagan Dunphy-Daly, Kirk Gastrich, and Aaron Wirsing. Thanks also to Joel Trexler and the FCE LTER for providing funding and logistical support for this project. Research was conducted under Everglades National Park permits EVER-2009-SCI-0024, EVER-2007-SCI-0025, and EVER-2005-SCI-0030, authorizations from the Department of Environment and Conservation, Western Australia and Fisheries WA and with Florida International University IACUC approval. This is publication 41 of the Shark Bay Ecosystem Research Project.

References

- Bearhop, S., Adams, C.E., Waldron, S., Fuller, R.A., and MacLeod, H. 2004. Determining trophic niche width: a novel approach using stable isotope analysis. *J. Anim. Ecol.* **73**: 1007-1012.
- Branstetter, S. and Stiles, R. 1987. Age and growth estimates of the bull shark, *Carcharhinus leucas*, from the northern Gulf of Mexico. *Environ. Biol. Fishes* **20**: 169-181.
- Chasar, L.C., Chanton, J.P., Koenig, C.C., and Coleman, F.C. 2005. Evaluating the effect of environmental disturbance on the trophic structure of Florida Bay, U.S.A.: Multiple stable isotope analyses of contemporary and historical specimens. *Limnol. Oceanogr.* **50**: 1059-1072.
- Cherel, Y., Hobson, K.A., Guinet, C., and Vanpe, C. 2007. Stable isotopes document seasonal changes in trophic niches and winter foraging individual specialization in diving predators from the Southern Ocean. *J. Anim. Ecol.* **76**: 826-836.
- Darimont, C.T., Paquet, P.C., and Reimchen, T.E. 2009. Landscape heterogeneity and marine subsidy generate extensive intrapopulation niche diversity in a large terrestrial vertebrate. *J. Anim. Ecol.* **78**: 126-133.

296 Domi, N., Bouquegneau, J.M., and Das, K. 2005. Feeding ecology of five commercial shark
 297 species of the Celtic Sea through stable isotope and trace metal analysis. Mar. Environ.
 298 Res. **60**: 551-569.

299 Dulvy, N.K., Baum, J.K., Clarke, S., Compagno, L.J.V., Cortes, E., Domingo, A., Fordham, S.,
 300 Fowler, S., Francis, M.P., Gibson, C., Martinez, J., Musick, J.A., Soldo, A., Stevens, J.D.,
 301 and Valenti, S. 2008 You can swim but you can't hide: the global status and conservation
 302 of oceanic pelagic sharks and rays. Aquat. Conserv.: Mar. Freshw. Ecosyst. **18**: 459-482.

303 Fisk, A.T., Tittlemier, S.A., Pranschke, J.L., and Norstrom, R.J. 2002. Using anthropogenic
 304 contaminants and stable isotopes to assess the feeding ecology of Greenland sharks.
 305 Ecology **83**: 2162-2172.

306 Haramis, G.M., Link, W.A., Osenton, P.C., Carter, D.B., Weber, R.G., Clark, N.A., Teece, M.A.,
 307 and Mizrahi, D.S. 2007. Stable isotope and pen feeding trial studies confirm the value of
 308 horseshoe crab *Limulus polyphemus* eggs to spring migrant shorebirds in Delaware Bay.
 309 J. Avian Biol. **38**: 367-376.

310 Heithaus, M.R, Frid, A., Wirsing, A.J., and Worm, B. 2008. Predicting ecological consequences
 311 of marine top predator declines. Trends Ecol. Evol. **23**: 202-210.

312 Heithaus, M.R., Delius, B.K., Wirsing, A.J., and Dunphy-Daly, M.M. 2009. Physical factors
 313 influencing the distribution of a top predator in a subtropical oligotrophic estuary.
 314 Limnol. Oceanogr. **54**: 472-482.

315 Heithaus M.R., Frid A., Vaudo J.J., Worm B., and Wirsing A.J. 2010. Unraveling the ecological
 316 importance of elasmobranchs. Pp 608-633 In: Carrier, J.C., Musick, J.A., and Heithaus,
 317 M.R. (eds) Sharks and Their Relatives II: Biodiversity, Adaptive Physiology, and
 318 Conservation. CRC Press, Boca Raton, FL.

319 Hobson, K.A. and Clark, R.G. 1992. Assessing avian diets using stable isotopes I: turnover of
 320 ^{13}C in tissues. *Condor* **94**: 181-188.

321 Inger, R., Ruxton, G.D., Newton, J., Colhoun, K., Robinson, J.A., Jackson, A.L., and Bearhop, S.
 322 2006. Temporal and intrapopulation variation in prey choice of wintering geese
 323 determined by stable isotope analysis. *J. Anim. Ecol.* **75**: 1190-1200.

324 Jardine, T.D., MacLatchy, D.L., Fairchild, W.L., Cunjak, R.A., and Brown, S.B. 2004. Rapid
 325 carbon turnover during growth of Atlantic salmon (*Salmo salar*) smolts in sea water, and
 326 evidence for reduced food consumption by growth-stunts. *Hydrobiologia* **527**: 63-75.

327 Knoff, A., Hohn, A., and Macko, S. 2008. Ontogenetic diet changes in bottlenose dolphins
 328 (*Tursiops truncatus*) reflected through stable isotopes. *Mar. Mamm. Sci.* **24**: 128-137.

329 Logan, J.M. and Lutcavage, M.E. Stable isotope dynamics in elasmobranch fishes. *Hydrobiologia*
 330 DOI 10.1007/s10750-010-0120-3.

331 Lowe, C.G., Wetherbee, B.M., Crow, G.L., and Tester, A.L. 1996. Ontogenetic dietary shifts and
 332 feeding behavior of the tiger shark, *Galeocerdo cuvier*, in Hawaiian waters. *Environ.*
 333 *Biol. Fishes* **47**: 203-211.

334 MacAvoy, S.E., Arneson, L.S., and Bassett, E. 2006. Correlation of metabolism with tissue
 335 carbon and nitrogen turnover rate in small mammals. *Oecologia* **150**: 190-201.

336 MacNeil, M.A., Skomal, G.B., and Fisk, A.T. 2005. Stable isotopes from multiple tissues reveal
 337 diet switching in sharks. *Mar. Ecol. Prog. Ser.* **302**: 199-206.

338 MacNeil, M.A., Drouillard, K.G., and Fisk, A.T. 2006. Variable uptake and elimination of stable
 339 nitrogen isotopes between tissues in fish. *Can. J. Fish. Aquat. Sci.* **63**: 345-353.

340 Martinez del Rio, C., Wolf, N., Carleton, S.A., and Gannes, L.Z. 2009. Isotopic ecology ten
 341 years after a call for more laboratory experiments. *Biol. Rev.* **84**: 91-111.

342 McIntyre, P.B. and Flecker, A.S. 2006. Rapid turnover of tissue nitrogen of primary consumers
 343 in tropical freshwaters. *Oecologia* **148**: 12-21.

344 Miller, J.F., Millar, J.S., and Longstaffe, F.J. 2008. Carbon- and nitrogen-isotope tissue-diet
 345 discrimination and turnover rates in deer mice, *Peromyscus maniculatus*. *Can. J. Zoo.* **86**:
 346 685-691.

347 Perga, M.E. and Gerdeaux, D. 2005. 'Are fish what they eat' all year round? *Oecologia* **144**:
 348 598-606.

349 Pinnegar, J.K. and Polunin, N.V.C. 1999. Differential fraction of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ among fish
 350 tissues: implications for the study of trophic interactions. *Funct. Ecol.* **13**: 225-231.

351 Porras-Peters, H., Aurióles-Gamboa, D., Cruz-Escalona, V.H., and Koch, P.L. 2008. Trophic
 352 level and overlap of sea lions (*Zalophus californianus*) in the Gulf of California, Mexico.
 353 *Mar. Mamm. Sci.* **24**: 554-576.

354 Post, D.M. 2002. Using stable isotopes to estimate trophic position: models, methods, and
 355 assumptions. *Ecology* **83**: 703-718.

356 Post, D.M., Layman, C.A., Arrington, D.A., Takimoto, G., Montaña, C.G., and Quattrochi, J.
 357 2007. Getting to the fat of the matter: models, methods and assumptions for dealing with
 358 lipids in stable isotope analyses. *Oecologia* **152**: 179-189.

359 Quevedo, M., Svanback, R., and Eklov, P. 2009. Intrapopulation niche partitioning in a
 360 generalist predator limits food web connectivity. *Ecology* **90**: 2263-2274.

361 Quillfeldt, P., Bugoni, L., McGill, R.A.R., Masello, J.F., and Furness, R.W. 2008. Differences in
 362 stable isotopes in blood and feathers of seabirds are consistent across species, age, and
 363 latitude: implications for food web studies. *Mar. Biol.* **155**: 593-598.

364 Simpfendorfer, C.A., Goodreid, A.B., and McAuley, R.B. 2001. Size, sex, and geographic
 365 variation in the diet of tiger sharks, *Galeocerdo cuvier*, from Western Australian waters.
 366 Environ. Biol. Fishes **61**: 37-46.

367 Sweeting, C.J., Jennings, S., and Polunin, N.V.C. 2005. Variance in isotopic signatures as a
 368 descriptor of tissue turnover and degree of omnivory. Funct. Ecol. **19**: 777-784.

369 Tierney, M., Southwell, C., Emmerson, L.M., and Hindell, M.A. 2008. Evaluating and using
 370 stable-isotope analysis to infer diet composition and foraging ecology of Adelie penguins
 371 *Pygoscelis adeliae*. Mar. Ecol. Prog. Ser. **355**: 297-307.

372 Tinker, M.T., Bentall, G., and Estes, J.A. 2008. Food limitation leads to behavioral
 373 diversification and dietary specialization in sea otters. PNAS **105**: 560-565.

374 Tucker, S., Bowen, W.D., Iverson, S.J., Blanchard, W., and Stenson, G.B. 2009. Sources of
 375 variation in the diets of harp and hooded seals estimated from quantitative fatty acid
 376 signature analysis (QFASA). Mar. Ecol. Prog. Ser. **384**: 287-302.

377 Vander Zanden, M. J. and Rasmussen, J.B. 2001. Variation in $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ trophic
 378 fractionation: implications for aquatic food web studies. Limnol. Oceanogr. **48**:2061-
 379 2066.

380 Weatherbee, B.M. and Cortes, E. 2004. Food consumption and feeding habits. *In*: Carrier, JC,
 381 JA Musick and MR Heithaus (eds) Biology of sharks and their relatives. Boca Raton,
 382 FL. CRC Press, pp 225-246.

383 Wiley, T.R. and Simpfendorfer, C.A. 2007. The ecology of elasmobranches occurring in the
 384 Everglades National Park, Florida: implications for conservation and management. Bull.
 385 Mar. Sci. **80**: 171-189.

386 Williams, A.J. and Trexler, J.C. 2006. A preliminary analysis of the correlation of food-web
387 characteristics with hydrology and nutrient gradients in the southern Everglades.
388 Hydrobiologia **569**:493–504.

389 Wirsing, A.J., Heithaus, M.R., and Dill, L.M. 2006. Tiger shark (*Galeocerdo cuvier*) abundance
390 and growth rates in a subtropical embayment: evidence from seven years of standardized
391 fishing efforts. Mar. Biol. **4**: 961-968.

392 Young, B.G., Loseto, L.L., and Ferguson, S.H. 2010. Diet differences among age classes of
393 Arctic seals: evidence from stable isotope and mercury biomarkers. Pol. Biol. **33**: 153-
394 162.

Table and Figure Legends

Table 1: Minimum and maximum values for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for blood, muscle, and fin for *Carcharhinus leucas* and blood and fin for *Galeocerdo cuvier* in ‰.

Figure 1: Comparisons of $\delta^{13}\text{C}$ for blood and fin (a), muscle and fin (c), and blood and muscle (e), and comparisons of $\delta^{15}\text{N}$ for blood and fin (b), muscle and fin (d), and blood and muscle (f) for *Carcharhinus leucas*, and $\delta^{13}\text{C}$ for blood and fin (g), and $\delta^{15}\text{N}$ for blood and fin (h) for *Galeocerdo cuvier*.

Figure 2: Comparisons of $\delta^{13}\text{C}$ and shark total length for fin (a), blood (c), and muscle (e), and comparisons of $\delta^{15}\text{N}$ and shark total length for fin (b), blood (d), and muscle (f) for *Carcharhinus leucas*, and $\delta^{13}\text{C}$ and shark total length for fin (g) and blood (i), and $\delta^{15}\text{N}$ and shark total length for fin (h) and blood (j) for *Galeocerdo cuvier*.

Figure 3: Paired differences of $\delta^{13}\text{C}$ for blood and fin (a), muscle and fin (c), and blood and muscle (e), and of $\delta^{15}\text{N}$ for blood and fin (b), muscle and fin (d), and blood and muscle (f) for *Carcharhinus leucas*, and $\delta^{13}\text{C}$ for blood and fin (g), and $\delta^{15}\text{N}$ for blood and fin (h) for *Galeocerdo cuvier*.

Table 1: Ranges of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in bull sharks and tiger sharks in ‰.

		Min $\delta^{13}\text{C}$	Max $\delta^{13}\text{C}$	Min $\delta^{15}\text{N}$	Max $\delta^{15}\text{N}$
Bull Sharks	Blood	-26.86	-16.27	9.91	12.53
	Muscle	-26.79	-16.51	11.07	13.26
	Fin	-24.62	-15.13	10.81	13.00
Tiger Sharks	Blood	-15.72	-9.56	10.57	13.09
	Fin	-14.69	-8.77	10.41	13.03

Figure 1

Figure 2

Figure 3