

2006

Flux of organic carbon in a riverine mangrove wetland in the Florida Coastal Everglades

Melissa M. Romigh

Department of Wildlife & Fisheries Sciences, Texas A&M University

Stephen E. Davis III

Department of Wildlife & Fisheries Sciences, Texas A&M University

Victor H. Rivera-Monroy

Department of Oceanography & Coastal Science, Wetland Biogeochemistry Institute, Louisiana State University

Robert R. Twilley

Department of Oceanography & Coastal Science, Wetland Biogeochemistry Institute, Louisiana State University

Follow this and additional works at: https://digitalcommons.fiu.edu/fce_lter_journal_articles

 Part of the [Hydrology Commons](#)

Recommended Citation

Romigh, M.M., S.E. Davis, V.H. Rivera-Monroy, R.R. Twilley. 2006. Flux of organic carbon in a riverine mangrove wetland in the Florida Coastal Everglades. *Hydrobiologia* 569(1): 505-516.

This material is based upon work supported by the National Science Foundation through the Florida Coastal Everglades Long-Term Ecological Research program under Cooperative Agreements #DBI-0620409 and #DEB-9910514. Any opinions, findings, conclusions, or recommendations expressed in the material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

This work is brought to you for free and open access by the FCE LTER at FIU Digital Commons. It has been accepted for inclusion in FCE LTER Journal Articles by an authorized administrator of FIU Digital Commons. For more information, please contact dcc@fiu.edu, jkrefft@fiu.edu.

Flux of Organic Carbon in a Riverine Mangrove Wetland in the Florida Coastal

2 Everglades

4 Melissa M. Romigh¹, Stephen E. Davis, III^{1a}, Victor H. Rivera-Monroy², and Robert R. Twilley²

6 ¹Department of Wildlife & Fisheries Sciences, 2258 TAMU, Texas A&M University, College
Station, TX 77843 USA

8

²Wetland Biogeochemistry Institute, Department of Oceanography & Coastal Science, Louisiana
10 State University, Baton Rouge, LA 70803 USA

12 *Keywords: DOC, mangrove forest, wetland-water column fluxes, Everglades National Park*

14 Note: This paper has not been submitted elsewhere in identical or similar form, nor will it be
during the first three months after its submission to *Hydrobiologia*.

16

18

20

22

^a Corresponding Author

24

ABSTRACT

2 Short term (daily) and seasonal variations in concentration and flux of dissolved organic carbon
(DOC) were examined over 15 tidal cycles in a riverine mangrove wetland along Shark River,
4 Florida in 2003. Due to the influence of seasonal rainfall and wind patterns on Shark River's
hydrology, samplings were made to include wet, dry and transitional (Norte) seasons. We used a
6 flume extending from a tidal creek to a basin forest to measure vertical (vegetated soil/water
column) and horizontal (mangrove forest/tidal creek) flux of DOC. We found significant
8 ($p < 0.05$) variations in surface water temperature, salinity, conductivity, pH and mean
concentration of DOC with season. Water temperature and salinity followed seasonal patterns of
10 air temperature and rainfall, while mean DOC concentration was highest during the dry season
(May), followed by the wet (October) and 'Norte' (December) seasons. This pattern of DOC
12 concentration may be due to a combination of litter production and inundation pattern of the
wetland. In contrast to daily (between tides) variation in DOC flux between the mangrove forest
14 and tidal creek, daily variations of mean water quality were not significant. However, within-
tide variation of DOC flux, dissolved oxygen content and salinity was observed. This indicated
16 that the length of inundation and water source (freshwater vs. saltwater) variation across tidal
cycles influenced water quality and DOC flux in the water column. Net DOC export was
18 measured in October and December, suggesting the mangrove forest was a source of DOC to the
adjacent tidal creek during these periods. Net annual export of DOC from the fringe mangrove
20 to both the tidal creek and basin mangrove forest was $56 \text{ g C m}^{-2} \text{ y}^{-1}$. The seasonal pattern in our
flux results indicates that DOC flux from this mangrove forest may be governed by both
22 freshwater discharge and tidal range.

INTRODUCTION

2 Coastal mangrove wetlands are valued because they provide food, shelter and nursery
habitat for a wide assortment of aquatic and terrestrial species (Robertson & Blaber, 1992;
4 Hogarth, 1999). Mangroves are also valued for their contributions to detritus-based food webs in
adjacent estuarine and coastal waters via the export of organic carbon (OC; Odum & Heald,
6 1975; Twilley, 1988). Litter, mainly leaves and twigs, produced in the mangrove canopy
represents a significant source of the organic carbon and nutrients available for export to adjacent
8 coastal waters (Robertson et al., 1992). Leaching of this litter is a rapid component of the
decomposition process and provides an important source of dissolved organic material to the soil
10 and water column (Davis et al., 2003; Romero et al., 2005). Given that this organic material is
highly labile, a sizable fraction of mangrove detritus is rapidly utilized by heterotrophic
12 microorganisms and made available to higher consumers in the detritus food web (Benner et al.,
1986).

14 Most mangrove wetlands show a net export of organic carbon, and DOC can account for
more than 80% of the organic carbon exported (Twilley, 1985; Lee, 1995; Furukawa et al., 1997;
16 Machiwa & Hallberg, 2002). For specific mangrove forest types, it has been shown that much of
the variation in export can be attributed to tidal amplitude, inundation frequency, and fauna
18 among other factors (Twilley, 1985; Robertson, 1986; Lee, 1995; Dittmar & Lara, 2001b; Feller,
2002). And these factors can vary seasonally, causing distinct patterns of organic carbon
20 dynamics in estuarine mangrove wetlands (Boto & Wellington, 1988; Dittmar & Lara, 2001a;
Dittmar et al., 2001; Davis et al., 2001a; Sutula et al., 2003). However, we still lack an
22 understanding of how small scale forces such as diurnal tides and seasonal variations in upland

runoff—as indicated by freshwater inflows—control organic carbon dynamics in these
2 ecosystems.

Few studies have quantified mangrove-water column interactions in situ, and there have
4 been even fewer such studies conducted in mangrove forests of Everglades National Park (ENP;
Davis et al., 2001a; Davis et al., 2001b; Sutula et al., 2003). This coastal region is of particular
6 interest where landscape-scale hydrologic restoration efforts are underway. The focus of our
study was on the short term (daily) and seasonal variability in organic carbon dynamics
8 (concentration and fluxes) between a riverine mangrove wetland and the Shark River estuary
(FL, USA). We hypothesized that seasonal variations in freshwater flow originating from the
10 Everglades and local diurnal tides were important in regulating organic carbon exchange in this
wetland. We further hypothesized that the high productivity common to riverine mangrove
12 forests and the high tidal range in Shark River (relative to other estuaries in the Gulf of Mexico)
would result in a net export of organic carbon for the year.

14 **METHODS**

16 *Site Description*

18 Everglades National Park is located in southern Florida and comprises 610,483 ha, of
which 144,447 ha are mangrove forest (Lewis et al., 1985; Welch et al., 1999). Average annual
20 rainfall in the region is 138 cm, with distinct wet (approximately 60% average annual rainfall
from June to September) and dry (approximately 25% average annual rainfall from November to
22 April) seasons (Figure 1). Shark River is a mangrove-dominated tidal river along the southwest
coast of ENP and is one of the largest estuaries in South Florida. Discharge of freshwater from
24 Shark River Slough to the estuary follows patterns of intra-annual (i.e., seasonal) and inter-

annual rainfall (Chen & Twilley, 1999; Childers et al., 2005). Tides in Shark River are
2 predominantly semi-diurnal with mean amplitude of 0.5–1.0 m.

Our site was located approximately 4 km upstream from the mouth of the estuary at SRS-
4 6 (25°21.852 N, 81°04.667 W), a site routinely sampled as part of the Florida Coastal Everglades
(FCE) LTER program. Specifically, our sampling site was located along a small tidal creek that
6 connects the mangrove wetland at SRS-6 to Shark River (Figure 2). The width of the tidal creek
is approximately 2 m and the distance from the sampling sites to Shark River was approximately
8 30 m. The riverine mangrove forest at this location is tidally inundated twice per day (Figure 3)
and is dominated by three species of mangroves (*Rhizophora mangle*, *Avicennia germinans*, and
10 *Laguncularia racemosa*).

12 *Field Sampling*

Samplings were made in 2003 in three seasons: dry (May), wet (October) and the
14 transitional ‘Norte’ season (December) to understand the importance of season in driving organic
carbon dynamics. Also, we sampled across multiple, consecutive tidal cycles (4-6 tides per
16 season) to capture the range of variability in concentration and fluxes within each season.

We used a flow-through flume to measure organic carbon fluxes between the wetland
18 surface and the water column (vertical) and exchanges between the wetland and Shark River
(horizontal). The experimental flume was open to surface water flow at both ends and measured
20 approximately 2 m wide and 12.5 m in length, extending from the tidal creek into the mangrove
forest. The flume walls were positioned perpendicular to the tidal creek and parallel to the
22 direction of flow of the flooding tidal water, so as to mimic natural flooding patterns in the
enclosed area of wetland. Panels on the flumes, consisting of clear, corrugated fiberglass, were

removed after each sampling to prevent long-term panel effects such as shading, edge scouring,
2 and detritus accumulation (Childers & Day, 1988).

During each sampling period, an ISCO auto-sampler was placed at each end of the flume
4 and programmed to collect a 1-liter water sample every 30 minutes from a single point inside the
flume. Samples were collected simultaneously from each end of the flume during flood and ebb
6 half tides. Samples were retrieved every 12 hours, packed in ice and transported to the
laboratory for processing. In addition, we continuously monitored temperature, salinity, pH,
8 conductivity and dissolved oxygen (DO) concentration of tidal creek surface water with a pre-
calibrated mini-sonde. We also utilized data on water level and salinity collected continuously at
10 SRS-6 as part of the FCE-LTER program. Lastly, the microtopography of the wetland within the
flume was surveyed in August 2003 to build a detailed map of elevation. These elevation data
12 were used in conjunction with water level data to estimate the volume of water in the flume
during each sampling and the change in volume (volume flux) between samplings.

Laboratory Analyses

16 Water samples were filtered through pre-rinsed (triplicate 5 ml rinses with de-ionized
water), pre-combusted (4 hr at 500°C) and pre-weighed 47 mm Whatman GF/F papers.
18 Filtration was performed within 24 hr of collection and both filtered and unfiltered samples were
refrigerated at 4 °C in 125 ml bottles. Particulate matter trapped on the filter was dried at 70 °C
20 for 24 hr and weighed for calculation of total suspended sediment (TSS) concentration, then
combusted (500 °C for 4 hr) to determine losses on ignition.

22 DOC and TOC concentrations in filtered and unfiltered samples were quantified using
high temperature catalytic oxidation (HTCO) and infrared detection with a Shimadzu TOC-5000.

Organic carbon concentrations were determined against potassium hydrogen phthalate standards.

2 Samples were measured in triplicate with a fixed coefficient of variation of 2%. Duplicate
samples were analyzed periodically to check for reproducibility of results and to evaluate the
4 precision of measurements.

6 *Statistical Analyses and Flux Calculations*

Tidal (flood, ebb) and seasonal (dry, wet, ‘Norte’) differences in temperature, salinity,
8 conductivity, pH, dissolved oxygen, [TSS] and [DOC] were analyzed using a two-way ANOVA
to test for interaction effects of the two factors. In the case of unequal variances during the two-
10 way ANOVA analysis, separate one-way ANOVA and LSD or Dunnett’s T3 post-hoc tests
($p < 0.05$) were used to analyze differences between seasons and simple one-way ANOVA tests
12 ($p < 0.05$) were used to analyze differences between flood and ebb tides.

We used formulas for ‘instantaneous flux’ and ‘net flux’ described in Childers & Day
14 (1988) for this study. Net fluxes were obtained by adding flood and ebb instantaneous fluxes for
each tidal cycle. We also employed an equation developed by Rivera-Monroy et al. (1995) for
16 calculating net areal fluxes in a tidally flooded mangrove forest in Mexico (Equation 1).

$$\text{Net areal flux (g m}^{-2}\text{ h}^{-1}\text{)} = \frac{\text{total flux}_{\text{upstream}} - \text{total flux}_{\text{downstream}}}{\text{Flume area} \times \text{total time}} \quad (1)$$

Using a hypsometric approach, we inferred fluxes of water from changes in water level
18 collected at 30-minute intervals during several whole-tides. Direct measurements of water level
within the flume over the course of one complete tidal inundation were regressed with the
20 continuous water level recorder data supplied by the FCE-LTER station along Shark River (SRS-
6). The relationship generated was used to predict water level within the flume for all additional

sampling periods. To best relate water level in a system with asymmetrical tides, we developed
2 separate correlation curves for flood and ebb tides (adjusted R^2 values of 0.999 and 0.997 for
ebb- and flood-tide correlations, respectively).

4 Paired t-tests between mean upstream and downstream organic carbon concentrations
over each half tide were used to determine if fluxes of organic carbon within the flume were
6 significant ($p < 0.05$). We used instantaneous flux measurements from individually paired organic
carbon concentrations to calculate total flux values independently for each half-tide. No
8 difference in concentration between upstream and downstream pairs was interpreted as no net
flux over a given half-tide (net area flux = $0 \text{ mg C m}^{-2}\text{h}^{-1}$). For example, higher concentrations
10 during flood tide at the streamside end (upstream) indicated a net uptake of organic carbon by the
wetland, while lower concentrations at the streamside end indicated a net release (Rivera-
12 Monroy et al., 1995). We calculated net areal fluxes only when water was present at both ends
of the flume and a significant difference in mean [DOC] was observed between paired upstream
14 and downstream sample pairs.

16 **RESULTS**

The two-way ANOVA tests for half tide or seasonal differences in temperature, salinity,
18 conductivity, pH, dissolved oxygen and [DOC] showed an interaction effect of half-tide and
season, though all data exhibited unequal variance. One-way ANOVA tests indicated none of
20 the variables, except pH, varied significantly between flood and ebb tides. The water column pH
had a small range; however, and was circum-neutral (7.3–7.6) throughout the study period
22 (Romigh, 2005).

We observed a trend of increasing salinity during flood tide and decreased salinity during ebb tide for all tides sampled, reflecting the estuarine nature of this site. The concentration and % saturation of DO in the water column was highest in May and lowest during October. DO saturation (%) also decreased with increasing duration of inundation, indicating oxygen depletion of the water column due to respiration.

Water temperature in the flume ranged from 17.9 to 30.6 °C and varied seasonally, typical of the regional climate (Figure 4a). Salinity in the flume ranged overall from 6.4 to 29.8 ‰. Lowest salinities were observed in the wet season when freshwater input from the Everglades was greatest (October), and highest salinities were recorded in the dry season (May; Figure 4b). Concentrations of both TSS and DOC were significantly different across samplings (Figures 4c,d). Overall, TSS concentrations varied between 1–192 mg l⁻¹. Significantly higher TSS concentrations occurred in May (mean = 112 mg l⁻¹), while concentrations were low in both October (mean = 22 mg l⁻¹) and December (mean = 24 mg l⁻¹; Figure 4d). DOC concentrations were significantly different between all seasons and ranged from 1.7 to 17.9 mg l⁻¹. Overall, [DOC] was highest in May and lowest in October (Figure 4c).

Significant changes in DOC concentration within the flume were observed for 50% of the tides sampled. More than 95% of the total organic carbon in the samples we collected was in the dissolved fraction. Given that the particulate fraction was so small and our sampling approach did not account for the macro-particulate fraction (leaves, seedlings, etc.), we only considered the dissolved organic fraction in the calculation of fluxes.

Fluxes

We were able to calculate net areal fluxes of DOC for 12 of the 15 tidal cycles measured in this study. Of those, export from the mangrove wetland was the majority of significant fluxes (Figure 5). Half-tide vertical DOC fluxes ranged from -5.52 to 8.9901 g C m⁻² h⁻¹ in May, -14.4591 to 1.3664 g C m⁻² h⁻¹ in October and -2.4017 to 3.7240 g C m⁻² h⁻¹ in December. Overall, total exchanges of DOC were greater at the creek-side end of the flume compared with the forest side.

Most fluxes calculated in the dry season indicated a net import of DOC to the mangrove wetland, with much of that occurring at the tidal creek end of the flume (Figure 5). The highest significant import of DOC to our mangrove site occurred during May sampling at 0.3227 g C m⁻² h⁻¹. The tidal creek was the source of this material (Figure 5). We also measured a single instance of significant DOC export from the forest in May (Figure 5). Overall, the average import rate of DOC to the mangrove wetland during this dry season sampling was 0.0908 g C m⁻² h⁻¹, and 86% of this import took place during flood tide.

There was a net export of DOC from our mangrove site at SRS-6 during the wet season. The highest export rate of DOC to the water column (-0.5784 g C m⁻² h⁻¹) occurred in October, and the average of significant fluxes for this sampling was -0.0606 g C m⁻² h⁻¹ (Figure 5). Approximately, 65% of the DOC exported from the mangrove occurred during ebb tide, with the remainder being exported from the wetland within the flume and transported to interior forest during flood tide. Difficulties in sampling attributed to low tidal range during the December sampling resulted in the calculation of net areal flux for two of the four tidal cycles measured. These fluxes indicated both import and export of DOC with an average DOC import of 0.0265 g C m⁻² h⁻¹ during the 'Norte' season.

Relationships between DOC flux and half tide, season, and length of inundation were
2 determined with one-way ANOVA, LSD post hoc analysis and correlation analysis ($p < 0.05$).
No significant relationship between DOC flux and half tide was determined. Net areal DOC flux
4 ($\text{g m}^{-2} \text{ h}^{-1}$) in the dry season was significantly different than in the wet and 'Norte' seasons, with
net import occurring during the dry season. DOC import was positively correlated with [DOC]
6 and DOC export was negatively correlated with [DOC] (Pearson's correlation, $p < 0.05$).
Instantaneous flux (g DOC s^{-1}) was negatively correlated with length of inundation. Within
8 seasons, no significant relationship between [DOC] and length of inundation was determined.

To calculate an average yearly DOC flux, we took an average of the significant net areal
10 fluxes measured in this study ($-0.0403 \text{ g C m}^{-2} \text{ h}^{-1}$), multiplied by the product of the average
inundation period per tide (2 hr) and the approximate number of tides at the site during 2003
12 (696). Due to higher than average precipitation during the wet and dry seasons in 2003 and
lower precipitation during the 'Norte' season (Figure 1), this estimate equally weights the length
14 of each season in the calculation of an average annual net areal flux. The average inundation
period per tide is the mean period of each high tide where we would be able to sample the flume
16 using the approach we describe earlier. This is the point of the tide where approximately 10 cm
of water covered each end of the flume. Note, this does not reflect the full duration of inundation
18 that occurs at SRS-6. As a result our estimate is only applicable to the period of time around
high tide and to the fringe forest around the area of the flume location. The result of this
20 estimation was a net annual export of DOC from the fringe mangrove to both the tidal creek and
basin mangrove forest ($56 \text{ g C m}^{-2} \text{ yr}^{-1}$).

22

DISCUSSION

The magnitude and direction of material fluxes in estuarine systems involve processes occurring at different spatial and temporal scales. Small-scale processes occurring directly at wetland interfaces (i.e. water-soil, water-atmosphere or soil-atmosphere) are influenced by short-term variability in environmental conditions (i.e., material concentration, wind and rain events, duration of inundation, flushing time). Intermediate-scale processes occurring across areas of similar or adjacent habitats (i.e. tidal creek-fringing wetland, fringing wetland-inland wetland) are influenced by longer temporal scale processes such as seasonal and annual variability in environmental conditions. At larger spatial scales, such as the interface between the mangrove forest and river, we see an incorporation of more general processes related to estuarine mixing and long-term climate patterns controlling carbon and nutrient dynamics.

Water flux estimates and paired upstream/downstream measurements of [DOC] in our study were used to quantify fluxes between the mangrove surface (soil, prop root, epibiont communities, etc.) and inundating water column of a riverine mangrove wetland. Sampling over multiple consecutive tides within a season allowed for the examination of short-term variability in flux that occurs in the natural environment. We measured vertical flux and found daily variability in DOC exchange between the mangrove soil and inundating water column. The differences in vertical fluxes between tides were attributed to the length of inundation and flushing rate of the mangrove soil and the DOC concentration in the water column. We also found seasonal variation in flux and found net import of DOC to the mangrove wetland during the dry season (May), net export to the inundating water column during the wet season (October), and import to the mangrove soil during the 'Norte' season (December) in 2003. We used vertical flux and directional flow measurements to calculate horizontal flux and found during the dry season, the net DOC import was mainly from the tidal creek during flood tide and

during the wet season, the net export of DOC was mainly to the tidal creek from the wetland
2 during ebb tide.

Particulate organic carbon (POC) can represent a significant amount of the organic matter
4 export from mangrove systems, though it most often varies with season. Export of POC
measured in an Australian riverine mangrove forest was $420 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Boto & Bunt, 1981)
6 whereas studies in Florida have measured $64 \text{ g C m}^{-2} \text{ yr}^{-1}$ (Twilley, 1985) and $186 \text{ g C m}^{-2} \text{ yr}^{-1}$
(Heald, 1971). For all sampling periods in our study, DOC accounted for >95% of the TOC in
8 the water column. However, nets were not used in this study to capture leaf-size particulate
matter, which may make up a significant portion of export from mangrove forests (Boto & Bunt,
10 1981; Twilley, 1985). Several other mangrove studies have also measured a high DOC fraction
of TOC in the water column. Twilley (1985) estimated that up to 75% of all carbon exchanged
12 in a basin mangrove forest in Florida was DOC, Davis et al. (2001b) estimated >90% of the TOC
in mangrove island enclosures was DOC and Sutula et al. (2003) found that in Taylor River,
14 Florida approximately 98% of TOC was dissolved.

Due to the sampling technique and asymmetrical tidal patterns, water samples could only
16 be collected in the mangrove flume during high tide, and only during those high tides when the
mangrove forest was inundated. Samples at the beginning and end of a high tide may have been
18 influenced by the re-suspension of particulates produced by stronger currents flowing across the
soil surface. However, no significant difference in organic carbon content was detected between
20 filtered and unfiltered samples, indicating very low POC in the water column and little influence
of particulate re-suspension on the water samples.

22 Concentrations of DOC in this riverine mangrove system were slightly higher than
reported values for other mangrove systems (Table 1). At SRS-6, DOC concentrations were

highest in May 2003, possibly due to relatively high litter fall production ($4 \text{ gdw m}^{-2} \text{ d}^{-1}$) that
2 month (Twilley & Rivera-Monroy, unpublished data). Coincidentally, freshwater inflow was
lowest—as indicated by salinity—and tidal amplitude was also quite low (0.7 m) during this
4 sampling period.

Tidal amplitude and freshwater inflow play a factor in regulating DOC flux from
6 estuarine wetland soils by influencing the amount of time the soil interacts with the inundating
water column and by regulating the amount of water discharged across the surface. Increased
8 freshwater input has been shown to increase DOC flux from estuaries to adjacent coastal waters,
sometimes by as much as 300% (Miller, 1999). Vertical flux data in this study indicate that the
10 highest export of DOC from the wetland soil to the inundating water column occurred during the
wet season, when tidal amplitude was greatest (0.9 m) and freshwater inflow from the
12 Everglades was highest. These flux results indicate DOC export from the wetland is controlled
to a greater extent by freshwater discharge and tidal range, and is not necessarily positively
14 correlated with DOC concentrations.

Flux estimates for our riverine fringe mangrove site were similar in magnitude to findings
16 of previous mangrove carbon flux studies (Table 2). Seasonal fluxes indicated net import of
DOC to the wetland soil during the dry season and export of DOC to the inundating water
18 column during the wet season, with an overall net DOC export of $56 \text{ g DOC m}^{-2} \text{ yr}^{-1}$ from the
mangrove wetland to the adjacent tidal creek. Based on leaching rates of mangrove litter (leaves
20 and wood) from this site, leaching losses under conditions of inundation can account for about
43% of the DOC exported from the forest (Romigh, 2005). Previous estimates of total carbon
22 export from mangroves range from 2 to $400 \text{ g C m}^{-2} \text{ yr}^{-1}$ with an average of about $200 \text{ g C m}^{-2} \text{ yr}^{-1}$,
while salt marshes typically export about half this amount (Twilley, 1998). In a ‘eulerian’,

creek flux study in Australia, Boto & Wellington (1988) were unable to detect a net flux of
2 dissolved materials (including DOC) from a fringe mangrove forest. However, the site they
sampled received little or no freshwater inflows from adjacent uplands (Boto & Wellington,
4 1988).

Generally, terrestrially derived DOM undergoes conservative mixing down the estuarine
6 gradient, forming an inverse linear relationship between salinity and concentration. A previous
study conducted in Shark River, FL (Jaffe et al., 2004) found that fluxes of DOC are controlled
8 to a great extent by discharge rather than concentration, indicating lower DOC concentration
with increasing freshwater input. The wet and 'Norte' season data from our study show a similar
10 inverse linear relationship of salinity vs. DOC, supporting these findings by Jaffe et al. (2004;
Figure 6). Consistent high salinities (25–30 ‰)—due to low freshwater input to Shark River
12 during May—coupled with high variation in DOC concentration indicate both import and export
processes occurring at the wetland soil-water column interface even during the dry season.
14 However, the non-linear relationship between salinity and DOC concentration during May might
be an indication that these processes are not occurring at the same rate as those during other
16 seasons. The energy signature of riverine mangrove forests is higher than in fringe, basin and
scrub forests (Twilley, 1995), which may help explain why fluxes from the forest at SRS-6 were
18 higher than those reported in other mangrove systems.

20 *Conclusions*

Results from our study support the hypothesis that highly productive, riverine mangrove
22 forests are sources of organic carbon to adjacent coastal waters. The flux of DOC at the
mangrove soil-water column interface is seasonal, perhaps in response to regional freshwater

inflow, as inferred from Shark River salinity patterns. There was variation in the direction and magnitude of organic carbon flux between consecutive tides. However, when sampling over multiple tidal cycles, trends in DOC flux emerged—with import of DOC to the mangrove soil during the dry season and export to the water column during the wet season. Overall, this mangrove forest showed a net annual export of 56 g DOC m⁻² yr⁻¹.

Import of DOC to the mangrove during the dry season and export during the wet season suggests that freshwater discharge may indeed have a strong influence on the direction of organic carbon flux to this system. It is unclear, however, whether carbon exchange between the forest canopy and atmosphere mimic these patterns. It is also unclear whether an increase in freshwater flow to the Everglades would significantly affect DOC flux.

This study only addressed a one-year period and cannot identify long-term trends in organic carbon flux for this system. Continued research would help determine the effect of long-term hydrologic influences (drought or regulated increases in freshwater flow) on carbon cycling in mangroves. Combining the quantitative flux data from the mangrove soil-water column interface with atmospheric flux data would certainly help to estimate the carbon budget in these wetlands.

ACKNOWLEDGEMENTS

We thank Justin Baker, Dan Bond and Edward Castañeda for their help in flume design and construction. We greatly appreciate the support and assistance provided by Dan Childers, the Wetland Ecosystems Lab at Florida International University, and Rachel Butzler. We would also like to thank Dan Childers and an anonymous referee for their comments on the manuscript. This study was partially funded by the Texas Water Resources Institute and the Sustainable

Coastal Margins Program at Texas A&M University. This material is based upon work

2 supported by the National Science Foundation under Grant No. 9910514.

LITERATURE CITED

- 2 Benner, R., E. Peele, & R. E. Hodson. 1986. Microbial utilization of dissolved organic
matter from leaves of the red mangrove, *Rhizophora mangle*, in the Fresh Creek Estuary,
4 Bahamas. *Estuarine, Coastal and Shelf Science* 23:607-619.
- 6 Boto, K. G. & J. S. Bunt. 1981. Tidal export of particulate organic matter from a northern
Australian mangrove system. *Estuarine, Coastal and Shelf Science* 13:247-255.
- 8 Boto, K. G. & J. T. Wellington. 1988. Seasonal variations in concentrations and fluxes of
10 dissolved organic and inorganic materials in a tropical, tidally-dominated, mangrove
waterway. *Marine Ecology Progress Series* 50:151-160.
- 12 Chen, R. & R. R. Twilley. 1999. Patterns of mangrove forest structure and soil nutrient
14 dynamics along the Shark River Estuary, Florida. *Estuaries* 22:955-970.
- 16 Childers, D. L. & J. W. Day Jr. 1988. A flow-through flume technique for quantifying
nutrient and materials fluxes in microtidal estuaries. *Estuarine, Coastal and Shelf Science*
18 27:483-494.
- 20 Childers, D. L., J. N. Boyer, S. E. Davis, C. J. Madden, D. T. Rudnick & F. H. Sklar.
2005. Relating precipitation and water management to nutrient concentrations in the
22 oligotrophic “upside down” estuaries of the Florida Everglades. *Limnology and
Oceanography in press*.
- 24 Davis, S., D. Childers, J. Day Jr., D. Rudnick, & F. Sklar. 2001a. Nutrient dynamics in
26 vegetated and unvegetated areas of a Southern Everglades mangrove creek. *Estuarine,
Coastal and Shelf Science* 52:753-768.
- 28 Davis, S., D. Childers, J. Day Jr., D. Rudnick, & F. Sklar. 2001b. Wetland-water column
30 exchanges of carbon, nitrogen, and phosphorus in a Southern Everglades dwarf
mangrove. *Estuaries* 24:610-622.
- 32 Davis, S., C. Corronado-Molina, D. Childers, & J. W. Day Jr. 2003. Temporally
34 dependent C, N, and P dynamics associated with the decay of *Rhizophora mangle* L. leaf
litter in oligotrophic mangrove wetlands of the Southern Everglades. *Aquatic Botany*
36 75:199-215.
- 38 Dittmar, T. & R. J. Lara. 2001a. Do mangroves rather than rivers provide nutrients to
coastal environments south of the Amazon River? Evidence from long-term flux
40 measurements. *Marine Ecology Progress Series* 213:67-77.
- 42 Dittmar, T. & R. J. Lara. 2001b. Driving forces behind nutrient and organic matter
dynamics in a mangrove tidal creek in North Brazil. *Estuarine, Coastal and Shelf Science*
44 52:249-259.

- 2 Dittmar, T., R. J. Lara, & G. Kattner. 2001. River or mangrove? Tracing major organic
matter sources in tropical Brazilian coastal waters. *Marine Chemistry* 73:253-271.
- 4 Feller, I. C. 2002. The role of herbivory by wood-boring insects in mangrove ecosystems
in Belize. *Oikos* 97:167-176.
- 6 Furukawa, K., E. Wolanski, & H. Mueller. 1997. Currents and sediment transport in
8 mangrove forests. *Estuarine, Coastal and Shelf Science* 44:301-310.
- 10 Heald, E. 1971. The production of organic detritus in a South Florida estuary. University
of Miami: Sea Grant Technical Bulletin 6:110.
- 12 Hogarth, P. J. 1999. *The Biology of Mangroves*. Oxford University Press. UK. 228 p.
- 14 Jaffe, R., J. N. Boyer, X. Lu, N. Maie, C. Yang, N. M. Scully, & S. Mock. 2004. Source
16 characterization of dissolved organic matter in a subtropical mangrove-dominated estuary
by fluorescence analysis. *Marine Chemistry* 84:195-210.
- 18 Lee, S. Y. 1995. Mangrove outwelling: a review. *Hydrobiologia* 295:203-212.
- 20 Lewis III, R. R., R. G. Gilmore Jr., D. W. Crewz, & W. E. Odum. 1985. Mangrove
22 habitat and fishery resources of Florida. p. 281-336. *In* Seaman Jr., W. E. (ed.) Florida
Aquatic Habitat and Fishery Resources. Florida Chapter of the American Fisheries
24 Society, Kissimmee, Florida.
- 26 Machiwa, J. F. & R. O. Hallberg. 2002. An empirical model of the fate of organic carbon
in a mangrove forest partly affected by anthropogenic activity. *Ecological Modelling*
28 147:69-83.
- 30 Miller, A. E. J. 1999. Seasonal investigations of dissolved organic carbon dynamics in the
Tamar Estuary, U.K. *Estuarine, Coastal and Shelf Science* 49:891-908.
- 32 Moran, M. A., R. J. Wicks, & R. E. Hodson. 1991. Export of dissolved organic matter
34 from a mangrove swamp ecosystem: evidence from natural fluorescence, dissolved lignin
phenols, and bacterial secondary production. *Marine Ecology Progress Series* 76:175-
36 184.
- 38 Odum, W. E. & E. J. Heald. 1975. The detritus-based food web of an estuarine mangrove
community. p. 265-286. *In* L. E. Cronin, editor. *Estuarine Research*, Volume 1.
40 Academic Press, Inc., New York.
- 42 Rivera-Monroy, V. H., J. Day, R. Twilley, F. Vera-Herrera, & C. Coronado-Molina.
1995. Flux of nitrogen and sediment in a fringe mangrove forest in Terminos Lagoon,
44 Mexico. *Estuarine, Coastal and Shelf Science* 40:139-160.

- 2 Robertson, A. I. 1986. Leaf-burying crabs: their influence on energy flow and export
4 from mixed mangrove forests (*Rhizophora* spp.) in northeastern Australia. *Journal of*
6 *Experimental Marine Biology and Ecology* 102:237-248.
- 8 Robertson, A. & S. Blaber. 1992. Plankton, epibenthos, and fish communities. p. 173-
10 224. *In* A. Robertson & D. Alongi (eds.) *Tropical Mangrove Ecosystems, Coastal and*
12 *Estuarine Studies Series*, American Geophysical Union, Washington, D.C.
- 14 Robertson, A., D. Alongi & K. Boto. 1992. Food chains and carbon fluxes. p. 293-326.
16 *In* A. Robertson & D. Alongi (eds.) *Tropical Mangrove Ecosystems, Coastal and*
18 *Estuarine Studies Series*, American Geophysical Union, Washington, D.C.
- 20 Romero, L.M., T.J. Smith, III, & J. Fourqurean. 2005. Changes in mass and nutrient
22 content of wood during decomposition in a south Florida mangrove forest. *Journal of*
24 *Ecology* 93:618-631.
- 26 Romigh, M. 2005. Organic carbon flux at the mangrove soil-water column interface in
28 the Florida Coastal Everglades. M.S. Thesis. Texas A&M University, College Station,
30 Texas. 45 pp.
- 32 Sutula, M., B. Perez, E. Reyes, D. Childers, S. Davis, J. Day, Jr., D. Rudnick, & F. Sklar.
34 2003. Factors affecting spatial and temporal variability in material exchange between the
36 Southern Everglades wetlands and Florida Bay (USA). *Estuarine, Coastal and Shelf*
38 *Science* 57:757-781.
- 40 Twilley, R. 1985. The exchange of organic carbon in basin mangrove forests in a
southwest Florida estuary. *Estuarine, Coastal and Shelf Science* 20:543-557.
- Twilley, R. 1988. Coupling of mangroves to the productivity of estuarine and coastal
waters. *Lecture Notes on Coastal and Estuarine Studies* 22:155-180.
- Twilley, R. 1995. Properties of mangrove ecosystems related to the energy signature of
coastal environments. p. 43-62. *In* C. A. S. Hall (ed.) *Maximum Power*.
University Press of Colorado, Niwot.
- Twilley, R. 1998. Mangrove Wetlands. p. 445-473. *In* M. G. Messina & W. H. Conner,
editors. *Southern Forested Wetlands: Ecology and Management*. Lewis Publishers, Boca
Raton, Florida.
- Welch, R., M. Madden, & R. Doren. 1999. Mapping the Everglades. *Photogrammetric*
Engineering and Remote Sensing. 65:163-170.

Table 1. Comparison of surface water DOC and TOC concentrations (in mg l⁻¹) from various mangrove studies.

Location	Wetland Type	Source	TOC	DOC
Florida	riverine mangrove	This study		1.7-17.9
Florida	dwarf & fringe mangrove	Sutula et al. (2003)		8.4-19.2
Zanzibar	fringe mangrove	Machiwa & Hallberg (2002)		0.78-1.28
Florida	fringe mangrove	Davis et al. (2001a)		8.4-21.6
Florida	dwarf mangrove	Davis et al. (2001b)	10.8-18	8.4-18
Brazil	riverine mangrove	Dittmar & Lara (2001)	7.20	4.32
Australia	fringe mangrove	Furukawa et al. (1997)	2.67	2.21
Bahamas	fringe mangrove	Moran et al. (1991)		2.3
Australia	fringe mangrove	Boto & Wellington (1988)		1-2
Florida	basin mangrove	Twilley (1985)	9.4-21	

Table 2. Estimates of net annual flux of TOC and DOC ($\text{g C m}^{-2} \text{ yr}^{-1}$) from various mangrove wetlands.

Location	TOC	DOC	Method	Source
Florida		-56	flume	This Study
Florida	-7.1		creek flux	Sutula et al. (2003)
Florida		3.04	in-channel flume	Davis et al. (2001a)
Florida		- 381	mangrove enclosures	Davis et al. (2001b)
Florida	-64		flume	Twilley (1985)
Australia		7.3	creek flux	Boto & Wellington (1988)
World Average	-210		review of literature	Twilley (1998)

List of Figures

2 Figure 1. Seasonal pattern of mean (\pm sd) temperature (open squares) and precipitation
(open circles) in south Florida from 1994–2003 (from USGS). Data from 2003 (closed
4 circles) mirrored the seasonal pattern exhibited across this 10-year record.


6 Figure 2. Map of South Florida with the location of the experimental flume study site
(SRS-6) along Shark River (Everglades National Park).


8
Figure 3. Plots of predicted water level in the flume during each 2003 sampling event.
10 Flume water levels calculated from a regression relationship between periodic flume
water level measurements and a permanent gauge along Shark River (SRS-6).


12
Figure 4. Box-and-whisker plots of (a) temperature, (b) salinity, (c) DOC concentration
14 and (d) TSS concentration at the study site for each 2003 sampling. Different letters
indicate significant differences among sampling periods (ANOVA, $p < 0.05$).





16
Figure 5. Net areal fluxes of DOC measured in each tidal cycle of the 2003 sampling
18 periods (M = May, O = October, and D = December). Positive values indicate uptake by
the wetland; negative values indicate release to the inundating water column. Asterisks
20 indicate fluxes that were not significantly different from zero. 'nd' = No flux data.

22 Figure 6. Scatter plot of salinity vs. DOC concentration at SRS-6 for all three sampling
periods in 2003.










2



▨ Import from tidal creek

▩ Import from mangrove forest

□ Export to tidal creek

■ Export to mangrove forest

