Florida International University FIU Digital Commons

All Faculty

6-4-2015

Draft Genome Sequence of Burkholderia sp. MR1, a Methylarsenate-Reducing Bacterial Isolate from Florida Golf Course Soil

Shashank S. Pawitwar Herbert Wertheim College of Medicine, Florida International University, spawitwa@fiu.edu

Sagar M. Utturkar University of Tennessee

Steven D. Brown University of Tennessee; Oak Ridge National Laboratory

Masafumi Yoshinaga Herbert Wertheim College of Medicine, Florida International University, myoshina@fiu.edu

Barry P. Rosen Herbert Wertheim College of Medicine, Florida International University, brosen@fu.edu

Follow this and additional works at: https://digitalcommons.fiu.edu/all_faculty

Recommended Citation

Pawitwar, Shashank S.; Utturkar, Sagar M.; Brown, Steven D.; Yoshinaga, Masafumi; and Rosen, Barry P., "Draft Genome Sequence of Burkholderia sp. MR1, a Methylarsenate-Reducing Bacterial Isolate from Florida Golf Course Soil" (2015). *All Faculty*. 85. https://digitalcommons.fu.edu/all_faculty/85

This work is brought to you for free and open access by FIU Digital Commons. It has been accepted for inclusion in All Faculty by an authorized administrator of FIU Digital Commons. For more information, please contact dcc@fu.edu.





Draft Genome Sequence of *Burkholderia* sp. MR1, a Methylarsenate-Reducing Bacterial Isolate from Florida Golf Course Soil

Shashank S. Pawitwar,^a Sagar M. Utturkar,^b Steven D. Brown,^{b,c} Masafumi Yoshinaga,^a Barry P. Rosen^a

Department of Cellular Biology and Pharmacology, Herbert Wertheim College of Medicine, Florida International University, Miami, Florida, USA^a; Graduate School of Genome Science and Technology, University of Tennessee, Knoxville, Tennessee, USA^b; Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA^c

To elucidate the environmental organoarsenical biocycle, we isolated a soil organism, *Burkholderia* sp. MR1, which reduces relatively nontoxic pentavalent methylarsenate to the more toxic trivalent methylarsenite, with the goal of identifying the gene for the reductase. Here, we report the draft genome sequence of *Burkholderia* sp. MR1.

Received 5 May 2015 Accepted 7 May 2015 Published 4 June 2015

Citation Pawitwar SS, Utturkar SM, Brown SD, Yoshinaga M, Rosen BP. 2015. Draft genome sequence of *Burkholderia* sp. MR1, a methylarsenate-reducing bacterial isolate from Florida golf course soil. Genome Announc 3(3):e00608-15. doi:10.1128/genomeA.00608-15.

Copyright © 2015 Pawitwar et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Address correspondence to Barry P. Rosen, brosen@fiu.edu.

rsenic is considered to be the most prevalent environmental toxin by the U.S. Environmental Protection Agency (EPA). Prolonged exposure to arsenic in drinking water results in diseases such as skin, bladder, and prostate cancer (1-3). Pentavalent organoarsenical herbicides such as monosodium methylarsenate (MSMA) [MAs(V)] undergo environmental reduction to the more toxic trivalent species and eventual C-As bond cleavage to produce arsenite [As(III)] (4), which contaminates our drinking water supplies (5). The poultry growth promoter roxarsone (4hydroxy-3-nitrophenylarsonic acid) has been proposed to undergo a similar pathway of reduction and C-As bond cleavage (6). The arsI gene of soil organisms encodes a C-As bond lyase that catalyzes the second reaction, but the pathway for reduction of MAs(V) to MAs(III) is unknown. The objective of this study was to identify the gene(s) and enzyme(s) responsible for MAs(V) reduction.

MSMA is applied as an herbicide to golf courses in Florida. We isolated an MAs(V)-reducing organism from simulated golf course soil. From the 16S rRNA gene sequence, it is a Gramnegative bacterial strain closely related to Burkholderia glathei (4). The environmental isolate has been named Burkholderia sp. MR1. Genome sequencing was performed using an Illumina HiSeq platform, with quality-based trimming, as described previously (7). After trimming, 6,727,114 paired-end reads remained, with an average read length of 90 bp, comprising genome coverage of $100 \times$. After evaluation of several approaches (8), optimal assembly was obtained through SPAdes software (version 3.1.1). The assembly consisted of 58 large contigs (\geq 500 bp), with a total genome size of 6.01 Mb, an N₅₀ contig size of 244 kb, and the largest contig of 432 kb. Gene prediction and annotation were performed at Oak Ridge National Laboratory, as described previously (9). The draft genome sequence has 5,554 candidate protein-coding genes and a G+C content of 63.2%.

The genomic sequence of *Burkholderia* sp. MR1 has a single *ars* operon comprising the genes *arsH* [encoding an NADPH-dependent flavin mononucleotide (FMN) MAs(III) oxidase] (10), *acr3* [encoding an As(III) efflux permease] (11), and *arsR* [encod-

ing an As(III)-responsive repressor protein] (12). The operon is on the complementary strand, so the putative operon structure is *arsR-acr3-arsH*. Note that ArsH detoxifies MAs(III) (10) and so may protect this organism from the toxicity of the product of the reductase. None of these gene products catalyze MAs(V) reduction, so the gene or genes for reductase activity must be elsewhere in the chromosome. In the genome there are 195 genes annotated as reductases that are candidates for the MAs(V) reductase. However, further biochemical or genetic analysis is required to identify the responsible gene(s).

Nucleotide sequence accession number. The draft genome sequence of strain AE038-8 has been deposited at DDBJ/EMBL/ GenBank (accession number JWHM00000000). The version described in this paper is the first version.

ACKNOWLEDGMENTS

This research was supported by NIH grant R37 GM55425 (to B.P.R.) and FIU HWCOM Pilot Grant ID800004594 (to M.Y.). We acknowledge the Oak Ridge National Laboratory, managed by UT-Battelle, LLC, for support under DOE contract DE-AC05-00OR22725.

REFERENCES

- 1. National Research Council. 2001. Arsenic in drinking water: 2001 update. National Academies Press, Washington, DC.
- Tchounwou PB, Centeno JA, Patlolla AK. 2004. Arsenic toxicity, mutagenesis, and carcinogenesis—a health risk assessment and management approach. Mol Cell Biochem 255:47–55. http://dx.doi.org/10.1023/ B:MCBI.0000007260.32981.b9.
- Abernathy CO, Thomas DJ, Calderon RL. 2003. Health effects and risk assessment of arsenic. J Nutr 133:15365–1538S.
- Yoshinaga M, Cai Y, Rosen BP. 2011. Demethylation of methylarsonic acid by a microbial community. Environ Microbiol 13:1205–1215. http:// dx.doi.org/10.1111/j.1462-2920.2010.02420.x.
- Feng M, Schrlau JE, Snyder R, Snyder GH, Chen M, Cisar JL, Cai Y. 2005. Arsenic transport and transformation associated with MSMA application on a golf course green. J Agric Food Chem 53:3556–3562. http:// dx.doi.org/10.1021/jf047908j.
- Yoshinaga M, Rosen BP. 2014. A C·As lyase for degradation of environmental organoarsenical herbicides and animal husbandry growth promoters. Proc Natl Acad Sci U S A 111:7701–7706. http://dx.doi.org/ 10.1073/pnas.1403057111.

- Woo HL, Utturkar S, Klingeman D, Simmons BA, Deangelis KM, Brown SD, Hazen TC. 2014. Draft genome sequence of the lignindegrading *Burkholderia* sp. strain LIG30, isolated from wet tropical forest soil. Genome Announc 2(3):e00637-14. http://dx.doi.org/10.1128/ genomeA.00637-14.
- Utturkar SM, Klingeman DM, Land ML, Schadt CW, Doktycz MJ, Pelletier DA, Brown SD. 2014. Evaluation and validation of *de novo* and hybrid assembly techniques to derive high-quality genome sequences. Bioinformatics 30:2709–2716. http://dx.doi.org/10.1093/bioinformatics/ btu391.
- Brown SD, Utturkar SM, Klingeman DM, Johnson CM, Martin SL, Land ML, Lu TY, Schadt CW, Doktycz MJ, Pelletier DA. 2012. Twentyone genome sequences from *Pseudomonas* species and 19 genome se-

quences from diverse bacteria isolated from the rhizosphere and endosphere of *Populus deltoides*. J Bacteriol **194**:5991–5993. http://dx.doi.org/ 10.1128/JB.01243-12.

- Chen J, Bhattacharjee H, Rosen BP. 3 March 2015. ArsH is an organoarsenical oxidase that confers resistance to trivalent forms of the herbicide monosodium methylarsenate and the poultry growth promoter roxarsone. Mol Microbiol. 2015. http://dx.doi.org/10.1111/mmi.12988.
- 11. Yang HC, Fu HL, Lin YF, Rosen BP. 2012. Pathways of arsenic uptake and efflux, p 325–358. *In* Lutsenko S, Argüello JM (ed), Metal transporters. Elsevier, Amsterdam, Netherlands.
- 12. Xu C, Rosen BP. 1999. Metalloregulation of soft metal resistance pumps, p 5–19. *In* Sarkar B (ed), Metals and genetics. Kluwer Academic/Plenum Publishers, New York, NY.