

7-9-2015

Genetics of primary ovarian insufficiency: new developments and opportunities

Yingying Qin

Center for Reproductive Medicine, Shandong Provincial Hospital, Shandong University, National Research Center for Assisted Reproductive Technology and Reproductive Genetics, The Key Laboratory for Reproductive Endocrinology of Ministry of Education, Shandong Provincial Key Laboratory of Reproductive Medicine

Xue Jiao

Center for Reproductive Medicine, Shandong Provincial Hospital, Shandong University, National Research Center for Assisted Reproductive Technology and Reproductive Genetics, The Key Laboratory for Reproductive Endocrinology of Ministry of Education, Shandong Provincial Key Laboratory of Reproductive Medicine

Joe Leigh Simpson

Research and Global Programs March of Dimes Foundation; Department of Human and Molecular Genetics, Herbert Wertheim College of Medicine, Florida International University, simpsonj@fiu.edu

Zi-Jiang Chen

Center for Reproductive Medicine, Shandong Provincial Hospital, Shandong University, National Research Center for Assisted Reproductive Technology and Reproductive Genetics, The Key Laboratory for Reproductive Endocrinology of Ministry of Education, Shandong Provincial Key Laboratory of Reproductive Medicine, Jinan 250001, China. Renji Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai

Follow this and additional works at: https://digitalcommons.fiu.edu/all_faculty

Recommended Citation

Qin, Yingying; Jiao, Xue; Simpson, Joe Leigh; and Chen, Zi-Jiang, "Genetics of primary ovarian insufficiency: new developments and opportunities" (2015). *All Faculty*. 94.
https://digitalcommons.fiu.edu/all_faculty/94

Genetics of primary ovarian insufficiency: new developments and opportunities

Yingying Qin¹, Xue Jiao¹, Joe Leigh Simpson^{2,3}, and Zi-jiang Chen^{1,4,*}

¹Center for Reproductive Medicine, Shandong Provincial Hospital, Shandong University, National Research Center for Assisted Reproductive Technology and Reproductive Genetics, The Key Laboratory for Reproductive Endocrinology of Ministry of Education, Shandong Provincial Key Laboratory of Reproductive Medicine, Jinan 250001, China ²Research and Global Programs March of Dimes Foundation, White Plains, NY, USA ³Department of Human and Molecular Genetics, Herbert Wertheim College of Medicine, Florida International University, Miami, FL, USA ⁴Renji Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai 200127, China

*Correspondence address. chenzijiang@vip.163.com

Submitted on March 6, 2015; resubmitted on June 2, 2015; accepted on July 9, 2015

TABLE OF CONTENTS

- Introduction
- Methods
- Results
 - Chromosomal abnormalities in POI
 - Single genes causing non-syndromic POI
 - Pleiotropic single gene disorders in POI
 - Mitochondrial genes causing POI
 - Multiple malformation syndromes characterized by POI
 - Genome-wide studies in POI
 - Regulatory genes and networks
- Conclusion: current status of genes causing POI

BACKGROUND: Primary ovarian insufficiency (POI) is characterized by marked heterogeneity, but with a significant genetic contribution. Identifying exact causative genes has been challenging, with many discoveries not replicated. It is timely to take stock of the field, outlining the progress made, framing the controversies and anticipating future directions in elucidating the genetics of POI.

METHODS: A search for original articles published up to May 2015 was performed using PubMed and Google Scholar, identifying studies on the genetic etiology of POI. Studies were included if chromosomal analysis, candidate gene screening and a genome-wide study were conducted. Articles identified were restricted to English language full-text papers.

RESULTS: Chromosomal abnormalities have long been recognized as a frequent cause of POI, with a currently estimated prevalence of 10–13%. Using the traditional karyotype methodology, monosomy X, mosaicism, X chromosome deletions and rearrangements, X-autosome translocations, and isochromosomes have been detected. Based on candidate gene studies, single gene perturbations unequivocally having a deleterious effect in at least one population include Bone morphogenetic protein 15 (*BMP15*), Progesterone receptor membrane component 1 (*PGRMC1*), and Fragile X mental retardation 1 (*FMR1*) premutation on the X chromosome; Growth differentiation factor 9 (*GDF9*), Folliculogenesis specific bHLH transcription factor (*FIGLA*), Newborn ovary homeobox gene (*NOBOX*), Nuclear receptor subfamily 5, group A, member 1 (*NR5A1*) and Nanos homolog 3 (*NANOS3*) seem likely as well, but mostly being found in no more than 1–2% of a single population studied. Whole genome approaches have utilized genome-wide association studies (GWAS) to reveal loci not predicted on the basis of a candidate gene, but it remains difficult to locate causative genes and susceptible loci were not always replicated. Cytogenomic methods (array CGH) have identified other regions of interest but studies have not shown consistent results, the resolution of arrays has varied and replication is uncommon. Whole-exome sequencing in non-syndromic POI kindreds has only recently begun, revealing mutations in the Stromal antigen

3 (*STAG3*), Synaptonemal complex central element 1 (*SYCE1*), minichromosome maintenance complex component 8 and 9 (*MCM8*, *MCM9*) and ATP-dependent DNA helicase homolog (*HFM1*) genes. Given the slow progress in candidate-gene analysis and relatively small sample sizes available for GWAS, family-based whole exome and whole genome sequencing appear to be the most promising approaches for detecting potential genes responsible for POI.

CONCLUSION: Taken together, the cytogenetic, cytogenomic (array CGH) and exome sequencing approaches have revealed a genetic causation in ~20–25% of POI cases. Uncovering the remainder of the causative genes will be facilitated not only by whole genome approaches involving larger cohorts in multiple populations but also incorporating environmental exposures and exploring signaling pathways in intragenic and intergenic regions that point to perturbations in regulatory genes and networks.

Key words: primary ovarian insufficiency / premature ovarian failure / chromosomal abnormality / gene mutation / genome-wide association studies / whole-exome sequencing / next generation sequencing

Introduction

Primary ovarian insufficiency (POI), also known as premature ovarian failure (POF) or premature menopause, is defined as cessation of menstruation before the expected age of menopause. This age is traditionally defined to be prior to 40 years and diagnosis is confirmed by elevated serum FSH levels ($>40\text{IU/l}$). Although frequently stated that ~1% of the population is affected with POI before the age of 40 years and 0.1% before age 30 years, the prevalence is actually less certain (Coulam et al., 1986).

The disorder is clearly heterogeneous, with a wide spectrum of causes, namely cytogenetic, genetic, infectious or iatrogenic. Auto-immune and metabolic etiologies may or may not be genetic. Irrespective, etiology remains to be elucidated in most cases and until a decade ago few specific causes were known beyond X-chromosomal abnormalities, Fragile X mental retardation 1 (*FMR1*) premutation and FSH receptor (*FSHR*) in the Finnish population (Simpson, 1975; Aittomäki et al., 1995; Wittenberger et al., 2007). Most cases of isolated POI still appear sporadically, but ~10–15% has an affected first-degree relative, indicating significant genetic etiology (Van Kasteren et al., 1999). Pedigrees with multiple affected relatives are not rare (recessive and dominant). Presence of POI as one component of a pleiotropic genetic disorder is also well recognized. Yet identifying precise causative genes has been challenging. Here, we enumerate known genetic causes of POI, most elucidated within the last 5–10 years.

Confusion exists concerning nomenclature, namely the use of POF or POI. It is the view of the authors that POI can be taken to encompass occult, biochemical and overt stages, whereas POF is best considered as only the final stage of POI. The designation POI is thus best reserved as alluding to the entire gamut of disorders having diminished ovarian reserve—occult, subclinical, iatrogenic. Although many authors espouse POI in lieu of POF, the canonical genetic reference—Online Mendelian inheritance in man (OMIM)—has long used, and continues to use POF, to nominate causative genes. These designations now apply to POF1–POF9 (Supplementary Table S1), and the list growing.

Methods

A search for original articles published up to May 2015 was performed using PubMed and Google Scholar to identify studies on genetic variants associated with the human disease. The key word combinations include 'premature ovarian failure', 'primary ovarian insufficiency', 'early menopause', 'genetic', 'gene mutation', 'variant' and 'genome wide study'. For a study to be included in our review, it had to focus on chromosomal analysis, candidate

gene screening, or a genome-wide study in different POI cohorts. In addition, studies on mitochondrial genes causing POI and multiple malformation syndromes characterized by POI were included. Reports on the role of candidate genes in animal models were not included. Where appropriate, reference lists of identified articles were also searched for further relevant papers. However, articles identified were restricted to English language full-text papers.

Results

Chromosomal abnormalities in POI

Chromosomal abnormalities have long been recognized as a cause of POI, but percentages vary widely among reported series. This clearly reflects biases of ascertainment, for example reflecting whether a cohort was derived from a referral cytogenetic lab, a gynecologic practice, or a pediatric practice. Numerous different karyotypic anomalies have been found, ranging from numerical defects (monosomy X; X chromosomal mosaicism), X-deletions, X-autosome translocations, and X-isochromosomes and other rearrangements. Aggregate frequency of chromosomal abnormalities in reported studies is summarized in Table 1. Small sample sizes as well as selection biases and differing ages of ascertainment probably account for different prevalence in different populations. However, each of the five largest studies with respect to sample size reported frequencies between 10.0 and 12.9% (Zhang et al., 2003; Lakhali et al., 2010a; Baronchelli et al., 2011; Jiao et al., 2012; Kalantari et al., 2013); thus, a prevalence of 10–13% seems reasonable.

Numerical defects

The X chromosome has long been known to play an essential role in the maintenance of ovarian development and function. Females lacking an X chromosome as well as those showing an extra X chromosome are predisposed to developing POI.

45,X and 45,X/46,XX

Turner syndrome, often but not universally associated with X monosomy, leads to ovarian dysgenesis and accelerated follicular atresia. X monosomy without mosaicism is more typically found in primary amenorrhea and cases were almost universally understood to present with this phenotype. However, many early series were recruited from pediatric clinics, not among adult women. In 1975, Simpson (1975) reported that 3% (5/178) of 45,X patients actually menstruated. 45,X/46,XX and other forms of association also are associated with secondary

Table 1 Frequency of chromosomal abnormalities (CA) in different population studies.

Reference	Frequency of CA (%)	No. of CA	Sample size	Clinical characteristics	Population
Ayed <i>et al.</i> (2014)	18.0	18	100	PA, SA	Tunisian
Kalantari <i>et al.</i> (2013)	10.05	18	179	PA, SA	Iranian
Jiao <i>et al.</i> (2012)	12.1	64	531	PA, SA	Chinese (Jinan, Beijing, Shenzhen)
Baronchelli <i>et al.</i> (2011)	10.0	27	269	PA, SA, EM	Italian
Lakhal <i>et al.</i> (2010a)	10.8	108	1000	PA, SA	Tunisian
Ceylaner <i>et al.</i> (2010)	25.3	19 ^a	75	SA	Turkish
Janse <i>et al.</i> (2010)	12.9	19	147	SA	Dutch
Portnoi <i>et al.</i> (2006)	8.8	8	90	PA, SA	French
Zhang <i>et al.</i> (2003)	12.5	13	104	POI	Chinese (Chongqing)
Devi and Benn (1999)	13.3	4	30	SA	American
Davison <i>et al.</i> (1998)	2.5	2	79	PA, SA FSH > 20 IU/l	English
Castillo <i>et al.</i> (1992)	32.0	15	47	POI	Chilean
Rebar and Connolly (1990)	25.4	16	63	PA, SA	American

Chromosomal 'abnormalities' means visible structural changes in karyotype that are sufficiently large to cause clinical abnormalities. Variants (e.g. prominent satellites) are not included. CA, chromosomal abnormalities; PA, primary amenorrhea; POI, primary ovarian insufficiency; SA, secondary amenorrhea; EM, early menopause.

^aIncluding 2 46,XY gonadal dysgenesis (Swyer syndrome).

amenorrhea (POI). Either haploinsufficiency of pivotal genes on the X chromosome or non-specific meiotic impairment could explain the accelerated atresia of 45,X oocytes. Variability would be expected given potential heterozygosity of alleles in genes subjected to X-inactivation.

47,XXX

47,XXX women may experience oligomenorrhea, secondary amenorrhea, and early menopause, but relative risk has not been well studied. Goswami *et al.* (2003) reported the prevalence of 47,XXX in 52 women with POI to be 3.8%, whereas in our much larger Chinese series we observed 1.5% (8/531) (Jiao *et al.*, 2012). The presence of three X chromosomes plausibly leads to meiotic disturbance and, secondarily, ovarian failure. Additionally, overexpression of genes that escape X-inactivation could cause POI in 47,XXX. Mechanisms remain to be defined (Tartaglia *et al.*, 2010). A confounder is that an association exists between 47,XXX and autoimmune diseases (Holland, 2001; Goswami *et al.*, 2003).

X-structural abnormalities and X-autosome translocations

X chromosome deletions and X-autosome balanced translocations have long been observed in POI and were once the only approach available to localize causative genes. This strategy was illustrated in the 1970s by a region on the X chromosome appearing critical for the POI phenotype (Sarto *et al.*, 1973). A critical region was delineated that gave boundaries for breakpoints of X-autosome translocations associated with ovarian failure. This region extends from Xq13-Xq21 (POI2) to Xq23-q27 (POI1). It has been proposed by Rizzolio and colleagues that Xq13-Xq21 governs epigenetic regulations that down-regulate oocyte-expressed autosomal genes (Rizzolio *et al.*, 2006, 2007, 2009).

Irrespective of mechanisms involved in the critical region, almost all terminal deletions originating at Xq13 are associated with primary amenorrhea, lack of breast development and complete ovarian failure (Simpson and Rajkovic, 1999; Simpson, 2008). By contrast, terminal

deletions arising at Xq25 or Xq26 are characterized by the more common phenotype being not primary amenorrhea but premature ovarian failure. The gene designation POI1 is applied to this region. The more distal deletions arise at Xq27 or Xq28 and exert a less severe effect on stature and reproductive function than do proximal deletions (Simpson, 1975; Simpson and Rajkovic, 1999). Gene(s) in this POI-causing region are considered distinct from Fragile X mental retardation 1 (*FMR1*), located at Xq27 and premutation of which is the most common single cause of POI.

Multiple genes on the X chromosome have been identified by X-autosomal translocations. These include Diaphanous-related formin 2 (*DIAPH2*, Xq22) (Bione *et al.*, 1998), X-prolyl aminopeptidase (aminopeptidase P) 2, membrane-bound (*XPNPEP2*, Xq25) (Prueitt *et al.*, 2000), Dachshund family transcription factor 2 (*DACH2*, Xq21.3) (Prueitt *et al.*, 2002), Premature ovarian failure, 1B (*POF1B*, Xq21.1) (Lorda-Sanchez *et al.*, 2000; Bione *et al.*, 2004), Choroideremia (*CHM*, Xq21.1) (Lorda-Sanchez *et al.*, 2000; Mansouri *et al.*, 2008), Progesterone receptor membrane component 1 (*PGRMC1*, Xq24) (Mansouri *et al.*, 2008), Collagen, type IV, alpha 6 (*COL4A6*, Xq22.3) (Nishimura-Tadaki *et al.*, 2011) and Nuclear RNA export factor 5 (*NXF5*, Xq22.1) (Bertini *et al.*, 2010). Some will individually be discussed below.

Autosomal rearrangements

Autosomal translocations—Robertsonian and reciprocal—have been observed in sporadic cases in Belgian, American, Japanese and Chinese women (Hens *et al.*, 1989; Orczyk *et al.*, 1989; Kawano *et al.*, 1998; Jiao *et al.*, 2012). Perturbations presumably confer haploinsufficiency or interrupt pivotal genes in these regions. Non-specific defective meiotic pairing or a position effect on contiguous genes is also a potential explanation (Simpson, 2008; Persani *et al.*, 2009). No autosomal region appears preferentially involved, long frustrating investigators seeking to use breakpoints to localize regions containing autosomal genes of relevance. Searches for autosomal regions disrupted in X-autosome translocations have similarly not proved fruitful in identifying autosomal roles in POI.

Single genes causing non-syndromic POI

Aside from regions of interest defined by chromosome deletions and translocations, the other traditional strategy to identify candidate genes in POI is to study genes whose product is known and plays a role in human folliculogenesis or shows an organ-specific effect based on murine knockout models (candidate genes). Many genes have been interrogated for these reasons. In this section we specifically review all genes for which data warrant strong consideration as a candidate gene for POI. Table II and Supplementary Table SII contain available details of reported studies.

Variants occurring in evolutionary conserved regions are more likely to carry functional significance. These include missense, nonsense, insertion or deletion variants and were considered as plausible causative variants with clinical significance. Thus, perturbations should yield a severe functional defect. A nonsense mutation that results in truncated protein should lead to haploinsufficiency; a splicing site mutation or insertion/deletion should result in a frameshift that leads to a different protein product; a missense mutation may change the amino acid and cause a dominant negative effect.

Those regions or genes found only by genome-wide studies, including genome-wide association study, cytogenomic study, whole-exome sequencing, and next generation sequencing (NGS), are cited separately.

Genes on the X chromosome

Bone morphogenetic protein 15 (BMP15) (Xp11.2). *BMP15* is located on chromosome Xp11.2. The possible involvement of *BMP15* in POI pathogenesis was initially supported by evidence from animal models. Inverdale and Hanna sheep with a naturally occurring *Bmp15* mutation had increased ovulation rate and twin and triplet births in heterozygotes, but ovarian failure results from impaired follicular development beyond the primary stage in homozygotes (Galloway et al., 2000). *Bmp15* knockout female mice also were subfertile, showing decreased ovulation rates, reduced litter size and decreased number of litters per lifetime (Yan et al., 2001).

In humans *BMP15* was first implicated in POI by Di Pasquale et al. (2004), who reported a heterozygous p.Y235C missense mutation in each of two sisters having ovarian failure. The authors presented *in vitro* evidence for a dominant negative mechanism. Other variants have been identified in Caucasian, Indian and Chinese women with POI, albeit with quite different frequencies (1.5–15%) (Di Pasquale et al., 2006; Dixit et al., 2006; Laissue et al., 2006; Ledig et al., 2008; Lakhali et al., 2009, 2010b; Rossetti et al., 2009; Wang et al., 2010b). Merely showing different frequencies between a given single nucleotide polymorphism (SNP) in POI and control is a less robust method than finding a unique perturbation with functional validation in a case (Zhang et al., 2007; Ledig et al., 2008). However, some variants found in higher frequency indeed show marked reduction of mature protein production (Rossetti et al., 2009).

Of relevance is that *BMP15* is a member of the transforming growth factor (TGF) family, with dimerization occurring with other TGF proteins such as *GDF9*, to be discussed below. Most reported *BMP15* variants are, in fact, in the region corresponding to the propeptide of the protein, which is essential for dimerization and subsequent post-translational processing into biologically active proteins.

Progesterone receptor membrane component 1 (PGRMC1) (Xq22-q24). *PGRMC1* was first described in 1998 as a putative progesterone-binding

membrane receptor (Losel et al., 2008). This protein is expressed in various tissues, e.g. liver, kidney, adrenal glands, uterus and leukocytes and involves progesterone signaling in the reproductive system (Cahill, 2007; Losel et al., 2008; Mansouri et al., 2008). *PGRMC1* mediates progesterone's anti-apoptotic effects on granulosa cells (Engmann et al., 2006; Peluso et al., 2006; Losel et al., 2008; Mansouri et al., 2008).

Mansouri et al. (2008) identified a mother and daughter with POI, both of whom carried an X;autosome translocation [t(X;11)(q24;q13)]. Systematic mapping of the Xq breakpoint and performing RNA expression studies revealed reduced expression of *PGRMC1*. Mutation screening of 67 females with idiopathic POI identified a third patient having a missense mutation (p.H165R), located in the cytochrome b5 domain. The p.H165R mutation abolishes binding of cytochrome P450 7A1 (CYP7A1) to *PGRMC1* and attenuates *PGRMC1*'s ability to mediate the anti-apoptotic action of progesterone in ovarian cells. These findings suggest that mutant or reduced levels of *PGRMC1* may cause POI through impaired activation of the microsomal cytochrome P450 and increased apoptosis of ovarian cells. A recent study in Chinese patients with POI identified a novel missense mutation (C.556C>T, p. P186S), but there was no functional study to confirm a deleterious effect (Wang et al., 2014b).

Androgen receptor (AR) (Xq12). The *AR* gene encodes the androgen receptor and is involved in sex differentiation and reproduction. Its perturbation in 46,XY individuals results in the well-known sex reversed phenotype of androgen insensitivity, testosterone produced by testis exerting no effect on androgen-dependent differentiation. In the ovary, AR is expressed in developing follicles, mainly granulosa cells. Deficiency of *Ar* in female mice results in a POI-like phenotype and dysregulation of a number of major genes critical for folliculogenesis, indicating that normal folliculogenesis requires AR-mediated androgen action (Shiina et al., 2006). An association between CAG repeat length in exon I of the *AR* gene and POI has been proposed but remains controversial (Bretherick et al., 2008; Chatterjee et al., 2009; Sugawa et al., 2009; Panda et al., 2010). An example is a repeat of two missense mutations (p.T650A and p.O658K) in Indian women with POI (Panda et al., 2010).

Forkhead box O4 (FOXO4) (Xq13.1). The *FOXO4* gene encodes a member of the O class of winged helix/forkhead transcription factor family (FOXO). *FOXO4* is expressed in granulosa cells in mice and human, and is involved in the PI3K (phosphoinositide 3-kinase)/Akt (v-akt murine thymoma viral oncogene homolog 1)/Cdkn1b (cyclin-dependent kinase inhibitor 1B) molecular pathway, which suggests a functional role in ovarian physiology (Pisarska et al., 2009). Mutation screening in 116 Tunisian patients identified only one intronic variant; IVS2 + 41T>G; therefore, *FOXO4* might not be a common cause of POI in the Tunisian population (Fonseca et al., 2012a).

Premature ovarian failure, 1B (POF1B) (Xq21.2). Alluded to previously, this 'gene' is actually a region, but codified by OMIM. Its significance is its location within the critical POI1 region. It was found to be interrupted by a breakpoint in an X-autosome translocation in a patient with secondary amenorrhea (POI). Subsequent mutation analysis in an Italian POI cohort ($N = 223$) only revealed 30 SNPs (Bione et al., 2004). In a Lebanese family having five sisters with POI, Lacombe et al. (2006) established linkage to Xq21 using whole-genome SNP typing and homozygosity-by-descent mapping. Sequencing identified a homozygous p.R329Q

Table II Variants identified in candidate genes on the X chromosome for idiopathic and sporadic POI.

Gene	Location	Cases (N)	Controls (N)	Ethnicity	MR ^a	Sequence variation	Amino acid change	FC	Mechanism	Reference	
BMP15	Xp11.2	50	214	Caucasian	2 (4.0%)	c.242A>G c.595G>A	p.H81R p.G199R			Tiotiu <i>et al.</i> (2010)	
				North Africa							
		100	100	Chinese	1 (1.0%)	c.985C>T	p.R329C	Yes	Slightly affects transactivation of BRE-luc in COV434 granulosa cells	Wang <i>et al.</i> (2010b)	
				Asia							
		300	216	Caucasian	12 (4.0%)	c.13A>C	p.S5R		Markedly reduces mature protein production and affects transactivation of BRE-luc in COV434 granulosa cells	Rossetti <i>et al.</i> (2009)	
		20	93	Germany	None	None	p.R68W			No effect on protein production or transactivation	Ledig <i>et al.</i> (2008)
				China							
		92	76	Chinese	None	None	p.R138H				Zhang <i>et al.</i> (2007)
203	54	Caucasian	3 (1.5%)	c.443T>C	p.L148P				Laissue <i>et al.</i> (2006)		
		African Asian									
133	197	Indian	14 (10.5%)	c.538G>A c.468G>A c.831T>C c.852C>T	p.A180T	Sense	Sense	Sense	Dixit <i>et al.</i> (2006c) and Inagaki and Shimasaki (2010)		
				c.181C>T c.182G>A c.226C>T	p.R61W p.R61E p.R76C			Decreased mature protein production, weaker Smad1/5/8 phosphorylation in COV434 cells and decreased granulosa cell proliferation	Yes		
				c.227G>A c.538G>A c.538G>T/ c.539C>T c.588T>A c.617G>A	p.R76H p.A180T p.A180F/S+V			Decreased mature protein production, weaker Smad1/5/8 phosphorylation in COV434 cells and decreased granulosa cell proliferation	Yes		
				c.631C>T c.661T>C c.727A>G c.381A>G c.*40dupG	p.E211X p.V221R p.L243G				Di Pasquale <i>et al.</i> (2004, 2006)		
166	181	Caucasian	7 (4.2%)	c.202C>T c.538G>A c.704A>G	p.R68W p.A180T p.Y235C			Diminished GC proliferation with a dominant negative effect	Yes		
38	51	New Zealand	None	None					Chand <i>et al.</i> (2006)		
		Japanese									
15	3		None	None					Takebayashi <i>et al.</i> (2000)		

Continued

Table II Continued

Gene	Location	Cases (N)	Controls (N)	Ethnicity	MR ^a	Sequence variation	Amino acid change	FC	Mechanism	Reference
AR	Xq12	133	200	Indian	2 (1.5%)	c.1948A>G c.1972C>A c.1885+9C>A	p.T650A p.O658K Intron			Panda et al. (2010)
FOXO4	Xq13.1	116	143	Tunisian		None				Fonseca et al. (2012a)
POF1B	Xq21.2	223	900	Italian	2 (0.9%)	c.G1477A	p.C444Y			Bione et al. (2004)
DACH2	Xq21.3	257	1110	Italian	2 (0.8%)	c.G274T c.T1111C	p.R37L p.F316S			Bione et al. (2004)
PGRMC1	Xq22-q24	67	153	Swedish Italian	1 (1.5%)	c.494A>G	p.H165R	Yes	Attenuates ability to transduce progesterone's anti-apoptotic action in granulosa cells and abolishes binding capacity to CYP7A1	Mansouri et al. (2008)
		196	200	Chinese	1 (0.51%)	c.556C>T	p.P186S			Wang et al. (2014b)

MR: mutation rate; FC: functional confirmation; BMP15: bone morphogenetic protein 15; AR: androgen receptor; FOXO4: forkhead box O4; POF1B: premature ovarian failure; IB; DACH2: dachshund family transcription factor 2; PGRMC1: progesterone receptor membrane component-1.

^aOnly refer to novel missense, frameshift, and nonsense mutations

mutation, which impaired the capacity to bind nonmuscle actin filaments, and might lead to exaggerated germ-cell apoptosis and POI.

Dachshund family transcription factor 2 (DACH2) (Xq21.3). DACH2, also named dachshund family transcription factor 2, is located on Xq21.3. It was first identified by fine mapping of the disrupted region in an X;auto-some translocation in POI patients (Prueitt et al., 2002). Subsequent mutation screening revealed two novel missense mutations—p.R37L and p.F316S—in an Italian cohort of POI patients (Bione et al., 2004). However, no subsequent evidence of involvement of DACH2 in mammalian gonads or additional mutations in other ethnic population has been reported.

Fragile X mental retardation 1 (FMR1) (Xq27.3). One of the commonest causes of POI is a premutation of FMR1, which when fully perturbed (>200 CGG repeats) causes fragile X syndrome but paradoxically not POI. A prototype of pleiotropic single gene disorders in which POI is one component, FMR1 is discussed in the 'Pleiotropic Single Gene Disorders Having POI' section, along with other pleiotropic genes.

Genes on autosomes

In this section we will review autosomal genes for which data appear to warrant strong consideration as a candidate gene for POI. Supplementary Table SII contains available details on studies generating this conclusion.

Growth differentiation factor 9 (GDF9) (5q31.1). Expressed in oocytes, GDF9 is an attractive candidate gene for POI because it is, like BMP15, a member of the TGF gene family. Increased frequencies of certain novel variants have been detected in European, Caucasian and Asian patients (Dixit et al., 2005; Laissue et al., 2006; Kovanci et al., 2007; Zhao et al., 2007), but not in Japanese and New Zealand populations (Takebayashi et al., 2000; Chand et al., 2006). All variants were heterozygous. Recently, high-resolution array comparative genomic hybridization (CGH) (2.2 kb resolution) was applied in 26 POI Swedish cases, finding one partial GDF9 gene duplication (475 bp) (Norling et al., 2014). Unfortunately, parents were not available to exclude a heritable copy number variant (CNV) less likely to carry significance. This pitfall is discussed further in 'Genome-Wide Studies in POI' section.

Heterozygous changes could result in a dominant negative effect, quite plausible given dimerization with fellow members of the TGF gene family (e.g. BMP15). The proportion of POI due to GDF9 perturbations is, however, unclear. If a hydrophobic amino acid replacing a hydrophilic amino acid were causative, GDF9 perturbations could account for a substantial number (1–4%) of POI cases.

Folliculogenesis specific bHLH transcription factor (FIGLA) (2p13.3). FIGLA, also named factor in the germline, alpha, is a germ-cell specific, basic helix-loop-helix (bHLH) transcription factor, that plays a crucial role in the formation of the primordial follicle and coordinates expression of zona pellucida genes. Zhao et al. (2008) screened 100 Chinese women with POI and identified three variants in four women: missense mutation p.A4E in two women; deletion p.G6fsX66 in one woman, resulting in a frameshift that leads to haploinsufficiency; and deletion p.I40delN in a fourth woman. Functional analyses by the yeast two-hybrid assay demonstrated that the p.I40delN mutation disrupted FIGLA binding to the TCF3 helix-loop-helix (HLH) domain. These findings show that

a subset of Chinese women with sporadic POI harbor mutations in *FIGLA*. Recently, another novel intronic variant was found in 219 Indian POI cases (Tosh *et al.*, 2015). Further functional validation is warranted.

Newborn ovary homeobox gene (*NOBOX*) (7q35). *NOBOX* is an oocyte-specific homeobox gene that plays a critical role in early folliculogenesis. The causative role was discovered by Rajkovic *et al.* (2004). In mice *Nobox* deficiency disrupted early folliculogenesis and oocyte-specific gene expression. Lack of *Nobox* accelerated post-natal oocyte loss and abolished the transition from primordial to growing follicles in mice. In female mice lacking *Nobox*, follicles are replaced by fibrous tissue in a manner similar to non-syndromic ovarian failure in women. Genes preferentially expressed in oocytes, including *Pou5f1* (POU class 5 homeobox 1) and *Gdf9*, are also down-regulated in *Nobox*^{-/-} mice. Lechowska *et al.* (2011) showed that POI in *Nobox* deficient mice results from faulty signaling between somatic and germ line components during embryonic development. In addition, the extremely unusual presence of abnormal adherens junctions between unseparated oocytes within syncytial follicles indicates that faulty communication between somatic and germ cells is involved in, or leads to, abnormalities in the cell adhesion program. Qin *et al.* (2007a) were the first to demonstrate that a perturbation (p.R355H) in *NOBOX* was responsible for human POI. The mutation disrupted *NOBOX* homeodomain binding to *NOBOX* DNA-binding element (NBE) and had a dominant negative effect. Our functional studies demonstrated that haploinsufficiency was involved in the genetic mechanism in humans for POI. Mutations in the homeobox domain of *NOBOX* proved not to be a common explanation for POI in Chinese women (0/200) (Qin *et al.*, 2009) but Bouilly *et al.* (2011, 2015) subsequently reported that novel *NOBOX* loss-of-function mutations accounted for 6.2 and 5.6%, respectively, of cases in two large 'primary ovarian insufficiency' cohorts of Caucasian and African ancestry.

Nuclear receptor subfamily 5, group A, member 1 (*NR5A1*); Steroidogenic factor-1 (*SF-1*) (9q33). *NR5A1* encodes an orphan nuclear receptor that regulates transcription of an array of genes involved in reproduction, steroidogenesis and male sexual differentiation. These include anti-Müllerian hormone (*AMH*), Nuclear receptor subfamily 0, group B, member 1 (*DAX1*), Cytochrome P450, family 11, subfamily A, polypeptide 1 (*CYP11A*), steroidogenic acute regulatory protein (*StAR*), as well as genes encoding steroid hydroxylases, gonadotrophins, and aromatase. Inactivation of *Nr5a1* specifically in mouse granulosa cells causes infertility associated with hypoplastic ovaries.

Philibert *et al.* (2010) identified *NR5A1* mutations as a frequent cause of 'primary amenorrhea' in 46,XY phenotypic female adolescents with a low testosterone concentration. Lourenco *et al.* (2009) sequenced *NR5A1* in four families (each having at least one family member with a 46,XY disorder of sex development and another with 46,XX POI) and 25 subjects with sporadic POI, and they identified 19 different mutations in the *NR5A1* gene. Functional studies indicated that these mutations substantially impaired the transactivational activity of *NR5A1*. Subsequently additional mutations were identified in different ethnicities with low frequencies (Supplementary Table SII). Janse *et al.* (2012) sequenced the coding regions of *NR5A1* in a large, well-phenotyped cohort of 356 Dutch women with POI, finding 9 different mutations in 10 patients. Functional prediction showed low to intermediate pathogenicity for all non-conserved mutations. However, the novel p.Y5D mutation,

detected in a non-domain region, was presumed to result in haploinsufficiency in Chinese patients with POI (Jiao *et al.*, 2013).

FSH receptor (FSHR) (2p21-p16). *FSH/FSHR* signaling plays a key role in normal gonadal function by regulating follicular growth, estrogen production and oocyte maturation. Mutation in *FSHR* was the first autosomal molecular explanation for POI, elucidated prior to the contemporary era. Aittomäki (1994) and Aittomäki *et al.* (1995, 1996) ascertained 75 primary or secondary amenorrhea cases, and found homozygous mutations (c.566C>T, p.A189V), in the extracellular portion of this G-protein receptor, in women of six Finnish families with hypergonadotrophic ovarian dysgenesis. This mutation resulted in a dramatic reduction of binding capacity and signal transduction, but with apparently normal ligand-binding affinity (Aittomäki *et al.*, 1995). The frequency of the c.566C>T mutation is 0.96% in a Finnish population (Jiang *et al.*, 1998). However, subsequent screening in cohorts of different ethnicities seldom found mutations (da Fonte Kohek *et al.*, 1998; Jiang *et al.*, 1998; Conway *et al.*, 1999; Takakura *et al.*, 2001; Tong *et al.*, 2001; Sundblad *et al.*, 2004; Chen *et al.*, 2006; Ledig *et al.*, 2008; Vilodre *et al.*, 2008; Prakash *et al.*, 2009; Woad *et al.*, 2013). Therefore, *FSHR* mutations are not uncommon in XX gonadal dysgenesis in Finland, but apparently rare elsewhere (Supplementary Table SII).

TGF, beta receptor III (TGFB3) (1p33-p32). Human *TGFB3* is located at 1p33-p32 and encodes the TGF-beta type III receptor. The encoded receptor is a membrane proteoglycan that often functions as a co-receptor with other TGF-beta receptor superfamily members. Two missense variants, p.E459G and p.P825L, were identified in Chinese women with idiopathic POI, both predicted to have functional and structural impacts on the *TGFB3* protein (Qin *et al.*, 2011). Another missense mutation—p.P775S—was found in an Indian POI case (Dixit *et al.*, 2006b).

G protein-coupled receptor 3 (GPR3) (1p36.1-p35). The *GPR3* gene, located in 1p36.1-p35 and having 2 exons, is a member of the G protein-coupled receptor family. Predominantly expressed in oocytes, *GPR3* maintains meiotic arrest in antral follicles until the LH surge through pathways involved in cAMP and cGMP regulation. In *Gpr3*^{-/-} mice, the majority of oocytes in antral follicles had unscheduled premature resumption of meiosis (Mehlmann *et al.*, 2004). A synonymous variant (c.135G>A, p.V45V) was found in one Chinese patient (Zhou *et al.*, 2010), and another study also failed to find any potential disease-associated changes in 82 North American Caucasian women with POI (Kovanci *et al.*, 2008).

Wingless-type MMTV integration site family, member 4 (WNT4) (1p36.23-p35.1). *WNT4* encodes a secreted extracellular signaling protein that is expressed in human ovaries early in fetal development (Jaaskelainen *et al.*, 2010), and plays a critical role in female sex determination and differentiation. In the ovaries of *Wnt4* mutant mice, the rate of apoptosis was similar to that of wild type mice at birth; however, apoptotic cells progressively increased during follicular development (Jaaskelainen *et al.*, 2010). By sequencing the coding region of *WNT4* in 55 Tunisian women with POI, a synonymous variant in exon 2 (c.99G>A, p.S33S) was identified (Lakhal *et al.*, 2012). Mutational analysis was also performed in 145 Chinese women with POI with no causative variants found (Chen *et al.*, 2011b).

Inhibins: inhibin, alpha (INHA) (2q35); inhibin, beta A (INHBA) (7p15-p13); inhibin, beta B (INHBB) (2cen-q13). Inhibin is a dimeric glycoprotein hormone. Belonging, like BMP-15 and GDF9, to the superfamily of TGF- β , inhibin is a negative regulator of FSH. Inhibin encompasses inhibin, alpha (INHA) (2q35), inhibin, beta A (INHBA) (7p15-p13), and inhibin, beta B (INHBB) (2cen-q13).

The missense mutation c.769G>A (p.A257T) in the *INHA* gene was more frequently found in patients with POI in New Zealand (7%) (Shelling et al., 2000), India (11.2%) (Dixit et al., 2004) and Italy (4.5%) (Marozzi et al., 2002). Increased susceptibility to POI was associated with impaired inhibin B bioactivity (Chand et al., 2007). The additional novel missense mutations c.275G>A (p.S92N), c.525C>G (p.H175Q) and c.545C>A (p.A182D) were exclusively identified in Indian POI patients (1.25% for each mutation) (Dixit et al., 2006a). It is unclear whether polymorphisms in the *INHA* promoter result in reduced inhibin expression, but the promoter variant c.-16C>T was significantly under-represented in patients with POI in New Zealand (Harris et al., 2005). The *INHBA* and *INHBB* gene encode inhibin β A and inhibin β B subunits. Causative mutations were not found in the two genes, however, except possibly for one synonymous mutation c.1032C>T in the *INHBA* gene (Shelling et al., 2000; Chand et al., 2010).

POU class 5 homeobox 1 (POU5F1) (6p21.31). The *POU5F1* transcription factor gene, located on 6p21.31, is significantly down-regulated in *Nobox* knockout mice. Thus, *POU5F1* becomes a potential candidate gene for POI, a downstream target of *NOBOX*. Wang et al. (2011) sequenced 175 Chinese POI cases and found one non-synonymous variant (p.P13T), a heterozygous hydrophobic to hydrophilic substitution.

MutS homolog 4 (MSH4) (1p31) and MSH5 (6p21.3). *MSH4* and *MSH5* belong to the DNA mismatch repair gene family, playing pivotal roles in meiotic recombination. Mammalian *MSH4* and *MSH5* proteins form a heterodimeric complex and exert essential functions for normal chromosome synapsis during zygotene. Disruption of *Msh4* or *Msh5* in female mice resulted in sterility, degenerated ovaries and progressive loss of oocytes due to meiotic failure (de Vries et al., 1999; Kneitz et al., 2000). In a case-control study in a Caucasian population, in which both genes were sequenced, a heterozygous mutation p.P29S in *MSH5* was found in 2 of 41 cases. The mutation was located in the Hmsh4-binding domain of *MSH5* which could disrupt the integrity of the protein interaction between *MSH5* and *MSH4* (Mandon-Pepin et al., 2008).

Forkhead box O3 (FOXO3) (6q21). Forkhead transcription factor *FOXO3*, located at 6q21, encodes a master regulator and potent suppressor of primordial follicle activation. Loss of *Foxo3* function in mice leads to POI due to global follicle activation (Liu et al., 2007). The frequency of variants in POI patients differs in different ethnic groups (Supplementary Table SII), but in several populations the frequencies of *FOXO3* variants are not insignificant (6% in French and 13.3% in Chinese). However, the pathological role of these variants needs to be determined by functional studies.

Cbp/p300-interacting transactivator, with Glu/Asp-rich carboxy-terminal domain, 2 (CITED2) (6q23.3). *CITED2* is essential for early embryonic development. This is evidenced by delayed differentiation of gonads in *Cited2*^{-/-} mice (Combes et al., 2010). Fonseca and colleagues reported

a novel missense mutation p.P202T in one of 116 Tunisian POI cases (Fonseca et al., 2012b). Further studies in other populations are warranted.

Spermatogenesis and oogenesis specific basic helix-loop-helix transcription factor 1 (SOHLH1) (9q34.3) and SOHLH2 (13q13.3). As germ cell specific master-master transcription factors, *SOHLH1* and *SOHLH2* orchestrate different oocyte-specific genes essential for early folliculogenesis. *Sohlh1/2*-deficient mice exhibit atrophied ovaries devoid of follicles due to defective primordial-to-primary follicle transition (Pangas et al., 2006; Choi et al., 2008). Novel distinct heterozygous variants were identified in both *SOHLH1* and *SOHLH2* in large cohorts of women with POI of Han Chinese and Serbian origin (Qin et al., 2014a; Zhao et al., 2015). Plausible pathogenesis might involve disturbing the expression, transactivation or homo-/hetero-dimerization of *SOHLH1* or *SOHLH2* proteins. No subsequent reports exist, to our knowledge.

Phosphatase and tensin homolog (PTEN) (10q23.3). Localized on chromosome 10q23.3, *PTEN* plays a causative role in early activation of primordial follicles by negatively regulating the PI3K pathway (Reddy et al., 2008). Primordial follicles become depleted in *Pten* null mice in early adulthood, mimicking the phenotype of POI in humans. However, no causative mutation was detected in coding regions of *PTEN* gene in Japanese and Chinese women with POI (Shimizu et al., 2009; Zhao et al., 2011).

Nanos homolog 1, 2, 3 (Drosophila) (NANOS1, 10q26.11; NANOS2, 19q13.32; NANOS3, 19p13.13). The *NANOS* gene family is known to be required for primordial germ cell (PGC) development and maintenance. Three homologs (*NANOS1*, *NANOS2* and *NANOS3*) exist. Disruption of *Nanos1* in mice did not affect germ cell development, but knockout of *Nanos2* or *Nanos3* resulted in infertility with decreased gonad size due to loss of PGC. *Nanos2* deficiency only resulted in spermatogonia loss whereas *Nanos3* impaired PGC maintenance in both males and females (Tsuda et al., 2003).

Mutations in *NANOS3* were investigated in 80 Chinese and 88 Caucasian women with POI (Qin et al., 2007b). No causative mutations were found in coding exons. However, one potentially relevant heterozygous mutation (c.457C>T; p.R153W) was identified in another study involving 100 Chinese POI patients (Wu et al., 2013). Functional studies showed decreased stability of *NANOS3*, potentially resulting in a hypomorph. And a homozygous mutation (c.358G>A, p.E120K) was found in two sisters with primary amenorrhea from 85 Brazilian women with POI. *In vitro* and *in silico* functional studies revealed that this mutation impaired the ability of *NANOS3* to prevent apoptosis, suggesting a mechanism for POI involving increased PGC apoptosis during embryonic cell migration (Santos et al., 2014). Taken together, these results suggest a role for *NANOS3* mutation in some cases of POI.

Cyclin-dependent kinase inhibitor 1B (CDKN1B) (12p13.1-p12). *CDKN1B*, also known as *P27* and *KIP1*, encodes a cyclin-dependent kinase inhibitor that regulates proliferation and differentiation in many tissues. It suppresses ovarian follicle endowment and activation, and promotes follicle atresia. Premature follicle depletion occurred due to accelerated activation in *Cdkn1b* knockout mice (Rajareddy et al., 2007). Sequence analysis of *CDKN1B* found one novel heterozygous mutation c.356T>C

(p.I119T) in one of 87 Tunisia POI patients (Ojeda *et al.*, 2011). However, no variants were identified in Chinese cohorts (Wang *et al.*, 2010a; Zhao *et al.*, 2013), suggesting that mutations in *CDKN1B* are not common in POI, at least in this population.

Anti-Mullerian hormone receptor, type II (AMHR2) (12q13). *AMHR2* encodes a receptor in the AMH pathway which plays a crucial role in the development and maintenance of reproductive organs in mammals. Polymorphism c.-482 A>G (rs2002555) in *AMHR2* was revealed to be associated with age at menopause in interaction with parity in Dutch women, but no association was found with POI in Korean and Chinese women (Yoon *et al.*, 2013; Qin *et al.*, 2014b). Negative results in *AMHR2* were also reported in 16 Japanese women with POI (Wang *et al.*, 2002). However, Qin *et al.* (2014b) identified two novel missense mutations (p.I209N and p.L354F) in a cohort of Chinese POI women.

KIT ligand (KITLG) (12q22). The human *KITLG* gene, located at 12q22, encodes the ligand of a tyrosine-kinase receptor. KIT/KITLG plays a critical role during oogenesis and folliculogenesis. Mice with a deficiency in *Kitlg* manifested impaired PGCs (Matsui *et al.*, 1991). However, no perturbations were reported in the coding region of *KITLG* from 40 Caucasian POI patients (Hui *et al.*, 2006).

Forkhead box O1 (FOXO1) (13q14.1). *FOXO1*, another member of the forkhead family of transcription factors, is important in granulosa cell function and follicle maturation. Watkins *et al.* (2006) identified one 5'UTR mutation (c.-30C>T) and one missense mutation (p.P84L) in 60 New Zealand and Slovenia POI patients.

Spalt-like transcription factor 4 (SALL4) (20q13.2). *SALL4*, a zinc finger transcription factor, is expressed in murine oocytes. *SALL4* binds to *POU5F1* and could regulate its expression. Both *Sall4* and *Pou5f1* are drastically down-regulated in *Nobox*^{-/-} newborn ovaries (Zhang *et al.*, 2006; Choi *et al.*, 2007). Wang *et al.* (2009) screened the coding regions of *SALL4* in 100 Han Chinese females with non-syndromic POI and identified two heterozygous missense mutations (p.V181M and p.T817A) in the conserved region. These may or may not be POI-associated gene variants. Further studies are needed to determine the functional effect of these variants.

Meiotic protein covalently bound to DSB (SPO11) (20q13.31). *SPO11* is involved in meiosis, forming the double-strand breaks (DSBs) that initiate meiotic recombination. *Spo11*^{-/-} mice are infertile with premature depletion of oocyte because of defective meiosis. However, no novel variants were found in 41 women with non-syndromic POI (Mandon-Pepin *et al.*, 2008). It is not clear whether an association between *SPO11* mutation and sporadic POI exists in human.

DNA meiotic recombinase I (DMC1) (22q13.1). Genes perturbing meiosis are logical candidates for non-syndromic POI. *DMC1* encodes a member of the superfamily of recombinases, which are important for repairing double-strand DNA breaks during mitosis and meiosis. Among 41 French women with POI, Mandon-Pepin *et al.* (2008) found one POI case with homozygous mutation p.M200V. However, a subsequent screening revealed no mutation but two known SNPs in 192 Chinese women with POI (Wang *et al.*, 2012).

Pleiotropic single gene disorders in POI

Distinct from non-syndromic POI, pleiotropic Mendelian disorders may manifest POI as part of their phenotypic spectrum. Indeed, the most common single genetic explanation for POI is represented by such a disorder—premutation for fragile X syndrome (Table III).

Fragile X syndrome: familial mental retardation I (FMR1) (Xq27.3)
Perturbations of *FMR1* are responsible for fragile X syndrome. Clinical features include mental retardation, characteristic facial features with large ears and prominent jaw, connective tissue findings (joint laxity), large testes after puberty, and behavioral abnormalities. Fragile X syndrome occurs in males when CGG repeats number above 200. In females ~70% of women with >200 CGG repeats show intellectual disability (de Vries *et al.*, 1996). The incidence of fragile X syndrome in males is approximately 1 in 4000, and in females 1 in 8000 (ACOG committee opinion, 2006a)

The normal number of CGG repeats in *FMR1* is 32. Thereafter, there is a stage (premutation) in which 54–200 CGG repeats exist. Pathogenic effects including mental retardation and ataxia may exist, as well as POI. About 15–20% of women with a *FMR1* premutation develop POI (Wittenberger *et al.*, 2007). Conversely, 5% of sporadic cases and 10–15% of familial cases in the Caucasian population are explained by *FMR1* premutations. For reasons that are not clear, the number of CGG repeats significantly correlates with risk of POI only within selected ranges. There is only a slightly increased risk of expansion associated with 40–79 repeats; but higher risk with 80–99 repeats, yet no further increased risk occurs after >100 repeats, and as noted POI is not observed with the full mutation (>200 CGG) (Allingham-Hawkins *et al.*, 1999). One possible explanation is that certain genes are suppressed in the 54–100 premutation range and link to POI, whereas other genes become suppressed only with higher numbers of CGG repeats. Perhaps, then, phenotypes in the two groups differ because some ovarian genes are inhibitory and others are the converse. Thus, ovarian function may be initially suppressed but later return to normal function.

Given the comparatively higher frequency of premutation of *FMR1* in POI than the general population (Supplemental Table SIII), *FMR1* testing has become part of the work-up for women with POI. It is formally recommended in Europe (Foresta *et al.*, 2002; European Society of Human Genetics; European Society of Human Reproduction and Embryology, 2006b). In other populations the prevalence is lower. Guo *et al.* (2014) reported that only 2 premutation carriers were found in 379 sporadic Chinese POI cases (0.49%); none were found in 402 controls. The frequency in Chinese women is thus considerably lower than in Caucasian women (3.3–6.7%). Frequencies are 1.56% in Japanese (Ishizuka *et al.*, 2011) and 4.8% in Slovenian women (Gersak *et al.*, 2003).

Blepharophimosis-ptosis-epicanthus syndrome (BPES): forkhead box L2 (FOXL2) (3q23)

Blepharophimosis-ptosis-epicanthus syndrome (BPES) is a pleiotropic autosomal dominant syndrome in which *FOXL2* is perturbed and premature ovarian failure occurs (Crisponi *et al.*, 2001). That *FOXL2* plays a key if not the pivotal role in ovarian development initially came from study of BPES kindreds. More than one hundred unique *FOXL2* mutations have now been described in BPES in different populations (Beysen *et al.*, 2004). By contrast, constitutional mutations are uncommon but

Table III Candidate genes responsible for Mendelian disorders that manifest POI.

Gene	Location	Mendelian syndrome	Somatic features	Reference
<i>FMR1</i>	Xq27.3	Fragile X syndrome	Attention deficits, hyperactivity, social deficits, anxiety disorder, deficits in cognitive flexibility.	Reiss and Hall (2007) and Spath <i>et al.</i> (2010)
<i>FOXL2</i>	3q23	Blepharophimosis-ptosis-epicanthus BPE type I syndrome, BPES I	BPES type I is a complex eyelid malformation associated with POI. The major features of the eyelid malformation involve (i) narrowed horizontal aperture of the eyelids (blepharophimosis), (ii) drooping of the upper eyelid (ptosis), (iii) the presence of a fold of skin arising from the lower eyelid that runs inward and upward (epicanthus inversus), and (iv) lateral displacement of the inner canthi (telecanthus).	Zlotogora <i>et al.</i> (1983) and Oley and Baraitser (1988)
<i>GALT</i>	9p13	Galactosemia	Cataracts, speech defects, poor growth, poor intellectual function, neurologic deficits (predominantly extrapyramidal findings with ataxia).	Schadewaldt <i>et al.</i> (2004)
<i>AIRE</i>	21q22.3	Autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome, APECED	Candidiasis, Addison's disease, hypoparathyroidism, type I diabetes, alopecia, vitiligo, ectodermal dystrophy, celiac disease and other intestinal dysfunctions, chronic atrophic gastritis, chronic active hepatitis, autoimmune thyroid disorders, pernicious anemia.	Fierabracci <i>et al.</i> (2012)
<i>EIF2B</i>	EIF2B2-14q24.3; EIF2B4-2p23.3; EIF2B5-3q27.1	Central nervous system leukodystrophy and ovarian failure, ovarioleukodystrophy	Neurological disorder characterized by involvement of the white matter of the central nervous system. When Leukodystrophies associated with premature ovarian failure referred to as ovarioleukodystrophy.	Mathis <i>et al.</i> (2008)
<i>POLG</i>	15q25	Progressive external ophthalmoplegia, PEO	Manifestations range from involvement limited to the eyelids and extraocular muscles.	Graziewicz <i>et al.</i> (2007)
<i>NOG</i>	17q22	Proximal symphalangism, SYM1	Ankylosis of the proximalinterphalangeal joints.	Kosaki <i>et al.</i> (2004)
<i>PMM2</i>	16p13	PMM2-CDG CDG-1 (a previously known as congenital disorder of glycosylation type 1a)	Cerebellar dysfunction (ataxia, dysarthria, dysmetria), non-progressive cognitive impairment, stroke-like episodes, peripheral neuropathy with or without muscle wasting, absent puberty in females, small testes in males, retinitis pigmentosa, progressive scoliosis with truncal shortening, joint contractures, and premature aging	Sparks and Krasnewich (2005)
<i>HSD17B4</i>	5q21	Perrault syndrome, PS	Sensorineural deafness in both males and females, and neurological manifestations in some patients.	Jenkinson <i>et al.</i> (2013), Morino <i>et al.</i> (2014), Pierce <i>et al.</i> (2011), Pierce <i>et al.</i> (2013) and Pierce <i>et al.</i> (2010)
<i>HARS2</i>	5q31.3			
<i>CLPP</i>	19p13.3			
<i>LARS2</i>	3p21.3			
<i>C10orf2</i>	10q24			
<i>BLM</i>	15q26.1	Bloom syndrome	Chromosomal breakage leading to early onset of aging, short stature and elevated rates of most cancers.	Ellis and German (1996)
<i>ATM</i>	11q22-q23	Ataxia telangiectasia, A-T	Progressive cerebellar degeneration, telangiectasias, immunodeficiency, recurrent infections, insulin-resistant diabetes, premature aging, radiosensitivity, and high risk for epithelial cancers in surviving adults.	Gatti <i>et al.</i> (1991) and Su and Swift (2000)
<i>WRN</i>	8p12	Werner syndrome	Premature aging of the skin, vasculature, and bone and elevated rates of certain cancers, particularly sarcomas.	Epstein <i>et al.</i> (1966)
<i>RECQL4</i>	8q24.3	Rothmund–Thomson syndrome, RTS	Cutaneous rash, sparse hair, small stature, skeletal and dental abnormalities, cataracts, premature aging, and an increased risk for cancer, especially malignancies originating from bone and skin tissue.	Wang <i>et al.</i> (2001)

FMR1: Fragile X mental retardation 1; *FOXL2*: forkhead box L2; *GALT*: galactose 1-phosphate uridyl transferase; *AIRE*: autoimmune regulator; *EIF2B*: eukaryotic translation initiation factor; *POLG*: polymerase (DNA directed), gamma; *NOG*: noggin; *PMM2*: Phosphomannomutase 2; *HSD17B4*: Hydroxysteroid (17-beta) dehydrogenase 4; *HARS2*: Histidyl-tRNA synthetase 2, mitochondrial; *CLPP*: caseinolytic mitochondrial matrix peptidase proteolytic subunit; *LARS2*: leucyl-tRNA synthetase 2, mitochondrial; *C10orf2*: Chromosome 10 open reading frame 2; *BLM*: Bloom syndrome, RecQ helicase-like; *ATM*: ATM serine/threonine kinase; *WRN*: Werner syndrome, RecQ helicase-like; *RECQL4*: RecQ protein-like 4.

reported in non-syndromic POI. Perhaps 2–3% of isolated POI cases have a *FOXL2* mutation (De Baere et al., 2002; Harris et al., 2002) (Supplementary Table SII).

Galactosemia: galactose 1-phosphate uridyl transferase (*GALT*) (9p13)

Galactosemia is caused by deficiency of galactose 1-phosphate uridyl transferase (*GALT*). Ovarian failure is a common long-term complication in girls and women with galactosemia, first described by Kaufman and coworkers (Kaufman et al., 1979, 1981), who observed POI in 12 of 18 (67%) galactosemic women. Later Waggoner et al. (1990) reported that 17% (8/47) galactosemic women presented with ovarian failure. However, with regard to duarte galactosemia, a mild variant of *GALT* deficiency, no apparent ovarian dysfunction was reported (Badik et al., 2011). Pathogenesis involves excess galactose toxicity that impairs folliculogenesis, induces resistance to gonadotrophins and accelerates follicular atresia (Fridovich-Keil et al., 2011; Banerjee et al., 2012).

Carbohydrate-deficient glycoprotein syndrome type I (*CDG-Ia*): phosphomannomutase 2 (*PMM2*) (16p13)

In type I carbohydrate-deficient glycoprotein deficiency, also named phosphomannomutase deficiency, mannose-6-phosphate cannot be converted to mannose-1-phosphate. This lipid linked oligosaccharide is necessary for formation of secretory glycoproteins. Neurologic abnormalities and ovarian failure occur (de Zegher and Jaeken, 1995; Kristianson et al., 1995). Located on 16p13, *PMM2* is typically caused by a missense mutation (Bjursell et al., 1997).

Proximal symphalangism (*SYM1*) and multiple synostoses syndrome (*SYNS1*): *noggin* (*NOG*) (17q22)

NOG encodes a secreted polypeptide that binds to and inactivates members of the TGF- β superfamily (i.e. BMP2, 4, 7, 14 and GDF5). *NOG* is expressed in various tissues including female reproductive organs. *NOG* mutations are known to explain proximal symphalangism (*SYM1*) and multiple synostoses syndrome (*SYNS1*) (Gong et al., 1999). In addition, Kosaki et al. (2004) described a heterozygous mutation (p.E48K) in *NOG* in a female presenting with *SYM1* and also having POI. However, Laissue et al. (2007) concluded the relationship between *NOG* mutations and non-syndromic POI was not clear, having screened the coding sequence of *NOG* in 100 non-syndromic sporadic POI patients and identifying only one heterozygous mutation (p.G92E) (Supplementary Table SII). Actually this experience mirrors the situation involving other pleiotropic genes causing POI.

Autoimmune regulation/autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (*APECED*): autoimmune regulator (*AIRE*) (21q22.3)

The *AIRE* gene can if perturbed lead to multi-system abnormalities: alopecia, vitiligo, keratopathy, malabsorption, hepatitis and mucocutaneous candidiasis. Ovarian hypoplasia often occurs, usually in the third decade (Wang et al., 1998). Many different *AIRE* perturbations have been found in this autosomal dominant disorder, not only nonsense mutations but also frame shifts. No particular mutation leads to ovarian failure as distinct from other autoimmune phenomena. *AIRE* mutations have not yet been sought in women with isolated POI (non-syndromic).

Ovarian leukodystrophy: eukaryotic translation initiation factor (*EIF2B*): subunit 2 beta, 39 kDa (*EIF2B2*) (14q24.3); subunit 4 delta, 67 kDa (*EIF2B4*) (2p23.3); subunit 5 epsilon, 82 kDa (*EIF2B5*) (3q27.1)

'Vanishing white matter' leads to variable but progressive neurological degeneration. Ovarian failure may coexist (Schiffmann et al., 1997; Boltshauser et al., 2002). That the causative gene (*EIF2B*) allows denatured stress-related proteins to accumulate makes plausible the relevance to oogenesis, given ongoing oocyte degeneration. Fogli et al. (2003) found variants in *EIF2B2*, *EIF2B4* and *EIF2B5* in seven POI cases with neurologic abnormalities, but in 93 non-syndromic POI none were found positive (Fogli et al., 2004). Therefore, *EIF2B* genes have not yet been shown to be responsible for non-syndromic POI.

Perrault syndrome: hydroxysteroid (17-beta) dehydrogenase 4, *HSD17B4*, 5q21; histidyl-tRNA synthetase 2, mitochondrial, *HARS2*, 5q31.3; leucyl-tRNA synthetase 2, mitochondrial, *LARS2*, 3p21.3; caseinolytic mitochondrial matrix peptidase proteolytic subunit, *CLPP*, 19p13.3; chromosome 10 open reading frame 2, *C10orf2*, 10q24

Perrault syndrome is a well-recognized pleiotropic autosomal recessive disorder characterized by ovarian failure in females, progressive sensorineural deafness in both males and females, and in some patients, neurological manifestations. Only recently have the underlying genes been identified and proved to be heterogeneous. In a family of mixed European ancestry with two sisters presenting with Perrault syndrome, compound heterozygous variants - c.650A>G (p.Y217C) and c.1704T>A (p.Y568X), in *HSD17B4* were found. Then with similar genomic strategies with linkage analysis or whole-exome sequencing (WES), mutations in *HARS2*, *LARS2*, *CLPP* and *C10orf2* have been found in the context of Perrault syndrome (Pierce et al., 2011, 2013; Jenkinson et al., 2013; Morino et al., 2014). All these genes are essential for normal mitochondrial function. This group of causative genes is detailed in 'Mitochondrial Genes Causing POI' Section.

Other pleiotropic disorders

Table III lists other nuclear genes causing syndromes in which POI is a feature. These include *POLG* associated with progressive external ophthalmoplegia (PEO, detailed in 'Mitochondrial Genes Causing POI' section), *BLM* with Bloom syndrome, *AIRE* with Ataxia telangiectasia, *WRN* with Werner syndrome and *RECQL4* with Rothmund–Thomson syndrome. A common feature of several is chromosomal breakage, best exemplified by Bloom syndrome (Simpson and Elias, 2003; Simpson, 2014).

In aggregate these and other conditions in Table III explain no more than 1% of cases of POI. From a scientific perspective however, elucidating the role these genes play in reproduction offers novel clues to integrity needed for normal ovarian differentiation. Clinicians caring for females with these syndromes should thoroughly investigate and evaluate any reported menstrual irregularities.

Mitochondrial genes causing POI

Perturbations of mitochondrial genes or nuclear genes affecting mitochondria are good candidates for POI because the mature oocyte has the greatest number of mitochondria of any human cell. Mature oocytes readily accumulate mitochondria during oogenesis, mitochondrial biogenesis

playing an essential role in oocyte maturation, fertilization and embryo development. Dysregulation of mitochondrial dynamics contributes to excess oxidative stress and initiation of apoptosis, thus accelerating follicle depletion. A marked quantitative decrease of mitochondrial DNA (mtDNA) in oocytes and peripheral blood cells has been well documented in women with ovarian insufficiency (May-Panloup et al., 2005; Bonomi et al., 2012). Thus, any gene affecting mitochondria involving muscular and neurological disturbance is a candidate, because these systems are so dependent on mitochondrial integrity. Genes governing mitochondrial functions may be located in the nucleus, like those discussed in 'Single Genes Causing Non-syndromic POI' section, or in mitochondria itself (mtDNA). To date, those of relevance to POI have been nuclear genes.

Progressive external ophthalmoplegia (PEO): polymerase (DNA directed), gamma (POLG) (15q25)

mtDNA mutations usually affect muscular and neurological systems. In progressive external ophthalmoplegia (PEO) clinical features are proximal myopathy, sensory ataxia and parkinsonism. The causative mutation lies in the gene encoding polymerase gamma (POLG), which is responsible for mtDNA replication and repair. In three of seven families studied by Luoma et al. (2004), POI cosegregated with PEO. A p.Y955C mutation in POLG was found in two of these three families. p.Y955C (tyrosine to cytosine) affects a highly conserved region, rendering functional plausibility. Compound heterozygosity (N468D/A1105T) was observed by Pagnamenta et al. (2006) in another 3-generation family in which multiple family members were affected with both PEO and POI. Mutation screening of POLG in isolated POI, has, however, yielded few perturbations. A single p.R953C mutation was found in 201 POI cases (0.5%) cumulatively reported by Tong et al. (2010). No novel mutations were identified in Italian and British women with POI (Pagnamenta et al., 2006; Bonomi et al., 2012; Duncan et al., 2012) (Supplementary Table SII). Therefore, POLG variation is not a common explanation for sporadic POI in the absence of clinical suspicion for other mitochondrial-associated physical signs.

Perrault syndrome: HARS2, 5q31.3; LARS2, 3p21.3; CLPP, 19p13.3; C10orf2, 10q24

As reviewed in 'Pleiotropic Single Gene Disorders Having POI' Section, ovarian failure is one of the characterized features in Perrault syndrome in females. The genes listed encode mitochondrial tRNA synthetase, chambered protease and primase-helicase and have been found to harbor mutations responsible for POI in Perrault syndrome (Pierce et al., 2011, 2013; Jenkinson et al., 2013; Morino et al., 2014). Through WES in a consanguineous Palestinian family and a nonconsanguineous Slovenian family with Perrault Syndrome, homozygous mutation c.1565C>A (p.T522N) and compound heterozygous mutation c.1077delT and c.1886C>T (p.T629M) in the LARS2 gene were identified, respectively (Pierce et al., 2013). WES was also performed in a nonconsanguineous family of mixed European ancestry, showing compound heterozygous mutations c.598C>G (p.L200V) and c.1102G>T (p.V368L) in the HARS2 gene (Pierce et al., 2011). The homozygous mutations c.433A>C (p.T145P), c.440G>C (p.C147S) and c.270+4A>G in the CLPP gene were observed in three consanguineous Pakistani families with Perrault Syndrome (Jenkinson et al., 2013). In a Japanese family, affected individuals carried compound heterozygous p.R391H and p.N585S in the C10orf2 gene essential for replication of mtDNA; meanwhile compound heterozygous mutations p.W441G and p.V507I were

responsible for this disease in another family of European ancestry (Morino et al., 2014).

In aggregate, the above supports a critical role for genes controlling mitochondria in the maintenance of ovarian function and their roles in non-syndromic POI warrant further investigation.

Multiple malformation syndromes characterized by POI

In other syndromes, POI or primary ovarian failure (gonadal dysgenesis) is accepted as one component but the causative gene has not been found. Supplementary Table SIV lists these conditions. Of particular interest is POI associated with cerebellar ataxia (Simpson, 2013).

Genome-wide studies in POI

Contemporary genetic strategies applied to locate susceptible loci or genes causing POI have extended beyond suspected candidate gene interrogations to genome-wide approaches. Approaches include linkage analysis in families with multiple affected members, CGH for CNV, genome-wide association studies (GWAS), genome-wide sequencing of exomes (WES) and, in the future, whole genome sequencing (WGS).

GWAS

In GWAS, also known as whole genome association study (WGAS), one examines many common genetic variants in different individuals to see if any variant is associated with a trait. In GWAS one searches agnostically through the entire genome to identify variants (SNPs) more common in cases than controls of similar ethnicity. No *a priori* expectations exist. Six GWAS have been performed to identify variants associated with POI, but not all adhered to accepted criteria. Sample sizes were often very small and replicates not sought (Table IV). The first GWAS showed association with the PTH-responsive B1 gene (*PTHBI*) in a small discovery set of 24 women and 24 controls (Kang et al., 2008). Knauff et al. (2009) identified an association of an intron SNP in ADAM metalloproteinase with thrombospondin type I motif, 19 (*ADAMTS19*) with POI in a discovery set of only 99 Dutch women and 181 controls; no replication set existed. Laminin, gamma 1 (*LAMC1*) was then reported to be significantly associated with POI in Korea in 122 cases versus 242 controls (Pyun et al., 2012).

Our group conducted the largest GWAS with an initial discovery set of 391 cases versus 895 controls; the independent replication set consisted of 400 cases and 800 controls. The most significant association was at 8q22.3 (1.6×10^{-6} – 3.86×10^{-6}). This falls short of the canonical 10^{-8} expected to confer unequivocal significance. This region does not contain a protein coding candidate gene (Qin et al., 2012b). However, it is now appreciated that 90% of significant GWAS associations are in intragenic or intergenic regions, portions of the 98.5% of the genome not coding for protein. These regions include a host of regulatory genes and networks (ENCODE). 8q22.3 may be an important yet undefined long-distance regulatory region affecting ovarian differentiation and oogenesis. Disruption might lead to ovarian failure, analogous to disruption of the region upstream of Sex determining region Y-box 9 (*SOX9*) that causes XY sex reversal in mice and humans (Qin et al., 2003, 2004). Replication in independent cohorts needs to be performed to determine potential causative roles.

GWAS have revealed multiple loci potentially associated with POI in Chinese, Korean, and Dutch women. However, in each it was difficult

Table IV Genome-wide association studies for POI.

	Kang et al. (2008)	Knauff et al. (2009)	Qin et al. (2012b)	Pyun et al. (2012)	Oldenburg et al. (2008)	Caburet et al. (2012)
Ethnicity	Korean	Caucasian (Dutch)	Chinese	Korean	Dutch	Middle-Eastern
Discovery set						
No. of cases	24	99	391	24	10	5
No. of controls	24	181	895	24	5	4
Associations in discovery set	<i>PTHB1</i> at 7p14 showed strongest association. Ht1 GAAAG: POI-susceptible haplotype; Ht2 TGTGC: POI-resistant haplotype.	rs246246 mapped to <i>ADAMTS19</i> intron	8q22.3 (10^{-6})	22 SNPs in <i>LAMC1</i> associated with POI	Susceptible locus: 5q14.1-q15	Susceptible loci: 7p21.1-15.3, 7q21.3-22.2
Replication set						
No. of cases	101	60	400	98	–	–
No. of controls	87	90	800	218	–	–
Results of replication set	<i>PTHB1</i> associated with POI; Ht1 confers susceptibility to POI.	Association not confirmed.		Frequencies of 9 SNPs and 1 haplotype were higher in POI than in control.	–	Sequencing three candidate genes <i>DLX5</i> , <i>DLX6</i> and <i>DSS1</i> did not reveal causal mutations

PTHB1: Bardet–Biedl syndrome 9 (BBS9); *ADAMTS19*: ADAM metalloproteinase with thrombospondin type 1 motif, 19; *LAMC1*: laminin, gamma 1; *DLX5*, 6: distal-less homeobox 5 and 6; *DSS1*: split hand/foot malformation (ectrodactyly) type 1.

to implicate specific novel genes, and in none did significance exceed 10^{-6} ; positive findings were not always replicated. The most likely explanation for these data is limitation based on small sample size. In no study did sample number exceed 1000, thus lacking statistical power sufficient to detect a modest association when evaluating hundreds of thousands of SNPs. That POI is a rarer condition than polycystic ovary syndrome or endometriosis makes it difficult to accrue the requisite sample size possible in studying those disorders (Uno et al., 2010; Chen et al., 2011a; Shi et al., 2012).

GWAS involving family linkage analysis

Table IV includes two genome-wide association linkage studies involving single extended families. In a Dutch POI family subjected to a genome-wide linkage scan 5q14.1-q15 was found to be a susceptibility region (Oldenburg et al., 2008). Caburet et al. (2012) performed genome-wide linkage and homozygosity analysis in a large consanguineous Middle-Eastern POI-affected family showing autosomal inheritance. Two regions—7p21.1-p15.3 and 7q21.3-q22.2—were identified as candidate regions by homozygosity mapping. However, sequencing the three most plausible candidate genes in this region—*DLX5*, *DLX6* and *DSS1*—failed to reveal mutations.

GWAS involving age of menopause

Insights derived from shared genetic susceptibility between POI and either age at natural menopause (AANM) or early menopause (EM) represent another potential path identifying novel entry points for unraveling genetic mechanism involved in POI. Thus, we gathered 36 SNPs shown from previous GWAS studies seeking SNPs associated with AANM or EM, plus 3 additional SNPs in *ESR1* and 2 additional SNPs in *PTHB1*. Differential association was then sought in 371 POI and 800

controls, all of Chinese Han origin (Qin et al., 2012c). Three SNPs, rs2278493 in hexokinase 3 (*HK3*), rs2234693 in estrogen receptor 1 (*ESR1*) and rs12611091 in BR serine/threonine kinase 1 (*BRSKI*), showed nominally significant association with POI. Thus, a plausible relationship could exist between POI and *ESR1*, *BRSKI*, *HK3*.

Cytogenomic studies (CNV)

There is increasing interest toward whole genome studies based on CNVs. Using CGH, one can efficiently search for small duplications or deletions potentially associated with a complex trait (such as POI). The purpose of identifying CNVs is that they may contain or be linked to a causative gene for the disorder studied. Seven array CGH studies have identified CNVs associated with POI (Supplementary Table SV).

In 2009, Aboura et al. (2009) reported that 8 previously known CNVs in 99 POI French patients located near 5 potential candidate genes—Dynein, axonemal, heavy chain 5 (*DNAH5*), NLR family, apoptosis inhibitory protein (*NAIP*), Dual specificity phosphatase 22 (*DUSP22*), Nuclear protein, transcriptional regulator, 1 (*NUPR1*), and *AKT1*. However, this and all other CGH studies have the major pitfall of usually lacking information on parents. A large, *de novo*, CNV deletion is more likely to be pathogenic than an inherited CNV. Another array CGH study involving 74 German patients with POI or ovarian dysgenesis identified 44 losses or gains at autosomes and X chromosome that might explain POI (Ledig et al., 2010). McGuire et al. (2011) identified 17 novel microduplications and 7 novel microdeletions among 89 POI patients, all but one located at autosomes. Included were two novel microdeletions causing haploinsufficiency for Synaptonemal complex central element protein 1 (*SYCE1*) and Cytoplasmic polyadenylation element binding protein 1 (*CPEB1*), genes known to cause ovarian failure in knockout mouse models (Bolcun-Filas et al., 2009; Novoa et al., 2010). Recently,

a high-resolution array CGH identified eleven unique CNVs in 11 patients with POI. Among these CNVs, a tandem duplication of 475 bp containing 3 *NOBOX*-binding elements and an E-box important for *GDF9* gene regulation in the promoter of *GDF9* is likely causative of POI (Norling et al., 2014).

Using a complete X chromosome tiling path array, Quilter et al. (2010) found 15 novel discrete X chromosome intervals in 20/42 (48%) women with POI in the UK. However, in patients from New Zealand, Dudding et al. (2010) detected only two microduplications (Xp22.33 and Xq13.3) in a low frequency of 4%. Knauff et al. (2011) found one CNV in Xq21.3 to be associated with POI, specifically in a region where Protocadherin 11 X-linked (*PCDH11X*) and TGF β -induced factor homeobox 2-like, X-linked (*TGIF2LX*) are located. Interestingly, no deletions were found in other regions, several considered on the basis of traditional cytogenetic studies to be pivotal (Simpson and Rajkovic, 1999).

Differing array CGH platforms and definitions for pathogenicity probably explain some of the discrepant results among the above studies. Also problematic is the limited sample size and lack of parental studies to exclude a CNV ostensibly associated with POI merely having been transmitted from a normal mother.

WES

Low prevalence and impaired fecundity resulting in limited numbers of POI pedigrees without associated somatic anomalies (non-syndromic) has led to increasing application of WES, another agnostic approach. WES has identified several genes in POI not previously anticipated.

In POI associated with somatic features, causative perturbations have been found by WES for *HSB17B4*, *LARS2*, *CLPP* and *C10orf2* in Perrault syndrome (Pierce et al., 2010, 2013; Jenkinson et al., 2013; Morino et al., 2014) (detailed in 'Mitochondrial Genes Causing POI' section and Table V). Up to the present, there have been six WES conducted in non-syndromic POI pedigrees. Interestingly, almost all plausible candidate genes identified were involved in meiosis and DNA repair (Table V).

ATP-dependent DNA helicase homolog (HFM1) (1p22.2). *HFM1*, a meiotic gene encoding DNA helicase preferentially expressed in germ-line tissues such as testis and ovary, is necessary for homologous recombination and synapsis during meiosis (Tanaka et al., 2006). *Hfm1*^{-/-} female mice had significantly reduced ovaries and follicle numbers and an increase in stromal cells (Guiraldelli et al., 2013). The recent report by Wang et al. (2014a) identified a compound heterozygous mutation (c.1686-1G>C and c.2651T>G, p.I884S) in the *HFM1* gene in two affected Chinese sisters. Screening for *HFM1* mutations in a cohort of 69 Chinese women with sporadic POI identified another patient with compound heterozygous mutations (c.2206G>A, p.G736S and c.3929_3930 delinsG, p.P1310R fs*41). These variants were not found in 316 matched controls or databases.

Minichromosome maintenance complex component 8 and 9 (MCM8, 20p12.3; MCM9, 6q22.31). *MCM8* and *MCM9*, recently discovered members of the highly conserved mini-chromosome maintenance proteins (MCM), are genes implicated in homologous recombination and repair of double-stranded DNA breaks. The *MCM8/MCM9* complex is required for the resolution of dsDNA breaks that occur during homologous recombination in pachytene of meiosis I. Failure to resolve breaks predictably leads to oocyte death and small or absent ovaries. *Mcm8* and *Mcm9*-deficient mice are infertile and have small gonads

due to germ cell depletion (Lutzmann et al., 2012). Recently, Rajkovic and colleagues (AlAsiri et al., 2015; Wood-Trageser et al., 2014) discovered homozygous mutations in *MCM8* and *MCM9* genes in consanguineous families with POI (primary amenorrhea), using WES. One of the two mutations found in *MCM9* is c.1732+2T>C, which resulted in abnormal alternative splicing and truncated forms of *MCM9* that are unable to be recruited to sites of DNA damage. The other mutation c.394C>T (p.R132*) results in loss of function of *MCM9*. It is suspected that preferential sensitivity of germline meiosis to *MCM9* functional deficiency and compromised DNA repair in the somatic component most likely account for the ovarian failure and short stature. A homozygous mutant c.446C>G (p.P149R) found in *MCM8* inhibited recruitment of *MCM8* to sites of DNA damage and led to genomic instability. ASNP rs16991615 in *MCM8* was also found associated with the age of natural menopause in the GWAS previously discussed (Chen et al., 2012). The role the novel *MCM8/MCM9* pathway plays in women with idiopathic POI needs to be explored further.

Stromal antigen 3 (STAG3) (7q22.1). *STAG3* encodes a subunit of cohesin, a large protein complex that is essential for proper pairing and segregation of chromosomes during meiosis. A homozygous frameshift mutation resulting from a 1-bp deletion (c.968delC, p.F187fs*7) in *STAG3* was identified in a highly consanguineous Palestinian pedigree; in which a 10-Mb region on 7q21.3–22.2 and a 3-Mb region on 7p21.1–15.3 had been previously identified to show significant linkage with the POI phenotype (Caburet et al., 2012, 2014). Deficiency of *Stag3* in female mice results in severe and early ovarian dysgenesis, with distinctive lack of oocytes and ovarian follicles. Early meiotic arrest and the centromeric chromosomal cohesion defects were observed in *Stag3*^{-/-} fetal oocytes providing further evidence that *Stag3* is essential for assembly of the meiotic cohesin ring and the synaptonemal complex (Caburet et al., 2014).

Synaptonemal complex central element 1 (SYCE1) (10q26.3). The *SYCE1* gene encodes a component of the synaptonemal complex where paired chromosome homologs closely associate in meiosis (synapsis) before crossover. *Syce1*^{-/-} mice were infertile with smaller gonads, and showed loss of follicles in ovaries and postmeiotic cells in testis. Early meiosis arrest at prophase I followed by cell apoptosis could explain the phenotype (Bolcun-Filas et al., 2009). Different microdeletions of *SYCE1* have also been reported in Caucasian and Chinese POI patients (McGuire et al., 2011; Zhen et al., 2013). In a consanguineous Israeli Arab family, a homozygous nonsense mutation (c.613C>T, p.Q205*) in *SYCE1* was identified in two sisters with primary amenorrhea, presumably inherited in autosomal recessive fashion (de Vries et al., 2014).

Eukaryotic translation initiation factor 4E nuclear import factor 1 (eIF4ENIF1) (22q11.2). *eIF4ENIF1*, a transport protein, plays an important role in repressing translation through *eIF4E*. Both genes appear to play important roles in ovarian germ cell development (Villaescusa et al., 2006). A heterozygous nonsense mutation (c.1286C>G, p.S429*) was identified in *eIF4ENIF1*, segregating with ovarian insufficiency (menopause age 29–35 years) in a large French Canadian family. No additional mutations were identified in *eIF4ENIF1* or *eIF4E* in 38 unrelated women with isolated POI. Haploinsufficiency or nuclear sequestration might disrupt the development of a normal oocyte complement (de Vries et al., 2014).

Table V Whole-exome sequencing for syndromic or isolated POI.

Gene	Location	Mutation	Function	Compound Het/Homo/Het	Reference
<i>HSB17B4</i>	5q21	c.650A>G (p.Y217C); c.1704T>A (p.Y568X)	Reduced expression with mutant protein	Compound heterozygous	Pierce et al. (2010)
<i>CLPP</i>	19p13.3	c.433A>C (p.T145P), c.440G>C (p.C147S)	Alter the structure of the CLPP barrel chamber that captures unfolded proteins and exposes them to proteolysis	Homozygous	Jenkinson et al. (2013)
<i>LARS2</i>	3p21.3	c.270+4A>G c.1565C>A (p.T522N) c.1077delT; c.1886C>T (p.T629M)	Partially functional in Yeast complementation assay Nonfunctional in Yeast complementation assay	Homozygous Compound heterozygous	Pierce et al. (2013)
<i>C10orf2</i>	10q24	c.1172G>A (p.R391H) c.1754A>G (p.N585S) c.1321T>G (p.W441G) c.1519G>A (p.V507I)	Affect interactions of the linker domain Likely affect enzyme activity Affect interactions of the linker domain	Compound heterozygous Compound heterozygous	Morino et al. (2014)
<i>HFM1</i>	1p22.2	c.1686-1G>C c.2651T>G (p.I884S) c.2206G>A (p.G736S); c.3929_3930 delinsG, (p.P1310R fs*41)		Compound heterozygous Compound heterozygous	Wang et al. (2014a)
<i>MCM9</i>	6q22.31	c.1732+2T>C	Abnormal splicing and truncated protein that are unable to be recruited to sites necessary for DNA damage	Homozygous	Wood-Trageser et al. (2014)
<i>STAG3</i>	7q22.1	c.394C>T (p.R132*) c.968delC (p.F187fs*7)	Repair of chromosome breaks impaired in lymphocytes	Homozygous	Caburet et al. (2014)
<i>SYCE1</i>	10q26.3	c.613C>T (p.Q205*)		Homozygous	de Vries et al. (2014)
<i>MCM8</i>	20p12.3	c.446C>G (p.P149R)	Impedes the repair of MMC-induced chromosomal breaks; inhibits MCM8 recruitment to sites of DNA damage; impairs DNA binding ability at the N-terminus	Homozygous	AlAsiri et al. (2015)
<i>eIF4ENIF1</i>	22q11.2	c.1286C>G (p.S429*)		Heterozygous	Kasipillai et al. (2013)

HSB17B4: Hydroxysteroid (17-beta) dehydrogenase 4; CLPP: Caseinolytic mitochondrial matrix peptidase proteolytic subunit; LARS2: leucyl-tRNA synthetase 2, mitochondrial; C10orf2: Chromosome 10 open reading frame 2; HFM1: ATP-dependent DNA helicase homolog; MCM8, MCM9: minichromosome maintenance complex component 8 and 9; STAG3: stromal antigen 3; SYCE1: synaptonemal complex central element; eIF4ENIF1: eukaryotic translation initiation factor 4E nuclear import factor 1.

The whole exome studies in familial POI mentioned above mainly involve genes crucial during meiosis, such as generating and repairing DSBs, chromosome synapsis and recombination, and sister chromatid cohesion. This implies that perturbation of genes encoding proteins that regulate meiosis can result in autosomal recessive primary ovarian insufficiency in humans. Despite some negative candidate gene results with meiotic genes (*DMC1*, *SPO11* and *MSH4*) in POI, WES results still provide reasons to pursue other genes participating in meiosis and DNA repair in ubiquitous pathways.

NGS

Future studies on targeted candidate genes of POI can be anticipated using NGS. A recent abstract (Fonseca et al., 2015) reported a study of 12 POI cases having non-syndromic POI and 176 control women whose menopause had occurred after age 50 years. A further 345 women from the same ethnic origin were stated to be recruited to assess allele frequency for potentially deleterious sequence variants. In the 12 POF cases, complete coding regions of 70 candidate genes were fully sequenced with mutations claimed in *ADAMTS19*, *BMPR2* and *LHCGR* genes. Full details are awaited.

Regulatory genes and networks

Seeking POF genes has to date largely focused on coding variants, presuming plausible protein disruption. However, only 1.5% of the genome is protein-coding. Indeed, many POF-associated variants in whole genome studies map within, or in linkage disequilibrium to, intronic or intergenic regions; thus, these regions likely contain causative regulatory genes or networks. In our own POI GWAS, for example, 'gene desert' 8q22.3 was the region of most significance by association (Qin et al., 2012b).

Non-coding variants must be more robustly interrogated. Perturbations sought should include involving non-coding RNA (microRNA [miRNA], long non-coding RNA [lncRNA]), disruption or creation of alternative splicing or transcription factor-binding sites, and epigenetic modifications (DNA methylation patterns, chromatin modification). Large epigenetic consortia, such as ENCODE (<http://genome.ucsc.edu/ENCODE/>), Roadmap (<http://www.roadmappigenomics.org/>) and iHEC (<http://www.ihec-epigenomes.org/>) point to approaches used to characterize the regulatory landscape of susceptibility regions for specific cell types. Unfortunately, ENCODE (Birney et al., 2007; ENCODE Project Consortium, 2012; Stamatoyanopoulos et al.,




Figure 1 Schematic representation of chromosomal location of plausible genes associated with primary ovarian insufficiency.

2012) bypassed the reproductive track (males and females), and thus an ovarian ENCODE does not yet exist.

In order to generate the reproductive ENCODE, human ovarian tissue of specific cell types must be studied. Then, differences in regulatory genes between affected and normal individuals can be determined. Massive parallel sequencing using RNA-seq, or Methyl-seq will facilitate a systematic study of the transcriptome in ovaries in relation to genotypes and variants in POI. Chromatin immunoprecipitation followed by sequencing (ChIP-Seq) and RNA-Seq can also help identify novel transcriptional and epigenetic regions and, hence, mechanisms that potentially underlie POI. To analyze the complete proteomes and quantification of post-translational modifications, mass spectrometry would be the most attractive approach.

Having accumulated the requisite information, network and pathway-based analyses of assembled datasets can be assembled using systems biology approaches. Cascades of transcriptional regulation can be evaluated simultaneously, identifying overlap among genes or putative networks that currently constrain key variables in model systems.

The ultimate proof that an identified variant has a pathological effect traditionally relies on constitutive or conditional knock in/out or cell lines with a mutation signature. Increasingly, other approaches will be utilized. Patient-specific induced pluripotent stem cells (iPSCs) lines might offer an individually targeted genetic model system to identify and manipulate pathologic pathway. Use of iPSCs should be of particular value in assessing regulatory genes, which may have a quantitative deleterious effect rather than 'all or none' qualitative effect, observed or assumed, in missense or nonsense protein-coding mutations.

Conclusion: current status of genes causing POI

The POI—causative genes surveyed and found to date allow us to reach several conclusions.

First, many genes have emerged as POI candidates (Fig. 1), but in non-syndromic POI only a minority have been proven equivocally causative by functional validation. These include *BMP15*, *PGRMC1*, and *FMR1* pre-mutation on the X chromosome and on autosomes *GDF9*, *FIGLA*, *FSHR*, *NOBOX*, *NRSA1*, *NANOS3*, *STAG3*, *SYCE1*, *MCM8/9* and *HFM1*. No perturbations have been found in a dozen other plausible candidates for which murine knockouts show ovarian failure, but this may simply reflect small sample sizes or interrogation restricted to a single ethnic group.

Second, notable differences in frequency exist among different populations. This is predictable for any genetic condition, and in POI this has already been observed in *FSHR*, *BMP15*, *NOBOX*, *FOXL2*, *TGFB3*, *CDKN1B* and *FOXO3A* (Supplementary Table SII). Future genetic studies should involve different ethnic groups and larger sample sizes. Clinically, caution should apply when counseling on the basis of data derived from an ethnic group different from that of the counseled individual.

Third, causative POI genes are increasingly being shown not only to be restricted to expression in ovaries but also expressed ubiquitously. True, *POF5* (*NOBOX*) and *POF6* (*FIGLA*) are restricted to the ovaries. However, *PTEN* is a major regulator of the PI3K pathway involved in systemic cell proliferation, survival, migration and metabolism. *PTEN* also plays a vital role in the activation of primordial follicles. The spectrum of candidate genes potentially causing POI is being enriched.

Fourth, many genes that currently appear isolated in function actually may be interrelated within yet to be defined pathways. It is logical to stratify by gene function in ostensibly distinct systems: endocrine, folliculogenesis, cell cycle, meiosis, mitochondrial, as examples. More difficult are gene-gene or protein-protein interactions, acting in ways not yet evident. This review thus unavoidably overestimates the role that protein-coding genes play to the detriment of upstream and downstream regulatory genes. Until recently, exome studies had to be restricted to sequencing individual candidate genes. Continued advances in sequencing techniques and contemporary bioinformatics will facilitate finding additional genes responsible for POI in other portions of the genome.

Fifth, elucidating the etiology and molecular basis of POI is of paramount importance not only in understanding ovarian physiology but also in providing genetic counseling and fertility guidance. Once additional variants are detected, it will be increasingly possible to predict the age of menopause. Women having certain perturbations of POI can be offered the option of oocyte cryopreservation, with later thawing and use in assisted reproductive technology at the appropriate age.

Supplementary data

Supplementary data are available at <http://humupd.oxfordjournals.org/>.

Authors' roles

Y.Q. contributed to literature searching and drafting the article. X.J. participated in literature searching and creating figures and tables. J.L.S. contributed the study design and revised the article critically. Z.-J.C. supervised the study design and revision.

Funding

This work was supported by the National Basic Research Program of China (973 program-2012CB944700); the National Natural Science Foundation of China (81270662, 81471509, 81370692, 81370687); the Key Program of National Natural Science Foundation of China (81430029). Funding to pay the Open Access publication charges for this article was provided by National Basic Research Program of China (973 program-2012CB944700).

Conflict of interest

None declared.

References

- Aboura A, Dupas C, Tachdjian G, Portnoi MF, Bourcigaux N, Dewailly D, Frydman R, Fauser B, Ronci-Chaix N, Donadille B et al. Array comparative genomic hybridization profiling analysis reveals deoxyribonucleic acid copy number variations associated with premature ovarian failure. *J Clin Endocrinol Metab* 2009; **11**:4540–4546.
- ACOG committee opinion. No. 338: Screening for fragile X syndrome. *Obstet Gynecol* 2006a; **6**:1483–1485.
- Aittomäki K. The genetics of XX gonadal dysgenesis. *Am J Hum Genet* 1994; **5**:844–851.
- Aittomäki K, Lucena JL, Pakarinen P, Sistonen P, Tapanainen J, Gromoll J, Kaskikari R, Sankila EM, Lehvaslaiho H, Engel AR et al. Mutation in the follicle-stimulating hormone receptor gene causes hereditary hypergonadotropic ovarian failure. *Cell* 1995; **6**:959–968.
- Aittomäki K, Herva R, Stenman UH, Juntunen K, Ylostalo P, Hovatta O, de la Chapelle A. Clinical features of primary ovarian failure caused by a point mutation in the follicle-stimulating hormone receptor gene. *J Clin Endocrinol Metab* 1996; **10**:3722–3726.
- AlAsiri S, Basit S, Wood-Trageser MA, Yatsenko SA, Jeffries EP, Surti U, Ketterer DM, Afzal S, Ramzan K, Faiyaz-Ul Haque M et al. Exome sequencing reveals MCM8 mutation underlies ovarian failure and chromosomal instability. *J Clin Invest* 2015; **125**:258–264.
- Allingham-Hawkins DJ, Babul-Hirji R, Chitayat D, Holden JJ, Yang KT, Lee C, Hudson R, Gorwill H, Nolin SL, Glicksman A et al. Fragile X premutation is a significant risk factor for premature ovarian failure: the International Collaborative POF in Fragile X study—preliminary data. *Am J Med Genet* 1999; **4**:322–325.
- Ayed W, Amouri A, Hammami W, Kilani O, Turki Z, Harzallah F, Bouayed-Abdelmoula N, Chemkhi I, Zhioua F, Slama CB. Cytogenetic abnormalities in Tunisian women with premature ovarian failure. *C R Biol* 2014; **12**:691–694.
- Badik JR, Castaneda U, Gleason TJ, Spencer JB, Epstein MP, Ficocioglu C, Fitzgerald K, Fridovich-Keil JL. Ovarian function in Duarte galactosemia. *Fertil Steril* 2011; **2**:469–473.
- Banerjee S, Chakraborty P, Saha P, Bandyopadhyay SA, Banerjee S, Kabir SN. Ovotoxic effects of galactose involve attenuation of follicle-stimulating hormone bioactivity and up-regulation of granulosa cell p53 expression. *PLoS One* 2012; **2**:e30709.
- Baronchelli S, Conconi D, Panzeri E, Bentivegna A, Redaelli S, Lissoni S, Saccheri F, Villa N, Crosti F, Sala E et al. Cytogenetics of premature ovarian failure: an investigation on 269 affected women. *J Biomed Biotechnol* 2011; **3**:70195.
- Bertini V, Ghirri P, Biccocchi MP, Simi P, Valetto A. Molecular cytogenetic definition of a translocation t(X;15) associated with premature ovarian failure. *Fertil Steril* 2010; **3**:1097.
- Beysen D, Vandesompele J, Messiaen L, De Paepe A, De Baere E. The human FOXL2 mutation database. *Hum Mutat* 2004; **3**:189–193.
- Bione S, Sala C, Manzini C, Arrigo G, Zuffardi O, Banfi S, Borsani G, Jonveaux P, Philippe C, Zuccotti M et al. A human homologue of the Drosophila melanogaster diaphanous gene is disrupted in a patient with premature ovarian failure: evidence for conserved function in oogenesis and implications for human sterility. *Am J Hum Genet* 1998; **3**:533–541.
- Bione S, Rizzolio F, Sala C, Ricotti R, Goegan M, Manzini MC, Battaglia R, Marozzi A, Vegetti W, Dalpra L et al. Mutation analysis of two candidate genes for premature ovarian failure, DACH2 and POF1B. *Hum Reprod* 2004; **12**:2759–2766.
- Birney E, Stamatoyannopoulos JA, Dutta A, Guigo R, Gingeras TR, Margulies EH, Weng Z, Snyder M, Dermitzakis ET, Thurman RE et al. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007; **7146**:799–816.
- Bjursell C, Stibler H, Wahlstrom J, Kristiansson B, Skovby F, Stromme P, Blennow G, Martinsson T. Fine mapping of the gene for carbohydrate-deficient glycoprotein syndrome, type I (CDG1): linkage disequilibrium and founder effect in Scandinavian families. *Genomics* 1997; **3**:247–253.
- Bolcun-Filas E, Hall E, Speed R, Taggart M, Grey C, de Massy B, Benavente R, Cooke HJ. Mutation of the mouse *Syce1* gene disrupts synapsis and suggests a link between synaptonemal complex structural components and DNA repair. *PLoS Genet* 2009; **2**:e1000393.
- Boltshauser E, Barth PG, Troost D, Martin E, Stallmach T. 'Vanishing white matter' and ovarian dysgenesis in an infant with cerebro-oculo-facio-skeletal phenotype. *Neuropediatrics* 2002; **2**:57–62.
- Bonomi M, Somigliana E, Cacciatori C, Busnelli M, Rossetti R, Bonetti S, Paffoni A, Mari D, Ragni G, Persani L. Blood cell mitochondrial DNA content and premature ovarian aging. *PLoS One* 2012; **8**:e42423.
- Bouilly J, Bachelot A, Broutin I, Touraine P, Binart N. Novel NOBOX loss-of-function mutations account for 6.2% of cases in a large primary ovarian insufficiency cohort. *Hum Mutat* 2011; **10**:1108–1113.
- Bouilly J, Roucher-Boulez F, Gompel A, Bry-Gauillard H, Azibi K, Beldjord C, Dode C, Bouligand J, Guiochon Mantel A, Hecart AC et al. New NOBOX mutations identified in a large cohort of women with primary ovarian insufficiency decrease KIT-L expression. *J Clin Endocrinol Metab* 2015; **3**:994–1001.
- Bretherick KL, Hanna CW, Currie LM, Fluker MR, Hammond GL, Robinson WP. Estrogen receptor alpha gene polymorphisms are associated with idiopathic premature ovarian failure. *Fertil Steril* 2008; **2**:318–324.
- Caburet S, Zavadakova P, Ben-Neriah Z, Bouhali K, Dipietromaria A, Charon C, Besse C, Laissue P, Chalifa-Caspi V, Christin-Maitre S et al. Genome-wide linkage in a highly consanguineous pedigree reveals two novel Loci on chromosome 7 for non-syndromic familial premature ovarian failure. *PLoS One* 2012; **3**:e33412.
- Caburet S, Arboleda VA, Llano E, Overbeek PA, Barbero JL, Oka K, Harrison W, Vaiman D, Ben-Neriah Z, Garcia-Tunon I et al. Mutant cohesin in premature ovarian failure. *N Engl J Med* 2014; **10**:943–949.
- Cahill MA. Progesterone receptor membrane component 1: an integrative review. *J Steroid Biochem Mol Biol* 2007; **1**–5:16–36.
- Castillo S, Lopez F, Tobella L, Salazar S, Daher V. The cytogenetics of premature ovarian failure. *Rev Chil Obstet Ginecol* 1992; **5**:341–345.
- Ceylaner G, Altinkaya SO, Mollamahmutoglu L, Ceylaner S. Genetic abnormalities in Turkish women with premature ovarian failure. *Int J Gynaecol Obstet* 2010; **2**:122–124.
- Chand AL, Ponnampalam AP, Harris SE, Winship IM, Shelling AN. Mutational analysis of BMP15 and GDF9 as candidate genes for premature ovarian failure. *Fertil Steril* 2006; **4**:1009–1012.

- Chand AL, Ooi GT, Harrison CA, Shelling AN, Robertson DM. Functional analysis of the human inhibin alpha subunit variant A257T and its potential role in premature ovarian failure. *Hum Reprod* 2007;**12**:3241–3248.
- Chand AL, Harrison CA, Shelling AN. Inhibin and premature ovarian failure. *Hum Reprod Update* 2010;**1**:39–50.
- Chatterjee S, Singh R, Kadam S, Maitra A, Thangaraj K, Meherji P, Modi D. Longer CAG repeat length in the androgen receptor gene is associated with premature ovarian failure. *Hum Reprod* 2009;**12**:3230–3235.
- Chen XN, Chen GA, Li MZ. Follicular stimulating hormone receptor gene C566T mutation in premature ovarian failure. *Zhonghua Fu Chan Ke Za Zhi* 2006;**5**:315–318.
- Chen ZJ, Zhao H, He L, Shi Y, Qin Y, Li Z, You L, Zhao J, Liu J, Liang X et al. Genome-wide association study identifies susceptibility loci for polycystic ovary syndrome on chromosome 2p16.3, 2p21 and 9q33.3. *Nat Genet* 2011a;**1**:55–59.
- Chen B, Suo P, Wang B, Wang J, Yang L, Zhou S, Zhu Y, Ma X, Cao Y. Mutation analysis of the WNT4 gene in Han Chinese women with premature ovarian failure. *Reprod Biol Endocrinol* 2011b;**1**:75.
- Chen CT, Fernandez-Rhodes L, Brzyski RG, Carlson CS, Chen Z, Heiss G, North KE, Woods NF, Rajkovic A, Kooperberg C et al. Replication of loci influencing ages at menarche and menopause in Hispanic women: the Women's Health Initiative SHARe Study. *Hum Mol Genet* 2012;**6**:1419–1432.
- Choi Y, Qin Y, Berger MF, Ballow DJ, Bulyk ML, Rajkovic A. Microarray analyses of newborn mouse ovaries lacking Nobox. *Biol Reprod* 2007;**2**:312–319.
- Choi Y, Yuan D, Rajkovic A. Germ cell-specific transcriptional regulator sohlh2 is essential for early mouse folliculogenesis and oocyte-specific gene expression. *Biol Reprod* 2008;**6**:1176–1182.
- Combes AN, Spiller CM, Harley VR, Sinclair AH, Dunwoodie SL, Wilhelm D, Koopman P. Gonadal defects in Cited2-mutant mice indicate a role for SFI in both testis and ovary differentiation. *Int J Dev Biol* 2010;**4**:683–689.
- Conway GS, Conway E, Walker C, Hoppner W, Gromoll J, Simoni M. Mutation screening and isoform prevalence of the follicle stimulating hormone receptor gene in women with premature ovarian failure, resistant ovary syndrome and polycystic ovary syndrome. *Clin Endocrinol* 1999;**1**:97–99.
- Coulam CB, Adamson SC, Annegers JF. Incidence of premature ovarian failure. *Obstet Gynecol* 1986;**4**:604–606.
- Crisponi L, Deiana M, Loi A, Chiappe F, Uda M, Amati P, Biscaglia L, Zelante L, Nagaraja R, Porcu S et al. The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome. *Nat Genet* 2001;**2**:159–166.
- da Fonte Kohek MB, Batista MC, Russell AJ, Vass K, Giacaglia LR, Mendonca BB, Latronico AC. No evidence of the inactivating mutation (C566T) in the follicle-stimulating hormone receptor gene in Brazilian women with premature ovarian failure. *Fertil Steril* 1998;**3**:565–567.
- Davison RM, Quilter CR, Webb J, Murray A, Fisher AM, Valentine A, Serhal P, Conway GS. A familial case of X chromosome deletion ascertained by cytogenetic screening of women with premature ovarian failure. *Hum Reprod* 1998;**11**:3039–3041.
- De Baere E, Lemerrier B, Christin-Maitre S, Durval D, Messiaen L, Fellous M, Veitia R. FOXL2 mutation screening in a large panel of POF patients and XX males. *J Med Genet* 2002;**8**:e43.
- Devi A, Benn PA. X-chromosome abnormalities in women with premature ovarian failure. *J Reprod Med* 1999;**4**:321–324.
- de Vries BB, Wiegers AM, Smits AP, Mohkamsing S, Duivenvoorden HJ, Fryns JP, Curfs LM, Halley DJ, Oostra BA, van den Ouweland AM et al. Mental status of females with an FMR1 gene full mutation. *Am J Hum Genet* 1996;**5**:1025–1032.
- de Vries SS, Baart EB, Dekker M, Siezen A, de Rooij DG, de Boer P, te Riele H. Mouse MutS-like protein Msh5 is required for proper chromosome synapsis in male and female meiosis. *Genes Dev* 1999;**5**:523–531.
- de Vries L, Behar DM, Smirin-Yosef P, Lagovsky I, Tzur S, Basel-Vanagaite L. Exome sequencing reveals SYCE1 mutation associated with autosomal recessive primary ovarian insufficiency. *J Clin Endocrinol Metab* 2014;**10**:E2129–E2132.
- de Zegher F, Jaeken J. Endocrinology of the carbohydrate-deficient glycoprotein syndrome type I from birth through adolescence. *Pediatr Res* 1995;**4 Pt 1**:395–401.
- Di Pasquale E, Beck-Peccoz P, Persani L. Hypergonadotropic ovarian failure associated with an inherited mutation of human bone morphogenetic protein-15 (BMP15) gene. *Am J Hum Genet* 2004;**1**:106–111.
- Di Pasquale E, Rossetti R, Marozzi A, Bodega B, Borgato S, Cavallo L, Einaudi S, Radetti G, Russo G, Sacco M et al. Identification of new variants of human BMP15 gene in a large cohort of women with premature ovarian failure. *J Clin Endocrinol Metab* 2006;**5**:1976–1979.
- Dixit H, Deendayal M, Singh L. Mutational analysis of the mature peptide region of inhibin genes in Indian women with ovarian failure. *Hum Reprod* 2004;**8**:1760–1764.
- Dixit H, Rao LK, Padmalatha V, Kanakavalli M, Deenadayal M, Gupta N, Chakravarty B, Singh L. Mutational analysis of the coding region of growth differentiation factor 9 gene in Indian women with ovarian failure. *Menopause* 2005;**6**:749–754.
- Dixit H, Rao KL, Padmalatha V, Kanakavalli M, Deenadayal M, Gupta N, Chakravarty BN, Singh L. Expansion of the germline analysis for the INHA gene in Indian women with ovarian failure. *Hum Reprod* 2006a;**6**:1643–1644.
- Dixit H, Rao KL, Padmalatha VV, Kanakavalli M, Deenadayal M, Gupta N, Chakravarty BN, Singh L. Mutational analysis of the betaglycan gene-coding region in susceptibility for ovarian failure. *Hum Reprod* 2006b;**8**:2041–2046.
- Dixit H, Rao LK, Padmalatha VV, Kanakavalli M, Deenadayal M, Gupta N, Chakravarty B, Singh L. Missense mutations in the BMP15 gene are associated with ovarian failure. *Hum Genet* 2006c;**4**:408–415.
- Dudding TE, Lawrence O, Winship I, Froyen G, Vandewalle J, Scott R, Shelling AN. Array comparative genomic hybridization for the detection of submicroscopic copy number variations of the X chromosome in women with premature ovarian failure. *Hum Reprod* 2010;**12**:3159–3160; author reply 3160–3151.
- Duncan AJ, Knight JA, Costello H, Conway GS, Rahman S. POLG mutations and age at menopause. *Hum Reprod* 2012;**7**:2243–2244.
- Ellis NA, German J. Molecular genetics of Bloom's syndrome. *Hum Mol Genet* 1996;**14**:57–74.
- Engmann L, Losel R, Wehling M, Peluso JJ. Progesterone regulation of human granulosa/luteal cell viability by an RU486-independent mechanism. *J Clin Endocrinol Metab* 2006;**12**:4962–4968.
- Epstein CJ, Martin GM, Schultz AL, Motulsky AG. Werner's syndrome a review of its symptomatology, natural history, pathologic features, genetics and relationship to the natural aging process. *Medicine* 1966;**3**:177–221.
- European Society of Human Genetics; European Society of Human Reproduction and Embryology. The need for interaction between assisted reproduction technology and genetics. Recommendations of the European Societies of Human Genetics and Human Reproduction and Embryology. *Eur J Hum Genet* 2006b;**5**:509–511.
- Fierabracci A, Bizzarri C, Palma A, Milillo A, Bellacchio E, Cappa M. A novel heterozygous mutation of the AIRE gene in a patient with autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy syndrome (APECED). *Gene* 2012;**1**:113–117.
- Fogli A, Rodriguez D, Eymard-Pierre E, Bouhour F, Labauge P, Meaney BF, Zeesman S, Kaneski CR, Schiffmann R, Boespflug-Tanguy O. Ovarian failure related to eukaryotic initiation factor 2B mutations. *Am J Hum Genet* 2003;**6**:1544–1550.
- Fogli A, Gauthier-Barichard F, Schiffmann R, Vanderhoof VH, Bakalov VK, Nelson LM, Boespflug-Tanguy O. Screening for known mutations in EIF2B genes in a large panel of patients with premature ovarian failure. *BMC Womens Health* 2004;**1**:8.
- Fonseca DJ, Garzon E, Lakkhal B, Braham R, Ojeda D, Elghezal H, Saad A, Restrepo CM, Laissue P. Screening for mutations of the FOXO4 gene in premature ovarian failure patients. *Reprod Biomed Online* 2012a;**3**:339–341.
- Fonseca DJ, Ojeda D, Lakkhal B, Braham R, Eggers S, Turbitt E, White S, Grover S, Warne G, Zacharin M et al. CITED2 mutations potentially cause idiopathic premature ovarian failure. *Transl Res* 2012b;**5**:384–388.
- Fonseca DJ, Patiño LCI, Suárez YC, de Jesús Rodríguez A, Mateus HE, Jiménez KM, Ortega-Recalde O, Díaz-Yamal I, Laissue P. Next generation sequencing in women affected by nonsyndromic premature ovarian failure displays new potential causative genes and mutations. *Fertil Steril* 2015;**104**:154–162.
- Foresta C, Ferlin A, Gianaroli L, Dallapiccola B. Guidelines for the appropriate use of genetic tests in infertile couples. *Eur J Hum Genet* 2002;**5**:303–312.
- Fridovich-Keil JL, Gubbels CS, Spencer JB, Sanders RD, Land JA, Rubio-Gozalbo E. Ovarian function in girls and women with GALT-deficiency galactosemia. *J Inherit Metab Dis* 2011;**2**:357–366.
- Galloway SM, McNatty KP, Cambridge LM, Laitinen MP, Juengel JL, Jokiranta TS, McLaren RJ, Luiro K, Dodds KG, Montgomery GW et al. Mutations in an oocyte-derived growth factor gene (BMP15) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nat Genet* 2000;**3**:279–283.
- Gatti RA, Boder E, Vinters HV, Sparkes RS, Norman A, Lange K. Ataxia-telangiectasia: an interdisciplinary approach to pathogenesis. *Medicine* 1991;**2**:99–117.
- Gersak K, Meden-Vrtovc H, Peterlin B. Fragile X premutation in women with sporadic premature ovarian failure in Slovenia. *Hum Reprod* 2003;**8**:1637–1640.

- Gong Y, Krakow D, Marcelino J, Wilkin D, Chitayat D, Babul-Hirji R, Hudgins L, Cremers CW, Cremers FP, Brunner HG et al. Heterozygous mutations in the gene encoding noggin affect human joint morphogenesis. *Nat Genet* 1999; **3**:302–304.
- Goswami R, Goswami D, Kabra M, Gupta N, Dubey S, Dadhwal V. Prevalence of the triple X syndrome in phenotypically normal women with premature ovarian failure and its association with autoimmune thyroid disorders. *Fertil Steril* 2003; **4**:1052–1054.
- Graziewicz MA, Bienstock RJ, Copeland WC. The DNA polymerase gamma Y955C disease variant associated with PEO and parkinsonism mediates the incorporation and translesion synthesis opposite 7,8-dihydro-8-oxo-2'-deoxyguanosine. *Hum Mol Genet* 2007; **22**:2729–2739.
- Guiraldelli MF, Eyster C, Wilkerson JL, Dresser ME, Pezza RJ. Mouse HFM1/Mer3 is required for crossover formation and complete synapsis of homologous chromosomes during meiosis. *PLoS Genet* 2013; **3**:e1003383.
- Guo T, Qin Y, Jiao X, Li G, Simpson JL, Chen ZJ. FMR1 premutation is an uncommon explanation for premature ovarian failure in Han Chinese. *PLoS One* 2014; **7**:e103316.
- Harris SE, Chand AL, Winship IM, Gersak K, Aittomaki K, Shelling AN. Identification of novel mutations in FOXL2 associated with premature ovarian failure. *Mol Hum Reprod* 2002; **8**:729–733.
- Harris SE, Chand AL, Winship IM, Gersak K, Nishi Y, Yanase T, Nawata H, Shelling AN. INHA promoter polymorphisms are associated with premature ovarian failure. *Mol Hum Reprod* 2005; **11**:779–784.
- Hens L, Devroey P, Van Waesberghe L, Bonduelle M, Van Steirteghem AC, Liebaers I. Chromosome studies and fertility treatment in women with ovarian failure. *Clin Genet* 1989; **2**:81–91.
- Holland CM. 47,XXX in an adolescent with premature ovarian failure and autoimmune disease. *J Pediatr Adolesc Gynecol* 2001; **7**:77–80.
- Hui ES, Udofa EA, Soto J, Vanderhoof VH, Zachman K, Tong ZB, Nelson LM. Investigation of the human stem cell factor KIT ligand gene, KITLG, in women with 46,XX spontaneous premature ovarian failure. *Fertil Steril* 2006; **5**:1502–1507.
- Inagaki K, Shimasaki S. Impaired production of BMP-15 and GDF-9 mature proteins derived from proproteins WITH mutations in the proregion. *Mol Cell Endocrinol* 2010; **1–2**:1–7.
- Ishizuka B, Okamoto N, Hamada N, Sugishita Y, Saito J, Takahashi N, Ogata T, Itoh MT. Number of CGG repeats in the FMR1 gene of Japanese patients with primary ovarian insufficiency. *Fertil Steril* 2011; **5**:1170–1174.
- Jaaskelainen M, Prunskaitė-Hyyryläinen R, Naillat F, Parviainen H, Anttonen M, Heikinheimo M, Liakka A, Ola R, Vainio S, Vaskivuo TE et al. WNT4 is expressed in human fetal and adult ovaries and its signaling contributes to ovarian cell survival. *Mol Cell Endocrinol* 2010; **1–2**:106–111.
- Janse F, Knauff EA, Niermeijer MF, Eijkemans MJ, Laven JS, Lambalk CB, Fauser BC, Goverde AJ. Similar phenotype characteristics comparing familial and sporadic premature ovarian failure. *Menopause* 2010; **4**:758–765.
- Janse F, de With LM, Duran KJ, Kloosterman WP, Goverde AJ, Lambalk CB, Laven JS, Fauser BC, Giltay JC; Dutch Primary Ovarian Insufficiency Consortium. Limited contribution of NR5A1 (SF-1) mutations in women with primary ovarian insufficiency (POI). *Fertil Steril* 2012; **1**:141–146.
- Jenkinson EM, Rehman AU, Walsh T, Clayton-Smith J, Lee K, Morell RJ, Drummond MC, Khan SN, Naeem MA, Rauf B et al. Perrault syndrome is caused by recessive mutations in CLPP, encoding a mitochondrial ATP-dependent chambered protease. *Am J Hum Genet* 2013; **4**:605–613.
- Jiang M, Aittomaki K, Nilsson C, Pakarinen P, Iitia A, Torresani T, Simonsen H, Goh V, Pettersson K, de la Chapelle A et al. The frequency of an inactivating point mutation (566C→T) of the human follicle-stimulating hormone receptor gene in four populations using allele-specific hybridization and time-resolved fluorometry. *J Clin Endocrinol Metab* 1998; **12**:4338–4343.
- Jiao X, Qin C, Li J, Qin Y, Gao X, Zhang B, Zhen X, Feng Y, Simpson JL, Chen ZJ. Cytogenetic analysis of 531 Chinese women with premature ovarian failure. *Hum Reprod* 2012; **7**:2201–2207.
- Jiao X, Qin Y, Li G, Zhao S, You L, Ma J, Simpson JL, Chen ZJ. Novel NR5A1 missense mutation in premature ovarian failure: detection in Han Chinese indicates causation in different ethnic groups. *PLoS One* 2013; **9**:e74759.
- Kalantari H, Madani T, Zari Moradi S, Mansouri Z, Almadani N, Gourabi H, Mohseni Meybodi A. Cytogenetic analysis of 179 Iranian women with premature ovarian failure. *Gynecol Endocrinol* 2013; **6**:588–591.
- Kang H, Lee SK, Kim MH, Song J, Bae SJ, Kim NK, Lee SH, Kwack K. Parathyroid hormone-responsive B1 gene is associated with premature ovarian failure. *Hum Reprod* 2008; **6**:1457–1465.
- Kasippillai T, MacArthur DG, Kirby A, Thomas B, Lambalk CB, Daly MJ, Welt CK. Mutations in *ELF4ENIF1* are associated with primary ovarian insufficiency. *J Clin Endocrinol Metab* 2013; **9**:E1534–E1539.
- Kaufman F, Kogut MD, Donnell GN, Koch H, Goebelsmann U. Ovarian failure in galactosaemia. *Lancet* 1979; **8145**:737–738.
- Kaufman FR, Kogut MD, Donnell GN, Goebelsmann U, March C, Koch R. Hypergonadotropic hypogonadism in female patients with galactosemia. *N Engl J Med* 1981; **17**:994–998.
- Kawano Y, Narahara H, Matsui N, Miyakawa I. Premature ovarian failure associated with a Robertsonian translocation. *Acta Obstet Gynecol Scand* 1998; **4**:467–469.
- Knauff EA, Franke L, van Es MA, van den Berg LH, van der Schouw YT, Laven JS, Lambalk CB, Hoek A, Goverde AJ, Christin-Maitre S et al. Genome-wide association study in premature ovarian failure patients suggests ADAMTS19 as a possible candidate gene. *Hum Reprod* 2009; **9**:2372–2378.
- Knauff EA, Blauw HM, Pearson PL, Kok K, Wijmenga C, Veldink JH, van den Berg LH, Bouchard P, Fauser BC, Franke L. Copy number variants on the X chromosome in women with primary ovarian insufficiency. *Fertil Steril* 2011; **5**:1584–1588.
- Kneitz B, Cohen PE, Avdievich E, Zhu L, Kane MF, Hou H Jr, Kolodner RD, Kuchlerapati R, Pollard JW, Edelmann W. MutS homolog 4 localization to meiotic chromosomes is required for chromosome pairing during meiosis in male and female mice. *Genes Dev* 2000; **9**:1085–1097.
- Kosaki K, Sato S, Hasegawa T, Matsuo N, Suzuki T, Ogata T. Premature ovarian failure in a female with proximal symphalangism and Noggin mutation. *Fertil Steril* 2004; **4**:1137–1139.
- Kovanci E, Rohozinski J, Simpson JL, Heard MJ, Bishop CE, Carson SA. Growth differentiating factor-9 mutations may be associated with premature ovarian failure. *Fertil Steril* 2007; **1**:143–146.
- Kovanci E, Simpson JL, Amato P, Rohozinski J, Heard MJ, Bishop CE, Carson SA. Oocyte-specific G-protein-coupled receptor 3 (GPR3): no perturbations found in 82 women with premature ovarian failure (first report). *Fertil Steril* 2008; **4**:1269–1271.
- Kristiansson M, Soop M, Saraste L, Sundqvist KG, Suontaka AM, Blomback M. Cytokine and coagulation characteristics of retrieved blood after arthroplasty. *Intensive Care Med* 1995; **12**:989–995.
- Lacombe A, Lee H, Zahed L, Choucair M, Muller JM, Nelson SF, Salameh WW, Vilain E. Disruption of POF1B binding to nonmuscle actin filaments is associated with premature ovarian failure. *Am J Hum Genet* 2006; **1**:113–119.
- Laissue P, Christin-Maitre S, Touraine P, Kuttann F, Ritvos O, Aittomaki K, Bourcigaux N, Jacquesson L, Bouchard P, Frydman R et al. Mutations and sequence variants in GDF9 and BMP15 in patients with premature ovarian failure. *Eur J Endocrinol* 2006; **5**:739–744.
- Laissue P, Christin-Maitre S, Bouchard P, Fellous M, Veitia RA. Mutations in the NOG gene are not a common cause of nonsyndromic premature ovarian failure. *Clin Endocrinol (Oxf)* 2007; **6**:900.
- Lakhal B, Laissue P, Braham R, Elghezal H, Saad A, Fellous M, Veitia RA. A novel BMP15 variant, potentially affecting the signal peptide, in a familial case of premature ovarian failure. *Clin Endocrinol (Oxf)* 2009; **5**:752–753.
- Lakhal B, Braham R, Berguigua R, Bouali N, Zaouali M, Chaieb M, Veitia RA, Saad A, Elghezal H. Cytogenetic analyses of premature ovarian failure using karyotyping and interphase fluorescence in situ hybridization (FISH) in a group of 1000 patients. *Clin Genet* 2010a; **2**:181–185.
- Lakhal B, Laissue P, Braham R, Elghezal H, Saad A, Fellous M, Veitia RA. BMP15 and premature ovarian failure: causal mutations, variants, polymorphisms? *Clin Endocrinol (Oxf)* 2010b; **3**:425–426.
- Lakhal B, Ben-Hadj-Khalifa S, Bouali N, Philipert P, Audran F, Braham R, Hatem E, Sultan C, Saad A. Mutational screening of SF1 and WNT4 in Tunisian women with premature ovarian failure. *Gene* 2012; **2**:298–301.
- Lechowska A, Bilinski S, Choi Y, Shin Y, Kloc M, Rajkovic A. Premature ovarian failure in nobox-deficient mice is caused by defects in somatic cell invasion and germ cell cyst breakdown. *J Assist Reprod Genet* 2011; **7**:583–589.
- Ledig S, Ropke A, Haeusler G, Hinney B, Wieacker P. BMP15 mutations in XX gonadal dysgenesis and premature ovarian failure. *Am J Obstet Gynecol* 2008; **1**:84.
- Ledig S, Ropke A, Wieacker P. Copy number variants in premature ovarian failure and ovarian dysgenesis. *Sex Dev* 2010; **4–5**:225–232.

- Liu L, Rajareddy S, Reddy P, Du C, Jagarlamudi K, Shen Y, Gunnarsson D, Selstam G, Boman K, Liu K. Infertility caused by retardation of follicular development in mice with oocyte-specific expression of Foxo3a. *Development* 2007; **1**:199–209.
- Lorda-Sanchez JJ, Ibanez AJ, Sanz RJ, Trujillo MJ, Anabitarte ME, Querejeta ME, Rodriguez de Alba M, Gimenez A, Infantes F, Ramos C et al. Choroideremia, sensorineural deafness, and primary ovarian failure in a woman with a balanced X-4 translocation. *Ophthalmic Genet* 2000; **3**:185–189.
- Losel RM, Besong D, Peluso JJ, Wehling M. Progesterone receptor membrane component 1—many tasks for a versatile protein. *Steroids* 2008; **9–10**:929–934.
- Lourenco D, Brauner R, Lin L, De Perdigo A, Weryha G, Muresan M, Boudjenah R, Guerra-Junior G, Maciel-Guerra AT, Achermann JC et al. Mutations in NR5A1 associated with ovarian insufficiency. *N Engl J Med* 2009; **12**:1200–1210.
- Luoma P, Melberg A, Rinne JO, Kaukonen JA, Nupponen NN, Chalmers RM, Oldfors A, Rautakorpi I, Peltonen L, Majamaa K et al. Parkinsonism, premature menopause, and mitochondrial DNA polymerase gamma mutations: clinical and molecular genetic study. *Lancet* 2004; **9437**:875–882.
- Lutzmann M, Grey C, Traver S, Ganier O, Maya-Mendoza A, Ranisavljevic N, Bernex F, Nishiyama A, Montel N, Gavois E et al. MCM8- and MCM9-deficient mice reveal gametogenesis defects and genome instability due to impaired homologous recombination. *Mol cell* 2012; **4**:523–534.
- Mandon-Pepin B, Touraine P, Kuttann F, Derbois C, Rouxel A, Matsuda F, Nicolas A, Cotinot C, Fellous M. Genetic investigation of four meiotic genes in women with premature ovarian failure. *Eur J Endocrinol* 2008; **1**:107–115.
- Mansouri MR, Schuster J, Badhai J, Stattin EL, Losel R, Wehling M, Carlsson B, Hovatta O, Karlstrom PO, Golovleva I et al. Alterations in the expression, structure and function of progesterone receptor membrane component-1 (PGRMC1) in premature ovarian failure. *Hum Mol Genet* 2008; **23**:3776–3783.
- Marozzi A, Porta C, Vegetti W, Crosignani PG, Tibiletti MG, Dalpra L, Ginelli E. Mutation analysis of the inhibin alpha gene in a cohort of Italian women affected by ovarian failure. *Hum Reprod* 2002; **7**:1741–1745.
- Mathis S, Scheper GC, Baumann N, Petit E, Gil R, van der Knaap MS, Neau JP. The ovarioleukodystrophy. *Clin Neurol Neurosurg* 2008; **10**:1035–1037.
- Matsui Y, Toksoz D, Nishikawa S, Nishikawa S, Williams D, Zsebo K, Hogan BL. Effect of Steel factor and leukaemia inhibitory factor on murine primordial germ cells in culture. *Nature* 1991; **6346**:750–752.
- May-Panloup P, Chretien MF, Jacques C, Vasseur C, Malthiery Y, Reynier P. Low oocyte mitochondrial DNA content in ovarian insufficiency. *Hum Reprod* 2005; **3**:593–597.
- McGuire MM, Bowden W, Engel NJ, Ahn HW, Kovanci E, Rajkovic A. Genomic analysis using high-resolution single-nucleotide polymorphism arrays reveals novel microdeletions associated with premature ovarian failure. *Fertil Steril* 2011; **5**:1595–1600.
- Mehlmann LM, Saeki Y, Tanaka S, Brennan TJ, Evsikov AV, Pendola FL, Knowles BB, Eppig JJ, Jaffe LA. The Gs-linked receptor GPR3 maintains meiotic arrest in mammalian oocytes. *Science* 2004; **5703**:1947–1950.
- Morino H, Pierce SB, Matsuda Y, Walsh T, Ohsawa R, Newby M, Hiraki-Kamon K, Kuramochi M, Lee MK, Klevit RE et al. Mutations in Twinkle primase-helicase cause Perrault syndrome with neurologic features. *Neurology* 2014; **22**:2054–2061.
- Nishimura-Tadaki A, Wada T, Bano G, Gough K, Warner J, Kosho T, Ando N, Hamanoue H, Sakakibara H, Nishimura G et al. Breakpoint determination of X;autosomal balanced translocations in four patients with premature ovarian failure. *J Hum Genet* 2011; **2**:156–160.
- Norling A, Hirschberg AL, Rodriguez-Wallberg KA, Iwarsson E, Wedell A, Barbaro M. Identification of a duplication within the GDF9 gene and novel candidate genes for primary ovarian insufficiency (POI) by a customized high-resolution array comparative genomic hybridization platform. *Hum Reprod* 2014; **8**:1818–1827.
- Novoa I, Gallego J, Ferreira PG, Mendez R. Mitotic cell-cycle progression is regulated by CPEB1 and CPEB4-dependent translational control. *Nat Cell Biol* 2010; **5**:447–456.
- Ojeda D, Lakhal B, Fonseca DJ, Braham R, Landolsi H, Mateus HE, Restrepo CM, Elghezal H, Saad A, Laissue P. Sequence analysis of the CDKN1B gene in patients with premature ovarian failure reveals a novel mutation potentially related to the phenotype. *Fertil Steril* 2011; **8**:2658–2660.
- Oldenburg RA, van Dooren MF, de Graaf B, Simons E, Govaerts L, Swagemakers S, Verkerk JM, Oostra BA, Bertoli-Avella AM. A genome-wide linkage scan in a Dutch family identifies a premature ovarian failure susceptibility locus. *Hum Reprod* 2008; **12**:2835–2841.
- Oley C, Baraitser M. Blepharophimosis, ptosis, epicanthus inversus syndrome (BPES syndrome). *J Med Genet* 1988; **1**:47–51.
- Orczyk GP, Pehrson J, Leventhal JM. Premature ovarian failure in a 35-year-old woman with a Robertsonian translocation. *Int J Fertil* 1989; **3**:184–187.
- Pagnamenta AT, Taanman JW, Wilson CJ, Anderson NE, Marotta R, Duncan AJ, Bitner-Glindzicz M, Taylor RW, Laskowski A, Thorburn DR et al. Dominant inheritance of premature ovarian failure associated with mutant mitochondrial DNA polymerase gamma. *Hum Reprod* 2006; **10**:2467–2473.
- Panda B, Rao L, Tosh D, Dixit H, Padmalatha V, Kanakavalli M, Raseswari T, Deenadayal M, Gupta N, Chakrabarty B et al. Germline study of AR gene of Indian women with ovarian failure. *Gynecol Endocrinol* 2010; **8**:572–578.
- Pangas SA, Choi Y, Ballou DJ, Zhao Y, Westphal H, Matzuk MM, Rajkovic A. Oogenesis requires germ cell-specific transcriptional regulators Sohlh1 and Lhx8. *Proc Natl Acad Sci USA* 2006; **21**:8090–8095.
- Peluso JJ, Pappalardo A, Losel R, Wehling M. Progesterone receptor membrane component 1 expression in the immature rat ovary and its role in mediating progesterone's antiapoptotic action. *Endocrinology* 2006; **6**:3133–3140.
- Persani L, Rossetti Z, Cacciatori C, Bonomi M. Primary Ovarian Insufficiency: X chromosome defects and autoimmunity. *J Autoimmun* 2009; **1**:35–41.
- Philibert P, Leprieux E, Zenaty D, Thibaud E, Polak M, Frances AM, Lespinasse J, Raingeard I, Servant N, Audran F et al. Steroidogenic factor-1 (SF-1) gene mutation as a frequent cause of primary amenorrhea in 46,XY female adolescents with low testosterone concentration. *Reprod Biol Endocrinol* 2010; **8**:28.
- Pierce SB, Walsh T, Chisholm KM, Lee MK, Thornton AM, Fiumara A, Opitz JM, Levy-Lahad E, Klevit RE, King MC. Mutations in the DBP-deficiency protein HSD17B4 cause ovarian dysgenesis, hearing loss, and ataxia of Perrault Syndrome. *Am J Hum Genet* 2010; **2**:282–288.
- Pierce SB, Chisholm KM, Lynch ED, Lee MK, Walsh T, Opitz JM, Li W, Klevit RE, King MC. Mutations in mitochondrial histidyl tRNA synthetase HARS2 cause ovarian dysgenesis and sensorineural hearing loss of Perrault syndrome. *Proc Natl Acad Sci USA* 2011; **16**:6543–6548.
- Pierce SB, Gersak K, Michaelson-Cohen R, Walsh T, Lee MK, Malach D, Klevit RE, King MC, Levy-Lahad E. Mutations in LARS2, encoding mitochondrial leucyl-tRNA synthetase, lead to premature ovarian failure and hearing loss in Perrault syndrome. *Am J Hum Genet* 2013; **4**:614–620.
- Pisarska MD, Kuo FT, Tang D, Zarrini P, Khan S, Ketefian A. Expression of forkhead transcription factors in human granulosa cells. *Fertil Steril* 2009; **4** Suppl:1392–1394.
- Portnoi MF, Aboura A, Tachdjian G, Bouchard P, Dewailly D, Bourcigaux N, Frydman R, Reys AC, Brisset S, Christin-Maitre S. Molecular cytogenetic studies of Xq critical regions in premature ovarian failure patients. *Hum Reprod* 2006; **9**:2329–2334.
- Prakash GJ, Kanth VV, Shelling AN, Rozati R, Sujatha M. Absence of 566C>T mutation in exon 7 of the FSHR gene in Indian women with premature ovarian failure. *Int J Gynaecol Obstet* 2009; **3**:265–266.
- Prueitt RL, Ross JL, Zinn AR. Physical mapping of nine Xq translocation breakpoints and identification of XPNPEP2 as a premature ovarian failure candidate gene. *Cytogenet Cell Genet* 2000; **1-2**:44–50.
- Prueitt RL, Chen H, Barnes RI, Zinn AR. Most X;autosomal translocations associated with premature ovarian failure do not interrupt X-linked genes. *Cytogenet Genome Res* 2002; **1-2**:32–38.
- Pyun JA, Cha DH, Kwack K. LAMC1 gene is associated with premature ovarian failure. *Maturitas* 2012; **4**:402–406.
- Qin Y, Poirier C, Truong C, Schumacher A, Agoulnik AI, Bishop CE. A major locus on mouse chromosome 18 controls XX sex reversal in Odd Sex (Ods) mice. *Hum Mol Genet* 2003; **5**:509–515.
- Qin Y, Kong LK, Poirier C, Truong C, Overbeek PA, Bishop CE. Long-range activation of Sox9 in Odd Sex (Ods) mice. *Hum Mol Genet* 2004; **12**:1213–1218.
- Qin Y, Choi Y, Zhao H, Simpson JL, Chen ZJ, Rajkovic A. NOBOX homeobox mutation causes premature ovarian failure. *Am J Hum Genet* 2007a; **3**:576–581.
- Qin Y, Zhao H, Kovanci E, Simpson JL, Chen ZJ, Rajkovic A. Mutation analysis of NANOS3 in 80 Chinese and 88 Caucasian women with premature ovarian failure. *Fertil Steril* 2007b; **5**:1465–1467.
- Qin Y, Shi Y, Zhao Y, Carson SA, Simpson JL, Chen ZJ. Mutation analysis of NOBOX homeodomain in Chinese women with premature ovarian failure. *Fertil Steril* 2009; **4**(Suppl):1507–1509.
- Qin CR, Chen SL, Yao JL, Wu WQ, Xie JS. Identification of novel missense mutations of the TGFBR3 gene in Chinese women with premature ovarian failure. *Reprod Biomed Online* 2011; **6**:697–703.
- Qin CR, Chen SL, Yao JL, Li T, Wu WQ. Haplotype and mutation analysis of the TGFBR3 gene in Chinese women with idiopathic premature ovarian failure. *Gynecol Endocrinol* 2012a; **1**:63–67.

- Qin Y, Zhao H, Xu J, Shi Y, Li Z, Qiao J, Liu J, Qin C, Ren C, Li J *et al.* Association of 8q22.3 locus in Chinese Han with idiopathic premature ovarian failure (POF). *Hum Mol Genet* 2012b;**21**:430–436.
- Qin Y, Sun M, You L, Wei D, Sun J, Liang X, Zhang B, Jiang H, Xu J, Chen Z. ESRI1, HK3 and BRSK1 gene variants are associated with both age at natural menopause and premature ovarian failure. *Orphanet J Rare Dis* 2012c;**7**:5.
- Qin Y, Jiao X, Dalgleish R, Vujovic S, Li J, Simpson JL, Al-Azzawi F, Chen ZJ. Novel variants in the SOHLH2 gene are implicated in human premature ovarian failure. *Fertil Steril* 2014a;**4**:1104–1109.
- Qin C, Yuan Z, Yao J, Zhu W, Wu W, Xie J. AMH and AMHR2 genetic variants in Chinese women with primary ovarian insufficiency and normal age at natural menopause. *Reprod Biomed Online* 2014b;**3**:311–318.
- Quilter CR, Karcnias AC, Bagga MR, Duncan S, Murray A, Conway GS, Sargent CA, Affara NA. Analysis of X chromosome genomic DNA sequence copy number variation associated with premature ovarian failure (POF). *Hum Reprod* 2010;**8**:2139–2150.
- Rajareddy S, Reddy P, Du C, Liu L, Jagarlamudi K, Tang W, Shen Y, Berthet C, Peng SL, Kaldis P *et al.* p27kip1 (cyclin-dependent kinase inhibitor 1B) controls ovarian development by suppressing follicle endowment and activation and promoting follicle atresia in mice. *Mol Endocrinol* 2007;**9**:2189–2202.
- Rajkovic A, Pangas SA, Ballow D, Suzumori N, Matzuk MM. NOBOX deficiency disrupts early folliculogenesis and oocyte-specific gene expression. *Science* 2004;**5687**:1157–1159.
- Rebar RW, Connolly HV. Clinical features of young women with hypergonadotropic amenorrhea. *Fertil Steril* 1990;**5**:804–810.
- Reddy P, Liu L, Adhikari D, Jagarlamudi K, Rajareddy S, Shen Y, Du C, Tang WL, Hamalainen T, Peng SL *et al.* Oocyte-specific deletion of Pten causes premature activation of the primordial follicle pool. *Science* 2008;**5863**:611–613.
- Reiss AL, Hall SS. Fragile X syndrome: assessment and treatment implications. *Child Adolesc Psychiatr Clin N Am* 2007;**3**:663–675.
- Rizzolio F, Bione S, Sala C, Goegan M, Gentile M, Gregato G, Rossi E, Pramparo T, Zuffardi O, Toniolo D. Chromosomal rearrangements in Xq and premature ovarian failure: mapping of 25 new cases and review of the literature. *Hum Reprod* 2006;**6**:1477–1483.
- Rizzolio F, Sala C, Alboresi S, Bione S, Gilli S, Goegan M, Pramparo T, Zuffardi O, Toniolo D. Epigenetic control of the critical region for premature ovarian failure on autosomal genes translocated to the X chromosome: a hypothesis. *Hum Genet* 2007;**3**:441–450.
- Rizzolio F, Pramparo T, Sala C, Zuffardi O, De Santis L, Rabellotti E, Calzi F, Fusi F, Bellazzi R, Toniolo D. Epigenetic analysis of the critical region 1 for premature ovarian failure: demonstration of a highly heterochromatic domain on the long arm of the mammalian X chromosome. *J Med Genet* 2009;**9**:585–592.
- Rossetti R, Di Pasquale E, Marozzi A, Bione S, Toniolo D, Grammatico P, Nelson LM, Beck-Peccoz P, Persani L. BMP15 mutations associated with primary ovarian insufficiency cause a defective production of bioactive protein. *Hum Mutat* 2009;**5**:804–810.
- Santos MG, Machado AZ, Martins CN, Domenice S, Costa EM, Nishi MY, Ferraz-de-Souza B, Jorge SA, Pereira CA, Soardi FC *et al.* Homozygous inactivating mutation in NANOS3 in two sisters with primary ovarian insufficiency. *Biomed Res Int* 2014;**7**:787465.
- Sarto GE, Therman E, Patau K. X inactivation in man: a woman with t(Xq-;12q+). *Am J Hum Genet* 1973;**3**:262–270.
- Schadewaldt P, Kamalanathan L, Hammen HW, Wendel U. Age dependence of endogenous galactose formation in Q188R homozygous galactosemic patients. *Mol Genet Metab* 2004;**1**:31–44.
- Schiffmann R, Tedeschi G, Kinkel RP, Trapp BD, Frank JA, Kaneski CR, Brady RO, Barton NW, Nelson L, Yanovski JA. Leukodystrophy in patients with ovarian dysgenesis. *Ann Neurol* 1997;**5**:654–661.
- Shelling AN, Burton KA, Chand AL, van Ee CC, France JT, Farquhar CM, Milsom SR, Love DR, Gersak K, Aittomaki K *et al.* Inhibin: a candidate gene for premature ovarian failure. *Hum Reprod* 2000;**12**:2644–2649.
- Shi Y, Zhao H, Cao Y, Yang D, Li Z, Zhang B, Liang X, Li T, Chen J, Shen J *et al.* Genome-wide association study identifies eight new risk loci for polycystic ovary syndrome. *Nat Genet* 2012;**9**:1020–1025.
- Shiina H, Matsumoto T, Sato T, Igarashi K, Miyamoto J, Takemasa S, Sakari M, Takada I, Nakamura T, Metzger D *et al.* Premature ovarian failure in androgen receptor-deficient mice. *Proc Natl Acad Sci USA* 2006;**1**:224–229.
- Shimizu Y, Kimura F, Takebayashi K, Fujiwara M, Takakura K, Takahashi K. Mutational analysis of the PTEN gene in women with premature ovarian failure. *Acta Obstet Gynecol Scand* 2009;**7**:824–825.
- Simpson JL. Gonadal dysgenesis and abnormalities of the human sex chromosomes: current status of phenotypic-karyotypic correlations. *Birth Defects Orig Artic Ser* 1975;**4**:23–59.
- Simpson JL. Genetic and phenotypic heterogeneity in ovarian failure: overview of selected candidate genes. *Ann N Y Acad Sci* 2008;**1135**:146–154.
- Simpson JL. Disorders of the gonads, genital tract, and genitalia. In: Rimoin DL, Connor JM, Pyertiz RE, Korf BR (eds). *Emery and Rimoin's Principles and Practice of Medical Genetics*. New York: Churchill-Livingstone, 2013, 4700 pp.
- Simpson JL. Genetics of female infertility due to anomalies of the ovary and mullerian ducts. *Methods Mol Biol* 2014;**1154**:39–73.
- Simpson JL, Elias S. *Genetics in Obstetrics and Gynecology*, 3rd edn. Philadelphia: Saunders, 2003.
- Simpson JL, Rajkovic A. Ovarian differentiation and gonadal failure. *Am J Med Genet* 1999;**4**:186–200.
- Sparks SE, Krasnewich DM. PMM2-CDG CDG-la GeneReviews™ [Internet]. <http://www.ncbi.nlm.nih.gov/books/NBK1110/>. 2005.
- Spath MA, Nillesen VN, Smits AP, Feuth TB, Braat DD, van Kessel AG, Yntema HG. X chromosome inactivation does not define the development of premature ovarian failure in fragile X premutation carriers. *Am J Med Genet A* 2010;**2**:387–393.
- Stamatoyannopoulos JA, Snyder M, Hardison R, Ren B, Gingeras T, Gilbert DM, Groudine M, Bender M, Kaul R, Canfield T *et al.* An encyclopedia of mouse DNA elements (Mouse ENCODE). *Genome Biol* 2012;**8**:418.
- Su Y, Swift M. Mortality rates among carriers of ataxia-telangiectasia mutant alleles. *Ann Intern Med* 2000;**10**:770–778.
- Sugawa F, Wada Y, Maruyama T, Uchida H, Ishizuka B, Ogata T. Premature ovarian failure and androgen receptor gene CAG repeat lengths weighted by X chromosome inactivation patterns. *Fertil Steril* 2009;**2**:649–652.
- Sundblad V, Chiauzzi VA, Escobar ME, Dain L, Charreau EH. Screening of FSH receptor gene in Argentine women with premature ovarian failure (POF). *Mol Cell Endocrinol* 2004;**1**:253–59.
- Takakura K, Takebayashi K, Wang HQ, Kimura F, Kasahara K, Noda Y. Follicle-stimulating hormone receptor gene mutations are rare in Japanese women with premature ovarian failure and polycystic ovary syndrome. *Fertil Steril* 2001;**1**:207–209.
- Takebayashi K, Takakura K, Wang HQ, Kimura F, Kasahara K, Noda Y. Mutation analysis of the growth differentiation factor-9 and-9B genes in patients with premature ovarian failure and polycystic ovary syndrome. *Fertil Steril* 2000;**5**:976–979.
- Tanaka K, Miyamoto N, Shouguchi-Miyata J, Ikeda JE. HFM1, the human homologue of yeast Mer3, encodes a putative DNA helicase expressed specifically in germ-line cells. *DNA Seq* 2006;**3**:242–246.
- Tartaglia NR, Howell S, Sutherland A, Wilson R, Wilson L. A review of trisomy X (47,XXX). *Orphanet J Rare Dis* 2010;**5**:8.
- Tiotiu D, Alvaro Mercadal B, Imbert R, Verbist J, Demeestere I, De Leener A, Englert Y, Vassart G, Costagliola S, Delbaere A. Variants of the BMP15 gene in a cohort of patients with premature ovarian failure. *Hum Reprod* 2010;**6**:1581–1587.
- Tong Y, Liao WX, Roy AC, Ng SC. Absence of mutations in the coding regions of follicle-stimulating hormone receptor gene in Singapore Chinese women with premature ovarian failure and polycystic ovary syndrome. *Horm Metab Res* 2001;**4**:221–226.
- Tong ZB, Sullivan SD, Lawless LM, Vanderhoof V, Zachman K, Nelson LM. Five mutations of mitochondrial DNA polymerase-gamma (POLG) are not a prevalent etiology for spontaneous 46,XX primary ovarian insufficiency. *Fertil Steril* 2010;**7**:2932–2934.
- Tosh D, Rani HS, Murty US, Deenadayal A, Grover P. Mutational analysis of the FIGLA gene in women with idiopathic premature ovarian failure. *Menopause* 2015;**22**:520–526.
- Tsuda M, Sasaoka Y, Kiso M, Abe K, Haraguchi S, Kobayashi S, Saga Y. Conserved role of nanos proteins in germ cell development. *Science* 2003;**5637**:1239–1241.
- Uno S, Zembutsu H, Hirasawa A, Takahashi A, Kubo M, Akahane T, Aoki D, Kamatani N, Hirata K, Nakamura Y. A genome-wide association study identifies genetic variants in the CDKN2BAS locus associated with endometriosis in Japanese. *Nat Genet* 2010;**8**:707–710.
- Van Kasteren YM, Hundscheid RD, Smits AP, Cremers FP, van Zonneveld P, Braat DD. Familial idiopathic premature ovarian failure: an overrated and underestimated genetic disease? *Hum Reprod* 1999;**10**:2455–2459.

- Villaescusa JC, Allard P, Carminati E, Kontogiannea M, Talarico D, Blasi F, Farookhi R, Verrotti AC. Clast4, the murine homologue of human elF4E-Transporter, is highly expressed in developing oocytes and post-translationally modified at meiotic maturation. *Gene* 2006;**367**:101–109.
- Vilodre LC, Kohek MB, Spritzer PM. Screening of follicle-stimulating hormone receptor gene in women with premature ovarian failure in southern Brazil and associations with phenotype. *J Endocrinol Invest* 2008;**6**:552–557.
- Waggoner DD, Buist NR, Donnell GN. Long-term prognosis in galactosaemia: results of a survey of 350 cases. *J Inherit Metab Dis* 1990;**6**:802–818.
- Wang CY, Davoodi-Semiromi A, Huang W, Connor E, Shi JD, She JX. Characterization of mutations in patients with autoimmune polyglandular syndrome type I (APS1). *Hum Genet* 1998;**6**:681–685.
- Wang LL, Levy ML, Lewis RA, Chintagumpala MM, Lev D, Rogers M, Plon SE. Clinical manifestations in a cohort of 41 Rothmund-Thomson syndrome patients. *Am J Med Genet* 2001;**1**:11–17.
- Wang HQ, Takakura K, Takebayashi K, Noda Y. Mutational analysis of the mullerian-inhibiting substance gene and its receptor gene in Japanese women with polycystic ovary syndrome and premature ovarian failure. *Fertil Steril* 2002;**6**:1329–1330.
- Wang B, Li L, Ni F, Song J, Wang J, Mu Y, Ma X, Cao Y. Mutational analysis of SAL-Like 4 (SALL4) in Han Chinese women with premature ovarian failure. *Mol Hum Reprod* 2009;**9**:557–562.
- Wang B, Ni F, Li L, Wei Z, Zhu X, Wang J, Cao Y, Ma X. Analysis of cyclin-dependent kinase inhibitor 1B mutation in Han Chinese women with premature ovarian failure. *Reprod Biomed Online* 2010a;**2**:212–214.
- Wang B, Wen Q, Ni F, Zhou S, Wang J, Cao Y, Ma X. Analyses of growth differentiation factor 9 (GDF9) and bone morphogenetic protein 15 (BMP15) mutation in Chinese women with premature ovarian failure. *Clin Endocrinol (Oxf)* 2010b;**1**:135–136.
- Wang J, Wang B, Song J, Suo P, Ni F, Chen B, Ma X, Cao Y. New candidate gene POU5F1 associated with premature ovarian failure in Chinese patients. *Reprod Biomed Online* 2011;**3**:312–316.
- Wang H, Sun M, Qin Y, Xia T, Ma J, Chen ZJ. Mutations in DMCI are not responsible for premature ovarian failure in Chinese women. *Reprod Biomed Online* 2012;**2**:175–178.
- Wang J, Zhang W, Jiang H, Wu BL. Mutations in HFM1 in recessive primary ovarian insufficiency. *N Engl J Med* 2014a;**10**:972–974.
- Wang JL, Li SL, Qin YY, Chen ZJ. Analysis of progesterone receptor membrane component 1 mutation in Han Chinese women with premature ovarian failure. *Reprod Biomed Online* 2014b;**5**:640–643.
- Watkins WJ, Umbers AJ, Woad KJ, Harris SE, Winship IM, Gersak K, Shelling AN. Mutational screening of FOXO3A and FOXO1A in women with premature ovarian failure. *Fertil Steril* 2006;**5**:1518–1521.
- Wittenberger MD, Hagerman RJ, Sherman SL, McConkie-Rosell A, Welt CK, Rebar RW, Corrigan EC, Simpson JL, Nelson LM. The FMR1 premutation and reproduction. *Fertil Steril* 2007;**3**:456–465.
- Woad KJ, Prendergast D, Winship IM, Shelling AN. FSH receptor gene variants are rarely associated with premature ovarian failure. *Reprod Biomed Online* 2013;**4**:396–399.
- Wood-Trageser MA, Gurbuz F, Yatsenko SA, Jeffries EP, Kotan LD, Surti U, Ketterer DM, Matic J, Chipkin J, Jiang H et al. MCM9 mutations are associated with ovarian failure, short stature, and chromosomal instability. *Am J Hum Genet* 2014;**6**:754–762.
- Wu X, Wang B, Dong Z, Zhou S, Liu Z, Shi G, Cao Y, Xu Y. A NANOS3 mutation linked to protein degradation causes premature ovarian insufficiency. *Cell Death Dis* 2013;**4**:e825.
- Yan C, Wang P, DeMayo J, DeMayo FJ, Elvin JA, Carino C, Prasad SV, Skinner SS, Dunbar BS, Dube JL et al. Synergistic roles of bone morphogenetic protein 15 and growth differentiation factor 9 in ovarian function. *Mol Endocrinol* 2001;**6**:854–866.
- Yoon SH, Choi YM, Hong MA, Kim JJ, Lee GH, Hwang KR, Moon SY. Association study of anti-Mullerian hormone and anti-Mullerian hormone type II receptor polymorphisms with idiopathic primary ovarian insufficiency. *Hum Reprod* 2013;**12**:3301–3305.
- Zhang J, Tang YY, Guo YP. A study of hypergonadotropic secondary amenorrhea with cytogenetics. *J Chongqing Med Univ* 2003;**28**:151–152.
- Zhang J, Tam WL, Tong GQ, Wu Q, Chan HY, Soh BS, Lou Y, Yang J, Ma Y, Chai L et al. Sall4 modulates embryonic stem cell pluripotency and early embryonic development by the transcriptional regulation of Pou5f1. *Nat Cell Biol* 2006;**10**:1114–1123.
- Zhang P, Shi YH, Wang LC, Chen ZJ. Sequence variants in exons of the BMP-15 gene in Chinese patients with premature ovarian failure. *Acta Obstet Gynecol Scand* 2007;**5**:585–589.
- Zhao H, Qin Y, Kovanci E, Simpson JL, Chen ZJ, Rajkovic A. Analyses of GDF9 mutation in 100 Chinese women with premature ovarian failure. *Fertil Steril* 2007;**5**:1474–1476.
- Zhao H, Chen ZJ, Qin Y, Shi Y, Wang S, Choi Y, Simpson JL, Rajkovic A. Transcription factor FIGLA is mutated in patients with premature ovarian failure. *Am J Hum Genet* 2008;**6**:1342–1348.
- Zhao Z, Qin Y, Ma J, Zhao H, Li J, Wang L, Ren C, Che L, Chen ZJ. PTEN gene analysis in premature ovarian failure patients. *Acta Obstet Gynecol Scand* 2011;**6**:678–679.
- Zhao Z, Wei D, Mu Y, Qin Y, Li G, Cui L, Chen ZJ. Mutational analysis of SKP2 and P27 in Chinese Han women with premature ovarian failure. *Reprod Biomed Online* 2013;**1**:104–106.
- Zhao S, Li G, Dalgleish R, Vujovic S, Jiao X, Li J, Simpson JL, Qin Y, Ivanisevic M, Ivovic M et al. Transcription factor SOHLH1 potentially associated with primary ovarian insufficiency. *Fertil Steril* 2015;**103**:548–553.
- Zhen XM, Sun YM, Qiao J, Li R, Wang LN, Liu P. Genome-wide copy number scan in Chinese patients with premature ovarian failure. *Beijing Da Xue Xue Bao* 2013;**6**:841–847.
- Zhou S, Wang B, Ni F, Wang J, Cao Y, Ma X. GPR3 may not be a potential candidate gene for premature ovarian failure. *Reprod Biomed Online* 2010;**1**:53–55.
- Zlotogora J, Sagi M, Cohen T. The blepharophimosis, ptosis, and epicanthus inversus syndrome: delineation of two types. *Am J Hum Genet* 1983;**5**:1020–1027.